



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 2, 2025 – 12:39 am BST

PDB ID : 1OIZ / pdb\_00001oiz  
Title : The Molecular Basis of Vitamin E Retention: Structure of Human Alpha-Tocopherol Transfer Protein  
Authors : Meier, R.; Tomizaki, T.; Schulze-Briesse, C.; Baumann, U.; Stocker, A.  
Deposited on : 2003-06-27  
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	278	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>14%</div> <div>• •</div> </div> </div>
1	B	278	<div> <div>4%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>• 5%</div> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4733 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-TOCOPHEROL TRANSFER PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	1
			2147	1393	372	376	6			
1	B	265	Total	C	N	O	S	0	0	1
			2142	1390	371	375	6			

- Molecule 2 is FRAGMENT OF TRITON X-100 (CCD ID: TRT) (formula:  $C_{21}H_{36}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			25	21	4		
2	A	1	Total	C	O	0	0
			25	21	4		
2	A	1	Total	C	O	0	0
			25	21	4		
2	B	1	Total	C	O	0	0
			25	21	4		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			25	21	4		
2	B	1	Total	C	O	0	0
			25	21	4		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	146	Total	O	0	0
			146	146		
3	B	148	Total	O	0	0
			148	148		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ALPHA-TOCOPHEROL TRANSFER PROTEIN



#### • Molecule 1: ALPHA-TOCOPHEROL TRANSFER PROTEIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.14Å 113.78Å 67.43Å 90.00° 98.95° 90.00°	Depositor
Resolution (Å)	30.00 – 1.88 30.00 – 1.88	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-1.88) 99.3 (30.00-1.88)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.195 , 0.226 0.196 , 0.225	Depositor DCC
$R_{free}$ test set	2711 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtriage
Anisotropy	0.341	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 46.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4733	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	1/2206 (0.0%)	0.70	3/2994 (0.1%)
1	B	0.48	0/2201	0.69	1/2987 (0.0%)
All	All	0.49	1/4407 (0.0%)	0.69	4/5981 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	90	ARG	CZ-NH1	6.22	1.41	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	ARG	NE-CZ-NH2	-6.89	116.86	120.30
1	A	268	ASP	CB-CG-OD2	5.86	123.57	118.30
1	A	185	ASP	CB-CG-OD2	5.29	123.06	118.30
1	B	185	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	43	ALA	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2147	0	2163	31	0
1	B	2142	0	2158	39	0
2	A	75	0	108	31	0
2	B	75	0	108	28	0
3	A	146	0	0	3	0
3	B	148	0	0	5	0
All	All	4733	0	4537	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:SER:OG	2:A:1275:TRT:H4C3	1.17	1.29
1:B:117:TYR:OH	2:B:1277:TRT:H7C2	1.42	1.20
2:A:1276:TRT:H4C3	2:A:1276:TRT:H8C2	1.39	1.05
2:B:1277:TRT:H10	2:B:1277:TRT:H3C3	1.40	1.03
1:B:195:HIS:HD2	1:B:225:HIS:HE1	1.13	0.95
2:B:1277:TRT:H3C3	2:B:1277:TRT:C10	1.97	0.94
2:B:1277:TRT:H4C1	3:B:2091:HOH:O	1.67	0.94
2:A:1276:TRT:H3C1	1:B:206:VAL:HG22	1.49	0.93
2:A:1276:TRT:H8C2	2:A:1276:TRT:C4	1.97	0.93
1:B:117:TYR:OH	2:B:1277:TRT:C7	2.18	0.91
2:A:1275:TRT:H2C3	2:A:1275:TRT:C10	2.02	0.89
1:B:179:ILE:HD13	2:B:1276:TRT:H201	1.53	0.88
1:B:42:LEU:HG	1:B:43:ALA:H	1.41	0.85
2:A:1276:TRT:H4C3	2:A:1276:TRT:C8	2.08	0.83
1:A:140:SER:HG	2:A:1275:TRT:C4	1.91	0.81
1:B:195:HIS:CD2	1:B:225:HIS:HE1	1.99	0.80
1:B:195:HIS:HD2	1:B:225:HIS:CE1	2.00	0.79

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1274:TRT:H191	2:A:1276:TRT:H7C2	1.66	0.77
2:A:1276:TRT:H2C1	1:B:206:VAL:HG13	1.66	0.75
1:A:115:LEU:HD21	2:A:1275:TRT:H3C1	1.67	0.74
1:A:163:TRP:HE1	2:A:1274:TRT:H221	1.50	0.73
2:B:1277:TRT:H2C2	2:B:1277:TRT:C9	2.21	0.71
2:A:1274:TRT:C19	2:A:1276:TRT:H7C2	2.22	0.70
2:A:1274:TRT:H14	2:A:1276:TRT:H161	1.75	0.67
1:A:241:LEU:O	1:A:247:GLY:HA3	1.97	0.65
1:B:227:ASN:HB3	3:B:2106:HOH:O	1.98	0.64
1:A:115:LEU:CD2	2:A:1275:TRT:H3C1	2.27	0.63
1:B:101:HIS:HE1	1:B:139:THR:OG1	1.80	0.62
2:A:1275:TRT:C10	2:A:1275:TRT:C2	2.79	0.61
1:B:115:LEU:CD2	2:B:1277:TRT:H7C1	2.30	0.61
1:B:93:ILE:HG13	1:B:97:LYS:HE3	1.84	0.60
2:B:1277:TRT:H10	2:B:1277:TRT:C3	2.26	0.60
1:B:264:MET:HA	1:B:264:MET:HE2	1.83	0.59
1:A:115:LEU:CD2	2:A:1275:TRT:H2C1	2.34	0.58
2:A:1276:TRT:H14	2:A:1276:TRT:H2C3	1.87	0.57
1:A:199:GLU:OE2	1:A:204:HIS:NE2	2.35	0.57
1:B:132:VAL:HG23	2:B:1277:TRT:H191	1.85	0.57
1:B:119:ILE:HG13	1:B:160:LEU:HD22	1.86	0.57
1:A:13:GLN:HB2	3:A:2002:HOH:O	2.05	0.56
1:A:76:TRP:CH2	1:A:137:LEU:HD21	2.39	0.56
2:A:1274:TRT:H7C3	2:A:1274:TRT:H3C3	1.88	0.56
1:B:28:LEU:HD21	1:B:53:LEU:HG	1.88	0.55
1:B:264:MET:HA	1:B:264:MET:CE	2.37	0.55
1:B:119:ILE:HD11	2:B:1277:TRT:H252	1.89	0.55
1:A:115:LEU:HD22	2:A:1275:TRT:H2C1	1.88	0.53
1:A:115:LEU:HD22	2:A:1275:TRT:C2	2.39	0.52
1:B:179:ILE:HG12	2:B:1277:TRT:H162	1.91	0.51
1:B:42:LEU:HG	1:B:43:ALA:N	2.19	0.51
1:B:37:GLU:C	1:B:39:GLY:H	2.14	0.51
1:A:160:LEU:HD13	1:A:163:TRP:CE3	2.46	0.50
2:B:1276:TRT:H4C2	2:B:1276:TRT:C8	2.41	0.50
2:A:1276:TRT:H4C1	1:B:206:VAL:HA	1.93	0.50
1:A:223:HIS:HE1	3:A:2098:HOH:O	1.94	0.50
1:A:28:LEU:HD21	1:A:53:LEU:HG	1.95	0.49
1:A:57:ARG:HH11	1:A:150:GLN:NE2	2.10	0.49
1:B:154:ILE:CD1	2:B:1277:TRT:H5C1	2.42	0.49
2:B:1275:TRT:H2C3	2:B:1276:TRT:H191	1.95	0.49
2:B:1277:TRT:C10	2:B:1277:TRT:C3	2.82	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:210:ILE:HD12	1:B:214:LEU:HD21	1.96	0.48
1:A:189:LEU:HD13	2:A:1275:TRT:H3C2	1.95	0.48
1:B:195:HIS:HE1	3:B:2104:HOH:O	1.96	0.48
1:B:223:HIS:HE1	3:B:2102:HOH:O	1.95	0.48
1:B:180:ALA:O	1:B:184:THR:HG23	2.14	0.47
1:B:166:SER:HB3	3:B:2084:HOH:O	2.15	0.47
1:A:210:ILE:HG23	2:B:1276:TRT:H10	1.97	0.47
1:A:203:PHE:HA	2:A:1274:TRT:H172	1.97	0.46
1:B:12:PRO:O	1:B:15:ASN:HB2	2.16	0.46
2:B:1277:TRT:C9	2:B:1277:TRT:C2	2.93	0.46
2:B:1276:TRT:H4C2	2:B:1276:TRT:H8C1	1.98	0.46
2:A:1275:TRT:H2C3	2:A:1275:TRT:C9	2.44	0.46
1:A:164:GLN:HG3	1:A:166:SER:OG	2.15	0.46
2:A:1274:TRT:H192	2:A:1276:TRT:H7C2	1.98	0.46
1:B:33:ARG:HG2	1:B:34:ARG:N	2.30	0.45
1:B:260:THR:O	1:B:264:MET:HG2	2.16	0.45
1:A:210:ILE:CG2	2:B:1276:TRT:H10	2.46	0.45
2:A:1276:TRT:C3	1:B:206:VAL:HG22	2.34	0.45
1:A:119:ILE:HD13	1:A:160:LEU:HD22	1.98	0.45
1:A:187:PHE:CG	1:A:188:PRO:HD2	2.52	0.44
2:A:1275:TRT:C4	2:A:1275:TRT:H8C2	2.48	0.44
1:B:154:ILE:HD11	2:B:1277:TRT:H4C2	2.00	0.44
2:B:1276:TRT:H2C3	2:B:1276:TRT:H14	1.99	0.44
1:A:45:LEU:HD12	1:A:45:LEU:HA	1.91	0.43
2:A:1274:TRT:H171	2:A:1276:TRT:H7C3	2.00	0.43
1:A:57:ARG:HH11	1:A:150:GLN:HE22	1.66	0.43
1:B:115:LEU:HD23	2:B:1277:TRT:H7C1	1.99	0.43
2:A:1274:TRT:H2C2	2:A:1274:TRT:C9	2.48	0.43
1:A:273:ILE:HG22	1:A:274:SER:N	2.33	0.42
2:A:1276:TRT:H2C2	2:B:1276:TRT:C7	2.49	0.42
2:B:1276:TRT:H10	2:B:1276:TRT:H7C3	1.84	0.42
2:B:1275:TRT:H4C2	2:B:1276:TRT:H191	2.01	0.42
2:A:1274:TRT:H162	2:A:1276:TRT:H7C3	2.01	0.42
1:B:142:LEU:CD1	1:B:263:ILE:HD13	2.50	0.42
2:B:1275:TRT:C9	2:B:1275:TRT:H2C2	2.49	0.42
1:A:43:ALA:O	1:A:45:LEU:N	2.53	0.41
1:B:136:SER:HB3	2:B:1277:TRT:C10	2.50	0.41
1:A:234:LEU:HD22	1:A:241:LEU:HD12	2.02	0.41
1:A:77:ARG:HH21	1:A:273:ILE:CG2	2.33	0.41
1:A:201:VAL:HG23	3:A:2103:HOH:O	2.20	0.41
1:B:24:LEU:HD22	1:B:61:PHE:CZ	2.56	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225:HIS:HD2	1:B:232:SER:OG	2.03	0.40
1:A:108:ASP:HB2	1:A:109:PRO:CD	2.51	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/278 (95%)	254 (96%)	7 (3%)	3 (1%)	12	3
1	B	263/278 (95%)	253 (96%)	9 (3%)	1 (0%)	30	18
All	All	527/556 (95%)	507 (96%)	16 (3%)	4 (1%)	16	6

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PRO
1	B	42	LEU
1	A	272	SER
1	A	273	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	232/243 (96%)	221 (95%)	11 (5%)	22 8
1	B	232/243 (96%)	225 (97%)	7 (3%)	36 19
All	All	464/486 (96%)	446 (96%)	18 (4%)	27 11

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG
1	A	37	GLU
1	A	40	VAL
1	A	42	LEU
1	A	45	LEU
1	A	133	PHE
1	A	137	LEU
1	A	160	LEU
1	A	202	ILE
1	A	234	LEU
1	A	271	SER
1	B	23	LEU
1	B	33	ARG
1	B	47	LEU
1	B	133	PHE
1	B	160	LEU
1	B	217	LYS
1	B	266	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	150	GLN
1	A	170	GLN
1	A	198	ASN
1	A	223	HIS
1	A	261	ASN
1	B	101	HIS
1	B	150	GLN
1	B	195	HIS
1	B	198	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	223	HIS
1	B	225	HIS
1	B	228	ASN
1	B	261	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	TRT	B	1276	-	25,25,25	0.50	0	33,33,33	0.58	0
2	TRT	A	1275	1	25,25,25	0.58	0	33,33,33	0.72	1 (3%)
2	TRT	B	1277	-	25,25,25	0.61	0	33,33,33	0.70	0
2	TRT	A	1276	-	25,25,25	0.59	0	33,33,33	0.82	1 (3%)
2	TRT	A	1274	-	25,25,25	0.53	0	33,33,33	0.57	0
2	TRT	B	1275	-	25,25,25	0.55	0	33,33,33	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRT	B	1276	-	-	8/23/23/23	0/1/1/1
2	TRT	A	1275	1	-	5/23/23/23	0/1/1/1
2	TRT	B	1277	-	-	13/23/23/23	0/1/1/1
2	TRT	A	1276	-	-	13/23/23/23	0/1/1/1
2	TRT	A	1274	-	-	3/23/23/23	0/1/1/1
2	TRT	B	1275	-	-	6/23/23/23	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1275	TRT	C16-O15-C12	2.49	124.44	117.93
2	A	1276	TRT	C16-O15-C12	2.01	123.18	117.93

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1276	TRT	C1-C5-C6-C9
2	B	1276	TRT	C1-C5-C6-C8
2	A	1276	TRT	C4-C1-C5-C6
2	A	1276	TRT	C3-C1-C5-C6
2	A	1276	TRT	O15-C16-C17-O18
2	B	1276	TRT	O15-C16-C17-O18
2	A	1275	TRT	O21-C22-C23-O24
2	A	1276	TRT	O21-C22-C23-O24
2	A	1276	TRT	C2-C1-C5-C6
2	A	1276	TRT	C13-C12-O15-C16
2	B	1277	TRT	O15-C16-C17-O18
2	A	1276	TRT	C11-C12-O15-C16
2	B	1275	TRT	O18-C19-C20-O21
2	A	1274	TRT	O18-C19-C20-O21
2	A	1275	TRT	O18-C19-C20-O21
2	A	1276	TRT	C1-C5-C6-C8
2	B	1277	TRT	C13-C12-O15-C16
2	B	1277	TRT	C17-C16-O15-C12
2	A	1276	TRT	C5-C6-C9-C14
2	A	1276	TRT	C5-C6-C9-C10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	1276	TRT	C1-C5-C6-C7
2	B	1277	TRT	C1-C5-C6-C7
2	B	1277	TRT	C11-C12-O15-C16
2	A	1274	TRT	O15-C16-C17-O18
2	B	1277	TRT	C1-C5-C6-C8
2	B	1277	TRT	C1-C5-C6-C9
2	A	1276	TRT	C1-C5-C6-C7
2	A	1276	TRT	O18-C19-C20-O21
2	B	1277	TRT	C4-C1-C5-C6
2	B	1276	TRT	O18-C19-C20-O21
2	B	1275	TRT	C17-C16-O15-C12
2	A	1275	TRT	O15-C16-C17-O18
2	A	1275	TRT	C16-C17-O18-C19
2	B	1276	TRT	O21-C22-C23-O24
2	B	1275	TRT	C23-C22-O21-C20
2	B	1275	TRT	C19-C20-O21-C22
2	A	1274	TRT	C22-C23-O24-C25
2	A	1276	TRT	C1-C5-C6-C9
2	B	1277	TRT	C19-C20-O21-C22
2	B	1276	TRT	C20-C19-O18-C17
2	B	1277	TRT	C20-C19-O18-C17
2	A	1275	TRT	C22-C23-O24-C25
2	B	1277	TRT	C2-C1-C5-C6
2	B	1275	TRT	C22-C23-O24-C25
2	B	1275	TRT	O15-C16-C17-O18
2	B	1277	TRT	C22-C23-O24-C25
2	B	1276	TRT	C22-C23-O24-C25
2	B	1277	TRT	C3-C1-C5-C6


There are no ring outliers.

6 monomers are involved in 58 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1276	TRT	10	0
2	A	1275	TRT	12	0
2	B	1277	TRT	17	0
2	A	1276	TRT	15	0
2	A	1274	TRT	10	0
2	B	1275	TRT	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/278 (95%)	0.25	16 (6%) 29 29	13, 22, 35, 42	0
1	B	265/278 (95%)	0.19	11 (4%) 41 42	13, 22, 34, 42	0
All	All	531/556 (95%)	0.22	27 (5%) 34 35	13, 22, 34, 42	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	44	PRO	6.2
1	B	42	LEU	6.2
1	B	43	ALA	5.8
1	A	43	ALA	5.0
1	B	11	GLY	4.6
1	B	45	LEU	3.5
1	A	41	PRO	3.4
1	B	44	PRO	3.4
1	A	273	ILE	3.4
1	B	41	PRO	3.3
1	A	130	TYR	3.0
1	A	35	ALA	2.9
1	A	45	LEU	2.9
1	A	274	SER	2.6
1	B	33	ARG	2.6
1	B	38	ALA	2.6
1	A	9	SER	2.6
1	A	80	CYS	2.6
1	A	46	PRO	2.6
1	B	271	SER	2.6
1	B	46	PRO	2.5
1	A	36	ARG	2.4
1	A	250	PHE	2.4
1	A	42	LEU	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	165	PHE	2.2
1	A	38	ALA	2.2
1	A	33	ARG	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	TRT	A	1276	25/25	0.62	0.24	60,63,66,66	0
2	TRT	A	1275	25/25	0.64	0.28	53,60,61,61	0
2	TRT	B	1276	25/25	0.73	0.22	62,63,64,64	0
2	TRT	B	1275	25/25	0.75	0.17	50,53,68,69	0
2	TRT	B	1277	25/25	0.75	0.23	53,59,65,66	0
2	TRT	A	1274	25/25	0.79	0.16	45,47,63,64	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around TRT A 1276:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around TRT A 1275:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around TRT B 1276:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TRT B 1275:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around TRT B 1277:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around TRT A 1274:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.