



wwPDB NMR Structure Validation Summary Report ⓘ

Apr 1, 2025 – 09:59 pm BST

PDB ID : 1OLN / pdb_00001ohn
Title : Model for thiostrepton antibiotic binding to L11 substrate from 50S ribosomal RNA
Authors : Lentzen, G.; Klinck, R.; Matassova, N.; Aboul-Ela, F.; Murchie, A.I.H.
Deposited on : 2003-08-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR, THEORETICAL MODEL

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463
RNA backbone	6643	756

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	140	<div> <div>41%</div> <div>41%</div> <div>11%</div> <div>• 5%</div> </div>
2	B	19	<div> <div>32%</div> <div>47%</div> <div>21%</div> </div>
3	C	58	<div> <div>62%</div> <div>38%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
2	B	BB9	13	-	1

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2350 atoms, of which 0 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called 50S RIBOSOMAL PROTEIN L11.

Mol	Chain	Residues	Atoms					Trace
1	A	133	Total	C	N	O	S	0
			999	642	169	182	6	

- Molecule 2 is a protein called THIOSTREPTON.

Mol	Chain	Residues	Atoms					Trace
2	B	19	Total	C	N	O	S	1
			114	72	19	18	5	

- Molecule 3 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					Trace
3	C	58	Total	C	N	O	P	0
			1237	554	225	401	57	

• Molecule 1: 50S RIBOSOMAL PROTEIN L11



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *DOCKING/MODELING*.

Of the ? calculated structures, 1 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATION AND BEST OVERALL DOCKING SCORE*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
rDOCK	refinement	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MH6, TS9, NH2, DBU, QUA, DCY, BB9, DHA

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
3	C	0	1
All	All	0	1

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

All planar outliers are listed below.

Mol	Chain	Res	Type	Group
3	C	1069	A	Sidechain

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	999	0	1071	61
2	B	114	0	79	11
3	C	1237	0	627	12
All	All	2350	0	1777	75

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

5 of 75 clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:22:PRO:HG3	2:B:11:BB9:SG	1.34	1.62
1:A:22:PRO:CG	2:B:11:BB9:SG	0.92	2.57
1:A:14:PRO:HG2	1:A:17:LYS:HG3	0.72	1.59
1:A:20:PRO:HB2	1:A:23:PRO:HD2	0.72	1.61
1:A:101:THR:HA	1:A:140:VAL:O	0.71	1.86

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/140 (94%)	112 (85%)	11 (8%)	8 (6%)	2	19
2	B	5/19 (26%)	4 (80%)	1 (20%)	0 (0%)	100	100
All	All	136/159 (86%)	116 (85%)	12 (9%)	8 (6%)	2	20

5 of 8 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	15	ALA
1	A	19	THR
1	A	23	PRO
1	A	24	VAL
1	A	25	GLY

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/112 (96%)	82 (76%)	26 (24%)	2	25
2	B	3/4 (75%)	2 (67%)	1 (33%)	1	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	111/116 (96%)	84 (76%)	27 (24%)	2 24

5 of 27 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	11	LEU
1	A	13	LEU
1	A	17	LYS
1	A	24	VAL
1	A	28	LEU

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
3	C	57/58 (98%)	3 (5%)	0 (0%)	0.67
All	All	57/58 (98%)	3 (5%)	0 (0%)	0.67

The overall RNA backbone suiteness is 0.67.

All RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	C	1070	A
3	C	1088	A
3	C	1090	U

There are no RNA pucker outliers to report.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

11 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	BB9	B	11	2	3,5,6	1.36	0 (0%)
2	DHA	B	16	2	4,4,5	4.20	2 (50%)
2	BB9	B	15	2	3,5,6	3.63	1 (33%)
2	MH6	B	14	2	3,3,6	1.55	1 (33%)
2	TS9	B	10	2	6,8,10	0.90	0 (0%)
2	DHA	B	17	2	4,4,5	1.79	1 (25%)
2	BB9	B	13	2	2,4,6	1.66	1 (50%)
2	BB9	B	6	2	3,5,6	1.82	1 (33%)
2	DHA	B	3	2	4,4,5	1.99	2 (50%)
2	DBU	B	8	2	4,4,6	3.13	2 (50%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	BB9	B	11	2	1,5,7	2.59	1 (100%)
2	DHA	B	16	2	2,4,6	5.47	1 (50%)
2	BB9	B	15	2	1,5,7	4.09	1 (100%)
2	MH6	B	14	2	1,3,7	0.45	0 (0%)
2	TS9	B	10	2	5,12,15	1.16	0 (0%)
2	DHA	B	17	2	2,4,6	2.59	1 (50%)
2	BB9	B	13	2	3,4,7	2.66	3 (100%)
2	BB9	B	6	2	1,5,7	2.63	1 (100%)
2	DHA	B	3	2	2,4,6	1.79	1 (50%)
2	DBU	B	8	2	4,4,7	1.51	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BB9	B	11	2	-	0,0,4,6	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DHA	B	16	2	-	0,0,2,4	-
2	BB9	B	15	2	-	0,0,4,6	-
2	TS9	B	10	2	-	0,9,12,16	-
2	DHA	B	17	2	-	0,0,2,4	-
2	BB9	B	13	2	-	0,0,2,6	-
2	BB9	B	6	2	-	0,0,4,6	-
2	DHA	B	3	2	-	0,0,2,4	-
2	DBU	B	8	2	-	0,1,2,6	-

5 of 11 bond outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	16	DHA	CA-N	7.90	1.55	1.35
2	B	15	BB9	O-C	6.24	1.07	1.22
2	B	8	DBU	CA-N	5.78	1.47	1.33
2	B	17	DHA	CA-N	3.05	1.42	1.35
2	B	6	BB9	O-C	2.90	1.28	1.22

5 of 10 angle outliers are listed below. They are sorted according to the Z-score.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	16	DHA	O-C-CA	7.70	111.18	125.54
2	B	15	BB9	O-C-CA	4.09	130.58	125.39
2	B	17	DHA	O-C-CA	3.08	119.81	125.54
2	B	13	BB9	C-CA-CB	2.91	126.69	121.39
2	B	8	DBU	CB-CA-N	2.77	124.55	122.87

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided