



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2025 – 04:12 PM EDT

PDB ID : 9ONO / pdb_00009ono
Title : Fe-bound B. pseudomallei rubrerythrin
Authors : Budziszewski, G.R.; Snell, M.E.; Monteiro, D.C.F.; Lynch, M.L.; Bowman, S.E.J.
Deposited on : 2025-05-15
Resolution : 1.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0rc1
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	<div><div></div><div>94%5%•</div></div>
1	B	140	<div><div>%</div><div>94%5%•</div></div>
1	C	140	<div><div>%</div><div>94%5%•</div></div>
1	D	140	<div><div>%</div><div>94%6%•</div></div>
1	E	140	<div><div></div><div>90%9%•</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	140	<div><div>%</div><div><div></div><div>96%</div><div></div></div><div><div></div><div></div><div></div></div></div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13927 atoms, of which 6193 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rubrerythrin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	139	Total	C	H	N	O	S	0	5	0
			2162	691	1046	195	228	2			
1	B	139	Total	C	H	N	O	S	0	2	0
			2106	674	1016	191	223	2			
1	C	139	Total	C	H	N	O	S	0	4	0
			2143	685	1035	194	227	2			
1	D	139	Total	C	H	N	O	S	0	6	0
			2176	695	1052	197	230	2			
1	E	139	Total	C	H	N	O	S	0	3	0
			2129	682	1028	192	225	2			
1	F	139	Total	C	H	N	O	S	0	2	0
			2106	674	1016	191	223	2			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Fe	0	0
			2	2		
2	B	2	Total	Fe	0	0
			2	2		
2	C	2	Total	Fe	0	0
			2	2		
2	D	2	Total	Fe	0	0
			2	2		
2	E	2	Total	Fe	0	0
			2	2		
2	F	2	Total	Fe	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	193	Total 195	O 195	0	2
3	B	185	Total 185	O 185	0	0
3	C	189	Total 189	O 189	0	0
3	D	169	Total 170	O 170	0	1
3	E	173	Total 174	O 174	0	1
3	F	179	Total 180	O 180	0	1

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rubrerythrin

Chain A: 



- Molecule 1: Rubrerythrin

Chain B: 




- Molecule 1: Rubrerythrin

Chain C: 




- Molecule 1: Rubrerythrin

Chain D: 



- Molecule 1: Rubrerythrin

Chain E: 



- Molecule 1: Rubrerythrin



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	202.79Å 202.79Å 68.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.32 – 1.66 38.32 – 1.66	Depositor EDS
% Data completeness (in resolution range)	97.7 (38.32-1.66) 97.6 (38.32-1.66)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.04 (at 1.66Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, R_{free}	0.172 , 0.206 0.172 , 0.206	Depositor DCC
R_{free} test set	6275 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.018 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13927	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/1138	0.52	0/1533
1	B	0.39	0/1112	0.55	0/1499
1	C	0.38	0/1130	0.53	0/1522
1	D	0.40	0/1146	0.50	0/1544
1	E	0.37	0/1123	0.49	0/1514
1	F	0.37	0/1112	0.52	0/1499
All	All	0.38	0/6761	0.52	0/9111

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	47	ARG	Sidechain
1	B	125	ARG	Sidechain
1	C	47	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1116	1046	1041	3	0
1	B	1090	1016	1014	3	0
1	C	1108	1035	1031	5	0
1	D	1124	1052	1046	6	0
1	E	1101	1028	1025	9	0
1	F	1090	1016	1014	3	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	195	0	0	0	0
3	B	185	0	0	0	0
3	C	189	0	0	0	0
3	D	170	0	0	1	0
3	E	174	0	0	1	0
3	F	180	0	0	0	0
All	All	7734	6193	6171	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42[B]:ILE:CD1	1:B:138:LEU:HD13	2.27	0.65
1:E:97:MET:O	1:E:101:MET:HG3	2.13	0.49
1:C:120:LEU:HD22	1:D:56:HIS:CD2	2.48	0.48
1:E:61[B]:LEU:HD23	1:F:29:TYR:CG	2.50	0.47
1:F:74:LEU:HD12	1:F:85:SER:HA	1.97	0.47
1:A:91:THR:HA	1:A:94:TYR:CE2	2.52	0.45
1:E:56:HIS:CD2	1:F:120:LEU:HD22	2.52	0.44
1:E:17:PHE:CE1	1:E:61[A]:LEU:HG	2.52	0.44
1:E:91:THR:HA	1:E:94:TYR:CE2	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5[A]:LYS:HE3	1:A:67:VAL:HG12	2.01	0.43
1:E:91:THR:O	1:E:95:THR:HG23	2.19	0.42
1:D:91:THR:HA	1:D:94:TYR:CE2	2.54	0.42
1:D:120[B]:LEU:O	1:D:124:GLU:HG2	2.19	0.42
1:B:120:LEU:O	1:B:124:GLU:HG2	2.20	0.41
1:B:65:GLU:HG2	1:B:76:PHE:CE2	2.56	0.41
1:C:56:HIS:CD2	1:D:120[A]:LEU:HD22	2.56	0.41
1:C:120:LEU:O	1:C:124:GLU:HG2	2.21	0.41
1:C:116:TRP:HE1	1:D:60:HIS:HD1	1.69	0.40
1:E:122:LYS:HE2	3:E:400:HOH:O	2.21	0.40
1:C:52:GLY:HA2	1:E:35:ASP:HB3	2.04	0.40
1:D:2:ALA:N	3:D:305:HOH:O	2.54	0.40
1:E:98:TYR:N	1:E:99:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	142/140 (101%)	140 (99%)	2 (1%)	0	100	100
1	B	139/140 (99%)	137 (99%)	2 (1%)	0	100	100
1	C	141/140 (101%)	139 (99%)	2 (1%)	0	100	100
1	D	143/140 (102%)	140 (98%)	3 (2%)	0	100	100
1	E	140/140 (100%)	136 (97%)	4 (3%)	0	100	100
1	F	139/140 (99%)	136 (98%)	3 (2%)	0	100	100
All	All	844/840 (100%)	828 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/106 (104%)	109 (99%)	1 (1%)	75	63
1	B	107/106 (101%)	106 (99%)	1 (1%)	75	63
1	C	109/106 (103%)	108 (99%)	1 (1%)	75	63
1	D	111/106 (105%)	110 (99%)	1 (1%)	75	63
1	E	108/106 (102%)	107 (99%)	1 (1%)	75	63
1	F	107/106 (101%)	106 (99%)	1 (1%)	75	63
All	All	652/636 (102%)	646 (99%)	6 (1%)	73	63

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25	ARG
1	B	25	ARG
1	C	25	ARG
1	D	25	ARG
1	E	25	ARG
1	F	25	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	3	GLN
1	E	92	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	139/140 (99%)	-0.68	0 100 100	10, 24, 40, 71	5 (3%)
1	B	139/140 (99%)	-0.65	1 (0%) 84 87	11, 25, 48, 73	2 (1%)
1	C	139/140 (99%)	-0.61	1 (0%) 84 87	11, 26, 42, 76	4 (2%)
1	D	139/140 (99%)	-0.48	1 (0%) 84 87	11, 28, 47, 60	6 (4%)
1	E	139/140 (99%)	-0.44	0 100 100	11, 29, 49, 59	3 (2%)
1	F	139/140 (99%)	-0.50	1 (0%) 84 87	12, 28, 45, 73	2 (1%)
All	All	834/840 (99%)	-0.56	4 (0%) 87 90	10, 27, 46, 76	22 (2%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	ALA	2.8
1	B	2	ALA	2.4
1	C	2	ALA	2.2
1	F	2	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FE	A	202	1/1	0.96	0.05	25,25,25,25	1
2	FE	D	202	1/1	0.98	0.06	27,27,27,27	1
2	FE	B	202	1/1	0.99	0.06	25,25,25,25	1
2	FE	C	202	1/1	0.99	0.04	26,26,26,26	1
2	FE	D	201	1/1	0.99	0.02	23,23,23,23	1
2	FE	B	201	1/1	0.99	0.04	25,25,25,25	1
2	FE	E	201	1/1	0.99	0.02	24,24,24,24	1
2	FE	E	202	1/1	0.99	0.03	29,29,29,29	1
2	FE	F	201	1/1	0.99	0.03	22,22,22,22	1
2	FE	A	201	1/1	1.00	0.05	23,23,23,23	1
2	FE	C	201	1/1	1.00	0.02	23,23,23,23	1
2	FE	F	202	1/1	1.00	0.02	25,25,25,25	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FE A 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE D 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE C 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE D 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE E 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE E 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE F 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE A 201:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.