



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 4, 2025 – 12:07 PM EDT

PDB ID : 9OPF / pdb_00009opf
Title : Context-Dependent Variability Of HIF Heterodimers Influences Interactions
With Macromolecular And Small Molecule Partners
Authors : Isiorho, E.A.; Xu, X.; Gardner, K.H.
Deposited on : 2025-05-19
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	8487 (2.30-2.26)
Clashscore	180529	9437 (2.30-2.26)
Ramachandran outliers	177936	9341 (2.30-2.26)
Sidechain outliers	177891	9342 (2.30-2.26)
RSRZ outliers	164620	8487 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	81	
1	B	81	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1837 atoms, of which 935 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transforming acidic coiled-coil-containing protein 3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	55	Total	C	H	N	O	S	0	0	0
			900	267	461	80	89	3			
1	B	57	Total	C	H	N	O	S	0	0	0
			931	277	474	82	95	3			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	O	0	0
			2	2		
2	B	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transforming acidic coiled-coil-containing protein 3



- Molecule 1: Transforming acidic coiled-coil-containing protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	263.11Å 23.83Å 31.85Å 90.00° 91.82° 90.00°	Depositor
Resolution (Å)	29.02 – 2.28 29.02 – 2.28	Depositor EDS
% Data completeness (in resolution range)	98.4 (29.02-2.28) 93.0 (29.02-2.28)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	-0.37 (at 2.26Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419-000	Depositor
R, R_{free}	0.266 , 0.295 0.266 , 0.296	Depositor DCC
R_{free} test set	8354 reflections (9.98%)	wwPDB-VP
Wilson B-factor (Å ²)	46.3	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.065 for -h,-k,l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	1837	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/438	1.01	0/582
1	B	0.65	0/456	0.91	0/606
All	All	0.67	0/894	0.96	0/1188

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	439	461	461	26	0
1	B	457	474	473	13	0
2	A	2	0	0	0	0
2	B	4	0	0	0	0
All	All	902	935	934	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:790:GLN:O	1:A:794:LEU:HD23	1.80	0.81

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:783:ALA:O	1:A:786:ARG:HD2	1.83	0.78
1:A:835:MET:CE	1:B:835:MET:HE3	2.26	0.65
1:B:797:GLN:HG3	1:B:798:ALA:N	2.12	0.65
1:A:797:GLN:O	1:A:797:GLN:OE1	2.16	0.64
1:A:835:MET:HE3	1:B:835:MET:HE3	1.84	0.58
1:A:834:LYS:O	1:A:835:MET:SD	2.64	0.56
1:B:783:ALA:HA	1:B:786:ARG:HB3	1.93	0.51
1:A:786:ARG:HG3	1:A:787:SER:H	1.76	0.50
1:A:834:LYS:C	1:A:835:MET:SD	2.96	0.48
1:A:821:ASN:OD1	1:B:821:ASN:HB2	2.13	0.47
1:B:790:GLN:HG3	1:B:791:ALA:N	2.28	0.47
1:A:787:SER:O	1:A:788:LYS:C	2.57	0.47
1:A:785:VAL:CG1	1:B:785:VAL:HG12	2.46	0.46
1:A:790:GLN:OE1	1:A:794:LEU:HD21	2.16	0.45
1:A:803:GLU:HA	1:A:803:GLU:OE1	2.16	0.45
1:A:789:ALA:O	1:A:792:GLU:HG3	2.17	0.45
1:A:788:LYS:O	1:A:792:GLU:HG3	2.18	0.44
1:B:787:SER:O	1:B:788:LYS:C	2.59	0.44
1:A:786:ARG:CG	1:A:787:SER:H	2.30	0.44
1:B:797:GLN:CG	1:B:798:ALA:N	2.80	0.44
1:A:782:ILE:HG22	1:A:783:ALA:N	2.33	0.43
1:A:790:GLN:O	1:A:794:LEU:CD2	2.60	0.43
1:B:788:LYS:HA	1:B:788:LYS:HD3	1.86	0.42
1:A:782:ILE:CG2	1:A:783:ALA:N	2.81	0.42
1:A:816:GLN:O	1:A:820:GLU:HG3	2.19	0.42
1:A:835:MET:HA	1:A:836:GLU:HA	1.79	0.42
1:A:785:VAL:O	1:A:786:ARG:C	2.62	0.42
1:B:816:GLN:O	1:B:820:GLU:HG3	2.20	0.42
1:A:786:ARG:HG3	1:A:787:SER:N	2.35	0.41
1:A:784:GLN:O	1:A:786:ARG:HG2	2.20	0.41
1:A:785:VAL:CG1	1:B:785:VAL:CG1	2.99	0.40
1:A:785:VAL:HG11	1:B:785:VAL:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	53/81 (65%)	46 (87%)	5 (9%)	2 (4%)	2	1
1	B	55/81 (68%)	51 (93%)	3 (6%)	1 (2%)	7	5
All	All	108/162 (67%)	97 (90%)	8 (7%)	3 (3%)	4	2

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	785	VAL
1	A	783	ALA
1	B	782	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	49/70 (70%)	48 (98%)	1 (2%)	50	65
1	B	51/70 (73%)	51 (100%)	0	100	100
All	All	100/140 (71%)	99 (99%)	1 (1%)	73	83

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	803	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	797	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	55/81 (67%)	1.99	17 (30%) 1 1	47, 74, 148, 155	0
1	B	57/81 (70%)	1.79	21 (36%) 1 1	48, 80, 150, 198	0
All	All	112/162 (69%)	1.89	38 (33%) 1 1	47, 76, 150, 198	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	785	VAL	7.5
1	A	783	ALA	7.1
1	A	792	GLU	6.9
1	A	782	ILE	5.3
1	B	835	MET	4.9
1	B	792	GLU	4.8
1	B	820	GLU	4.3
1	A	796	LEU	4.2
1	B	836	GLU	4.1
1	A	835	MET	3.8
1	A	784	GLN	3.7
1	B	784	GLN	3.5
1	B	781	GLU	3.4
1	A	836	GLU	3.4
1	B	783	ALA	3.2
1	A	786	ARG	3.0
1	A	795	ALA	3.0
1	B	806	ARG	2.9
1	B	816	GLN	2.9
1	B	821	ASN	2.8
1	B	782	ILE	2.6
1	A	787	SER	2.6
1	A	791	ALA	2.6
1	A	831	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	815	GLU	2.5
1	B	785	VAL	2.5
1	B	811	GLU	2.5
1	A	788	LYS	2.5
1	B	780	GLU	2.3
1	B	817	LYS	2.1
1	B	834	LYS	2.1
1	A	800	LEU	2.1
1	B	809	SER	2.1
1	A	832	ILE	2.1
1	B	819	LYS	2.1
1	B	824	LEU	2.1
1	A	820	GLU	2.0
1	B	808	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.