



wwPDB EM Validation Summary Report ⓘ

Jul 15, 2024 – 07:48 am BST

PDB ID : 8OSJ
EMDB ID : EMD-17155
Title : Cryo-EM structure of CLOCK-BMAL1 bound to a nucleosomal E-box at position SHL-6.2 (DNA conformation 1)
Authors : Michael, A.K.; Stoos, L.; Kempf, G.; Cavadini, S.; Thoma, N.H.
Deposited on : 2023-04-19
Resolution : 6.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 6.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	64% 36%
1	E	139	62% 36%
2	B	106	73% 25%
2	F	106	73% 25%
3	C	133	73% 25%
3	G	133	74% 25%
4	D	128	70% 30%
4	H	128	70% 30%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	I	153	<div><div></div><div>81%</div><div></div><div></div><div>16%</div></div>
6	J	153	<div><div></div><div>81%</div><div></div><div></div><div>16%</div></div>
7	M	375	<div><div></div><div>17%</div><div></div><div></div><div>83%</div></div>
8	N	384	<div><div></div><div>14%</div><div></div><div></div><div>86%</div></div>

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 11493 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	89	Total	C	N	O	S	0	0
			728	460	136	128	4		
1	E	89	Total	C	N	O	S	0	0
			728	460	136	128	4		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P68431
A	-2	SER	-	expression tag	UNP P68431
A	-1	HIS	-	expression tag	UNP P68431
E	-3	GLY	-	expression tag	UNP P68431
E	-2	SER	-	expression tag	UNP P68431
E	-1	HIS	-	expression tag	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	79	Total	C	N	O	S	0	0
			633	399	124	109	1		
2	F	79	Total	C	N	O	S	0	0
			633	399	124	109	1		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P62805
B	-2	SER	-	expression tag	UNP P62805
B	-1	HIS	-	expression tag	UNP P62805
F	-3	GLY	-	expression tag	UNP P62805
F	-2	SER	-	expression tag	UNP P62805
F	-1	HIS	-	expression tag	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	100	Total	C	N	O	0	0
			769	486	148	135		
3	G	100	Total	C	N	O	0	0
			769	486	148	135		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P04908
C	-2	SER	-	expression tag	UNP P04908
C	-1	HIS	-	expression tag	UNP P04908
G	-3	GLY	-	expression tag	UNP P04908
G	-2	SER	-	expression tag	UNP P04908
G	-1	HIS	-	expression tag	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	90	Total	C	N	O	S	0	0
			699	441	123	133	2		
4	H	90	Total	C	N	O	S	0	0
			699	441	123	133	2		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P06899
D	-2	SER	-	expression tag	UNP P06899
D	-1	HIS	-	expression tag	UNP P06899
H	-3	GLY	-	expression tag	UNP P06899
H	-2	SER	-	expression tag	UNP P06899
H	-1	HIS	-	expression tag	UNP P06899

- Molecule 5 is a DNA chain called DNA (124-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	128	Total	C	N	O	P	0	0
			2611	1238	475	770	128		

- Molecule 6 is a DNA chain called DNA (124-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	128	Total	C	N	O	P	0	0
			2637	1246	497	766	128		

- Molecule 7 is a protein called Circadian locomoter output cycles protein kaput.

Mol	Chain	Residues	Atoms				AltConf	Trace
7	M	63	Total	C	N	O	0	0
			315	189	63	63		

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	21	GLY	-	expression tag	UNP O08785
M	22	ALA	-	expression tag	UNP O08785
M	23	MET	-	expression tag	UNP O08785
M	24	ASN	-	expression tag	UNP O08785
M	25	PRO	-	expression tag	UNP O08785

- Molecule 8 is a protein called Basic helix-loop-helix ARNT-like protein 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
8	N	54	Total	C	N	O	0	0
			272	164	54	54		

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
N	64	GLY	-	expression tag	UNP Q9WTL8
N	65	ALA	-	expression tag	UNP Q9WTL8
N	66	MET	-	expression tag	UNP Q9WTL8
N	67	ASN	-	expression tag	UNP Q9WTL8
N	68	PRO	-	expression tag	UNP Q9WTL8

- Molecule 4: Histone H2B type 1-J

- Molecule 4: Histone H2B type 1-J

- Molecule 5: DNA (124-MER)

- Molecule 6: DNA (124-MER)

- Molecule 7: Circadian locomoter output cycles protein kaput

THR	PRO	GLN	PHE	ILE	LYS	GLU	MET	CYS	THR	VAL	GLU	GLU	PRO	ASN	GLU	GLU	PHE	THR	SER	HIS	ARG	SER	LEU	GLU	TRP	LYS	PHE	PHE	LEU	ASP	HIS	ARG	ALA	PRO	PRO	ILE	ILE	GLY	TYR	LEU	PRO	PHE	GLU	VAL	LEU	GLY	THR	SER	GLY	TYR	HIS	VAL	ASP	ASP
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

GLU ASN
GLU LEU
ALA ARG
ALA ALA
CYS CYS
HIS HIS
GLU HIS
HIS MET
LEU MET
GLN GLY
TYR TYR
GLY GLY
LYS GLY
LYS LYS
LYS LYS
SER SER
CYS CYS
TYR TYR
ARG ARG
PHE PHE
LEU LEU
THR THR
LYS LYS
GLY GLY
TRP TRP
LEU LEU
GLN GLN
GLN GLN
ILE ILE
TYR TYR
LEU LEU
GLN GLN
TYR TYR
HIS HIS
HIS HIS
GLN GLN
TRP TRP
ASN ASN
SER SER
ARG ARG
PRO PRO
GLU GLU
PHE PHE
ILE ILE
VAL VAL
CYS CYS
THR THR
HIS HIS
VAL VAL
SER SER
TYR TYR

ALA
GLU
VAL
ARG
ALA
GLU
ARG
ARG
ARG
GLU
LEU
GLY
ILE
GLU
GLY
SER
LEU

● Molecule 8: Basic helix-loop-helix ARNT-like protein 1

Chain N: 14% 86%

GLY
ALA
MET
ASN
PRO
GLU
TYR
ALA
ALA
HIS
HIS
GLN
GLY
H76
H129
LYS
LYS
THR
LEU
LEU
ARG
GLY
ALA
THR
ASN
ASN
PRO
TYR
TYR
THR
GLU
GLY
ALA
ALA
ASN
ASN
LYS
PRO
LYS
THR
PHE
LEU
LEU
ASP
TYR
ALA
HIS
HIS
SER
SER
ASP
ASP
GLU
LYS
LEU
HIS
HIS
LEU
SER
SER
ILE
LEU
LEU
ARG
ALA
ALA
PRO
ASP
GLY
PHE
LEU
PHE
VAL
VAL
CYS
ASP
GLY

LYS
ILE
LEU
PHE
VAL
SER
GLU
SER
VAL
VAL
PHE
HIS
LYS
ILE
LEU
ASN
TYR
SER
GLN
ASN
ASP
SER
GLY
ILE
PHE
LEU
PHE
LEU
SER
PRO
ARG
GLU
ARG
PHE
LEU
ILE
ASP
ALA
LYS
THR
GLY
PRO

VAL
LYS
THR
ASP
THR
ILE
THR
PRO
GLY
VAL
VAL
PHE
SER
ARG
ARG
LEU
CYS
SER
SER
CYS
SER
GLY
ALA
ARG
ASN
CYS
PHE
PHE
GLY
SER
SER
MET
LEU
CYS
CYS
PHE
ASP
VAL
VAL
LYS
LYS
THR
ALA
ALA
GLN
GLU
PHE
LEU
VAL
SER
SER
VAL
PHE
ASP
LYS
LYS
SER
THR
ALA
GLY
THR
SER
HIS
ILE
THR
PHE
LEU

LYS
SER
TRP
PRO
PRO
THR
LYS
MET
GLY
LEU
ASP
ASP
GLU
ASN
GLU
PRO
ASP
ASN
ASN
GLY
GLY
CYS
ASN
GLY
LEU
SER
CYS
CYS
LEU
VAL
ALA
PHE
ILE
GLY
SER
ARG
VAL
LEU
HIS
ASP
GLU
HIS
SER
HIS
GLY
ASP
GLY
ILE
GLU
ILE
ARG
VAL
LYS
SER
MET
GLU
TYR
VAL
SER
SER
ARG
HIS
ALA
ASP
GLY

LYS
PHE
VAL
PHE
SER
PHE
ASP
GLN
ARG
ALA
THR
SER
ARG
ALA
ILE
TRP
PHE
LEU
ALA
TYR
LEU
PRO
GLN
GLU
LEU
LEU
GLY
THR
GLY
VAL
SER
CYS
TYR
GLU
TYR
PHE
ILE
SER
SER
HIS
HIS
LEU
ALA
GLU
CYS
HIS
HIS
GLY
HIS
ASP
GLY
VAL
LEU
VAL
GLN
ASN
GLY
ILE
LEU
GLN
THR
THR
ARG
GLU
LYS
ILE
THR
THR
ASN
CYS
TYR
LYS
PHE
LYS
LYS
ILE

LYS
ASP
GLY
SER
PHE
ILE
THR
LEU
ARG
ARG
SER
ARG
TRP
PHE
SER
PHE
MET
ASN
PRO
TRP
THR
LYS
GLY
VAL
VAL
TYR
ILE
VAL
SER
THR
ASN
THR
VAL
VAL

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	152914	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	200	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	0.013	Depositor
Minimum map value	-0.005	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.00126	Depositor
Map size (\AA)	242.08, 242.08, 242.08	wwPDB
Map dimensions	356, 356, 356	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.68, 0.68, 0.68	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/736	0.48	0/987
1	E	0.72	0/736	0.48	0/987
2	B	0.65	0/640	0.55	0/857
2	F	0.65	0/640	0.55	0/857
3	C	0.67	0/779	0.50	0/1054
3	G	0.66	0/779	0.50	0/1054
4	D	0.66	0/710	0.53	0/957
4	H	0.66	0/710	0.51	0/957
5	I	0.33	0/2925	1.00	0/4509
6	J	0.35	0/2961	1.00	0/4571
7	M	0.25	0/315	0.35	0/439
8	N	0.23	0/272	0.33	0/380
All	All	0.52	0/12203	0.80	0/17609

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	728	0	766	0	0
1	E	728	0	766	1	0
2	B	633	0	673	1	0
2	F	633	0	673	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	769	0	815	1	0
3	G	769	0	815	1	0
4	D	699	0	714	0	0
4	H	699	0	714	0	0
5	I	2611	0	1436	3	0
6	J	2637	0	1434	5	0
7	M	315	0	144	0	0
8	N	272	0	127	0	0
All	All	11493	0	9077	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 11 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:J:117:DA:H2''	6:J:118:DT:C5'	2.38	0.53
3:G:47:ALA:N	3:G:48:PRO:CD	2.75	0.48
5:I:120:DA:H2''	5:I:121:DG:C8	2.51	0.46
1:E:65:LEU:N	1:E:66:PRO:HD2	2.29	0.46
3:C:47:ALA:N	3:C:48:PRO:CD	2.82	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	87/139 (63%)	87 (100%)	0	0	100	100
1	E	87/139 (63%)	87 (100%)	0	0	100	100
2	B	77/106 (73%)	77 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	F	77/106 (73%)	77 (100%)	0	0	100	100
3	C	98/133 (74%)	98 (100%)	0	0	100	100
3	G	98/133 (74%)	96 (98%)	2 (2%)	0	100	100
4	D	88/128 (69%)	88 (100%)	0	0	100	100
4	H	88/128 (69%)	88 (100%)	0	0	100	100
7	M	61/375 (16%)	61 (100%)	0	0	100	100
8	N	52/384 (14%)	52 (100%)	0	0	100	100
All	All	813/1771 (46%)	811 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	78/113 (69%)	78 (100%)	0	100	100
1	E	78/113 (69%)	77 (99%)	1 (1%)	69	82
2	B	65/81 (80%)	65 (100%)	0	100	100
2	F	65/81 (80%)	65 (100%)	0	100	100
3	C	79/102 (78%)	78 (99%)	1 (1%)	69	82
3	G	79/102 (78%)	79 (100%)	0	100	100
4	D	76/106 (72%)	76 (100%)	0	100	100
4	H	76/106 (72%)	76 (100%)	0	100	100
7	M	1/345 (0%)	1 (100%)	0	100	100
8	N	1/342 (0%)	1 (100%)	0	100	100
All	All	598/1491 (40%)	596 (100%)	2 (0%)	92	95

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	20	ARG
1	E	129	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17155. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 178




Y Index: 178




Z Index: 178


6.2.2 Raw map



X Index: 178



Y Index: 178




Z Index: 178


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 174




Y Index: 203




Z Index: 170


6.3.2 Raw map



X Index: 176



Y Index: 182




Z Index: 174


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)


6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.00126. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 446 nm³; this corresponds to an approximate mass of 403 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.161 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.161 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	6.20	-	-
Author-provided FSC curve	6.10	7.89	6.84
Unmasked-calculated*	7.37	9.69	7.59

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.37 differs from the reported value 6.2 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17155 and PDB model 8OSJ. Per-residue inclusion information can be found in section [3](#) on page [7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.00126 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00126).

9.4 Atom inclusion ⓘ



At the recommended contour level, 100% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.00126) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.9670	<div><div></div></div> 0.2190
A	<div><div></div></div> 0.9400	<div><div></div></div> 0.2220
B	<div><div></div></div> 0.9460	<div><div></div></div> 0.2470
C	<div><div></div></div> 0.9170	<div><div></div></div> 0.2000
D	<div><div></div></div> 0.9280	<div><div></div></div> 0.2130
E	<div><div></div></div> 0.9450	<div><div></div></div> 0.2080
F	<div><div></div></div> 0.9290	<div><div></div></div> 0.2360
G	<div><div></div></div> 0.9420	<div><div></div></div> 0.2100
H	<div><div></div></div> 0.9560	<div><div></div></div> 0.2060
I	<div><div></div></div> 1.0000	<div><div></div></div> 0.2410
J	<div><div></div></div> 0.9950	<div><div></div></div> 0.2400
M	<div><div></div></div> 0.9430	<div><div></div></div> 0.0620
N	<div><div></div></div> 0.9930	<div><div></div></div> 0.0390

1.0

0.0

<0.0