



wwPDB EM Validation Summary Report ⓘ

Dec 31, 2024 – 07:40 PM EST

PDB ID : 8P0J
EMDB ID : EMD-17334
Title : Cryo EM map and model of the vaccinia RNA polymerase intermediate pre-initiation open promoter complex
Authors : Grimm, C.; Jungwirth, S.; Fischer, U.
Deposited on : 2023-05-10
Resolution : 2.39 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1286	
2	B	1164	
3	C	305	
4	E	185	
5	F	164	
6	G	161	
7	H	382	
8	J	63	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	L	287	
10	N	70	
11	O	844	
12	S	259	
13	T	70	
14	I	288	

2 Entry composition

There are 17 unique types of molecules in this entry. The entry contains 86467 atoms, of which 42799 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase 147 kDa polypeptide.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	1277	Total	C	H	N	O	S	0	0
			20617	6593	10357	1690	1931	46		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	258	THR	SER	variant	UNP P20504
A	489	GLU	LYS	variant	UNP P20504
A	1015	LYS	ARG	variant	UNP P20504

- Molecule 2 is a protein called DNA-directed RNA polymerase 133 kDa polypeptide.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	B	1155	Total	C	H	N	O	S	2	0
			18659	5934	9340	1595	1741	49		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	ASN	ASP	variant	UNP P68694
B	343	PHE	TYR	variant	UNP P68694

- Molecule 3 is a protein called DNA-directed RNA polymerase 35 kDa subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	C	304	Total	C	H	N	O	S	0	0
			4947	1608	2463	399	464	13		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	236	ASN	ASP	variant	UNP P21087

- Molecule 4 is a protein called DNA-directed RNA polymerase 22 kDa subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	E	184	Total	C	H	N	O	S	0	0
			3041	966	1546	248	276	5		

- Molecule 5 is a protein called DNA-directed RNA polymerase 19 kDa subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	F	104	Total	C	H	N	O	S	0	0
			1736	551	877	151	154	3		

- Molecule 6 is a protein called DNA-directed RNA polymerase 18 kDa subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	G	153	Total	C	H	N	O	S	0	0
			2370	753	1178	198	235	6		

- Molecule 7 is a protein called Intermediate transcription factor 3 large subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	H	382	Total	C	H	N	O	S	0	0
			6298	2004	3164	522	590	18		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	58	PRO	HIS	variant	UNP P20998

- Molecule 8 is a protein called DNA-directed RNA polymerase 7 kDa subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
8	J	61	Total	C	H	N	O	S	0	0
			1017	310	527	88	88	4		

- Molecule 9 is a protein called mRNA-capping enzyme regulatory subunit OPG124.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	L	284	Total	C	H	N	O	S	0	0
			4678	1492	2358	385	430	13		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	235	ASN	ASP	variant	UNP P20980

- Molecule 10 is a DNA chain called Non-template DNA oligomer.

Mol	Chain	Residues	Atoms						AltConf	Trace
10	N	39	Total	C	H	N	O	P	0	0
			1236	383	442	142	230	39		

- Molecule 11 is a protein called mRNA-capping enzyme catalytic subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
11	O	844	Total	C	H	N	O	S	0	0
			13715	4399	6884	1123	1290	19		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	8	PHE	SER	variant	UNP P20979
O	202	THR	LYS	variant	UNP P20979

- Molecule 12 is a protein called DNA-directed RNA polymerase 30 kDa polypeptide.

Mol	Chain	Residues	Atoms						AltConf	Trace
12	S	116	Total	C	H	N	O	S	0	0
			1916	609	961	155	187	4		

- Molecule 13 is a DNA chain called Template DNA oligomer.

Mol	Chain	Residues	Atoms						AltConf	Trace
13	T	36	Total	C	H	N	O	P	0	0
			1161	360	409	141	215	36		

- Molecule 14 is a protein called Intermediate transcription factor 3 small subunit.

Mol	Chain	Residues	Atoms						AltConf	Trace
14	I	276	Total	C	H	N	O	S	0	0
			4563	1469	2293	370	424	7		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	247	LYS	GLU	variant	UNP P20986

- Molecule 15 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
15	A	1	Total 1	Mg 1	0

- Molecule 16 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
16	A	2	Total 2	Zn 2	0
16	B	1	Total 1	Zn 1	0
16	J	1	Total 1	Zn 1	0

- Molecule 17 is water.

Mol	Chain	Residues	Atoms		AltConf
17	A	161	Total 161	O 161	0
17	B	165	Total 165	O 165	0
17	C	42	Total 42	O 42	0
17	E	18	Total 18	O 18	0
17	F	17	Total 17	O 17	0
17	G	18	Total 18	O 18	0
17	H	1	Total 1	O 1	0
17	J	6	Total 6	O 6	0
17	L	7	Total 7	O 7	0
17	N	2	Total 2	O 2	0
17	O	42	Total 42	O 42	0
17	S	24	Total 24	O 24	0
17	T	2	Total 2	O 2	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
17	I	3	Total	O	0
			3	3	


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-directed RNA polymerase 147 kDa polypeptide




- Molecule 2: DNA-directed RNA polymerase 133 kDa polypeptide







- Molecule 3: DNA-directed RNA polymerase 35 kDa subunit





- Molecule 4: DNA-directed RNA polymerase 22 kDa subunit





- Molecule 5: DNA-directed RNA polymerase 19 kDa subunit



- Molecule 6: DNA-directed RNA polymerase 18 kDa subunit




- Molecule 7: Intermediate transcription factor 3 large subunit





- Molecule 8: DNA-directed RNA polymerase 7 kDa subunit

Chain J: 79% 16% . .





- Molecule 9: mRNA-capping enzyme regulatory subunit OPG124

Chain L:  53% 85% 14%




- Molecule 10: Non-template DNA oligomer


Chain N:  29% 27% 44%




- Molecule 11: mRNA-capping enzyme catalytic subunit

Chain O:  5% 89% 11%







- Molecule 12: DNA-directed RNA polymerase 30 kDa polypeptide



- Molecule 13: Template DNA oligomer



- Molecule 14: Intermediate transcription factor 3 small subunit



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1421158	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	70	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	4.295	Depositor
Minimum map value	-1.708	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.110	Depositor
Recommended contour level	0.733	Depositor
Map size (Å)	408.38403, 408.38403, 408.38403	wwPDB
Map dimensions	384, 384, 384	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0635, 1.0635, 1.0635	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.25	0/10462	0.44	0/14135
2	B	0.25	0/9525	0.48	0/12866
3	C	0.26	0/2540	0.46	0/3440
4	E	0.25	0/1522	0.49	0/2069
5	F	0.23	0/874	0.47	0/1173
6	G	0.24	0/1209	0.48	0/1639
7	H	0.24	0/3189	0.44	0/4288
8	J	0.24	0/494	0.47	0/663
9	L	0.24	0/2365	0.45	0/3189
10	N	0.51	0/889	0.94	0/1365
11	O	0.25	0/6973	0.46	0/9433
12	S	0.24	0/970	0.44	0/1298
13	T	0.49	0/845	0.93	0/1303
14	I	0.24	0/2315	0.43	0/3123
All	All	0.26	0/44172	0.49	0/59984

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10260	10357	10377	70	0
2	B	9319	9340	9359	84	0
3	C	2484	2463	2473	21	0
4	E	1495	1546	1548	13	0
5	F	859	877	881	3	0
6	G	1192	1178	1181	11	0
7	H	3134	3164	3176	25	0
8	J	490	527	528	8	0
9	L	2320	2358	2363	25	0
10	N	794	442	444	14	0
11	O	6831	6884	6899	60	0
12	S	955	961	961	9	0
13	T	752	409	411	8	0
14	I	2270	2293	2299	16	0
15	A	1	0	0	0	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	J	1	0	0	0	0
17	A	161	0	0	7	0
17	B	165	0	0	14	0
17	C	42	0	0	0	0
17	E	18	0	0	1	0
17	F	17	0	0	1	0
17	G	18	0	0	0	0
17	H	1	0	0	0	0
17	I	3	0	0	0	0
17	J	6	0	0	1	0
17	L	7	0	0	0	0
17	N	2	0	0	0	0
17	O	42	0	0	2	0
17	S	24	0	0	1	0
17	T	2	0	0	0	0
All	All	43668	42799	42900	339	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 339 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:239:TYR:O	2:B:242:ARG:NH1	2.07	0.88
11:O:519:ARG:O	11:O:541:TYR:OH	1.94	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:745:ILE:O	1:A:749:THR:OG1	1.96	0.83
13:T:50:DT:OP1	14:I:128:LYS:NZ	2.12	0.81
2:B:1058:ARG:NH2	2:B:1059:ASP:OD1	2.17	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1273/1286 (99%)	1233 (97%)	40 (3%)	0	100	100
2	B	1155/1164 (99%)	1110 (96%)	45 (4%)	0	100	100
3	C	302/305 (99%)	288 (95%)	14 (5%)	0	100	100
4	E	182/185 (98%)	172 (94%)	10 (6%)	0	100	100
5	F	102/164 (62%)	99 (97%)	3 (3%)	0	100	100
6	G	149/161 (92%)	141 (95%)	8 (5%)	0	100	100
7	H	380/382 (100%)	372 (98%)	8 (2%)	0	100	100
8	J	59/63 (94%)	55 (93%)	4 (7%)	0	100	100
9	L	280/287 (98%)	270 (96%)	10 (4%)	0	100	100
11	O	842/844 (100%)	821 (98%)	21 (2%)	0	100	100
12	S	114/259 (44%)	109 (96%)	5 (4%)	0	100	100
14	I	272/288 (94%)	261 (96%)	11 (4%)	0	100	100
All	All	5110/5388 (95%)	4931 (96%)	179 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1148/1157 (99%)	1137 (99%)	11 (1%)	73	86
2	B	1057/1064 (99%)	1055 (100%)	2 (0%)	92	97
3	C	286/287 (100%)	286 (100%)	0	100	100
4	E	174/175 (99%)	173 (99%)	1 (1%)	84	92
5	F	95/151 (63%)	95 (100%)	0	100	100
6	G	136/144 (94%)	136 (100%)	0	100	100
7	H	363/363 (100%)	361 (99%)	2 (1%)	84	92
8	J	60/62 (97%)	58 (97%)	2 (3%)	33	53
9	L	269/272 (99%)	269 (100%)	0	100	100
11	O	774/774 (100%)	766 (99%)	8 (1%)	73	86
12	S	108/240 (45%)	108 (100%)	0	100	100
14	I	257/268 (96%)	256 (100%)	1 (0%)	89	95
All	All	4727/4957 (95%)	4700 (99%)	27 (1%)	82	92

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	H	20	PHE
8	J	9	THR
11	O	805	MET
8	J	3	PHE
11	O	281	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
11	O	21	ASN
11	O	35	ASN
11	O	147	ASN
6	G	127	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	159	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17334. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 188




Y Index: 188




Z Index: 160


6.3.2 Raw map



X Index: 188



Y Index: 188



Z Index: 193


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.733. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 269 nm³; this corresponds to an approximate mass of 243 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.418 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.418 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.39	-	-
Author-provided FSC curve	2.39	2.71	2.42
Unmasked-calculated*	2.93	3.50	2.99

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 2.93 differs from the reported value 2.39 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17334 and PDB model 8P0J. Per-residue inclusion information can be found in section [3](#) on page [9](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.733 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.733).































9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 83% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.733) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8280	 0.5580
A	 0.8860	 0.6050
B	 0.9090	 0.6160
C	 0.9180	 0.6140
E	 0.9140	 0.6080
F	 0.9290	 0.6530
G	 0.6650	 0.4810
H	 0.7580	 0.4690
I	 0.8040	 0.5460
J	 0.9480	 0.6220
L	 0.4020	 0.3010
N	 0.8290	 0.4540
O	 0.8100	 0.5350
S	 0.6680	 0.5100
T	 0.8380	 0.4620

