



## Full wwPDB EM Validation Report ⓘ

Jan 13, 2025 – 05:24 pm GMT

PDB ID : 8PEY  
EMDB ID : EMD-17641  
Title : Rho P167L-ATPgS-Psu complex II locked  
Authors : Gjorgjevikj, D.; Wahl, M.C.; Hilal, T.; Loll, B.  
Deposited on : 2023-06-15  
Resolution : 3.00 Å(reported)  
Based on initial models : 3RX6, 1PV4

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div> <div>11%</div> <div>57%</div> <div>43%</div> </div>
1	B	419	<div> <div>8%</div> <div>100%</div> </div>
1	C	419	<div> <div>100%</div> </div>
1	D	419	<div> <div>100%</div> </div>
1	E	419	<div> <div>100%</div> </div>
1	F	419	<div> <div>15%</div> <div>100%</div> </div>
1	G	419	<div> <div>33%</div> <div>88%</div> <div>11%</div> </div>
1	I	419	<div> <div>22%</div> <div>90%</div> <div>9%</div> </div>
1	J	419	<div> <div>100%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	K	419	 100%
1	L	419	 100%
1	M	419	 100%
1	N	419	 17% 99% .
2	a	190	 12% 97% ..
2	b	190	 7% 98% .
2	c	190	 98% .
2	d	190	 98% .
2	e	190	 98% .
2	f	190	 98% .
2	g	190	 98% .
2	h	190	 98% .
2	i	190	 98% ..
2	j	190	 98% .

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 55957 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription termination factor Rho.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	240	Total	C	N	O	S	0	0
			1875	1185	332	348	10		
1	B	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	C	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	D	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	E	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	F	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	G	374	Total	C	N	O	S	0	0
			2944	1852	523	555	14		
1	I	380	Total	C	N	O	S	0	0
			2984	1886	526	559	13		
1	J	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	K	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	L	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	M	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		
1	N	419	Total	C	N	O	S	0	0
			3299	2075	586	621	17		

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	167	LEU	PRO	engineered mutation	UNP P0AG30
B	167	LEU	PRO	engineered mutation	UNP P0AG30
C	167	LEU	PRO	engineered mutation	UNP P0AG30
D	167	LEU	PRO	engineered mutation	UNP P0AG30

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	167	LEU	PRO	engineered mutation	UNP P0AG30
F	167	LEU	PRO	engineered mutation	UNP P0AG30
G	167	LEU	PRO	engineered mutation	UNP P0AG30
I	167	LEU	PRO	engineered mutation	UNP P0AG30
J	167	LEU	PRO	engineered mutation	UNP P0AG30
K	167	LEU	PRO	engineered mutation	UNP P0AG30
L	167	LEU	PRO	engineered mutation	UNP P0AG30
M	167	LEU	PRO	engineered mutation	UNP P0AG30
N	167	LEU	PRO	engineered mutation	UNP P0AG30

- Molecule 2 is a protein called Polarity suppression protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	a	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	b	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	c	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	d	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	e	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	f	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	g	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	h	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	i	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		
2	j	187	Total	C	N	O	S	0	0
			1478	911	279	283	5		

- Molecule 3 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter code: AGS) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>S).



Idol	Chain	Residues	Atoms						AltConf
3	B	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	C	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	D	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	E	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	F	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	G	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	I	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	J	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	K	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	L	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	M	1	Total 31	C 10	N 5	O 12	P 3	S 1	0
3	N	1	Total 31	C 10	N 5	O 12	P 3	S 1	0


- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
4	B	1	Total 1	Mg 1	0
4	C	1	Total 1	Mg 1	0
4	D	1	Total 1	Mg 1	0
4	E	1	Total 1	Mg 1	0
4	F	1	Total 1	Mg 1	0
4	G	1	Total 1	Mg 1	0
4	I	1	Total 1	Mg 1	0
4	J	1	Total 1	Mg 1	0
4	K	1	Total 1	Mg 1	0
4	L	1	Total 1	Mg 1	0
4	M	1	Total 1	Mg 1	0
4	N	1	Total 1	Mg 1	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription termination factor Rho



- Molecule 1: Transcription termination factor Rho




- Molecule 1: Transcription termination factor Rho



- Molecule 1: Transcription termination factor Rho












- Molecule 1: Transcription termination factor Rho





- Molecule 1: Transcription termination factor Rho



- Molecule 1: Transcription termination factor Rho




- Molecule 1: Transcription termination factor Rho





- Molecule 1: Transcription termination factor Rho

Chain J: 100%



- Molecule 1: Transcription termination factor Rho

Chain K: 100%




- Molecule 1: Transcription termination factor Rho

Chain L: 100%





- Molecule 1: Transcription termination factor Rho

Chain M: 100%




- Molecule 1: Transcription termination factor Rho


Chain N: 17% 99%




- Molecule 2: Polarity suppression protein




- Molecule 2: Polarity suppression protein




- Molecule 2: Polarity suppression protein




- Molecule 2: Polarity suppression protein




- Molecule 2: Polarity suppression protein



- Molecule 2: Polarity suppression protein




- Molecule 2: Polarity suppression protein




## • Molecule 2: Polarity suppression protein

Chain h:  98% .




## • Molecule 2: Polarity suppression protein

Chain i:  98% ..



## • Molecule 2: Polarity suppression protein

Chain j:  98% .



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	317635	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	42	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	2.245	Depositor
Minimum map value	-0.000	Depositor
Average map value	0.010	Depositor
Map value standard deviation	0.059	Depositor
Recommended contour level	0.25	Depositor
Map size (Å)	372.73602, 372.73602, 372.73602	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.8320001, 0.8320001, 0.8320001	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AGS, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/1901	0.58	0/2563
1	B	0.26	0/3347	0.55	1/4504 (0.0%)
1	C	0.25	0/3347	0.52	0/4504
1	D	0.25	0/3347	0.52	0/4504
1	E	0.25	0/3347	0.51	0/4504
1	F	0.24	0/3347	0.53	0/4504
1	G	0.26	0/2989	0.59	0/4024
1	I	0.26	0/3029	0.55	0/4082
1	J	0.25	0/3347	0.53	0/4504
1	K	0.25	0/3347	0.51	0/4504
1	L	0.25	0/3347	0.51	0/4504
1	M	0.26	0/3347	0.54	0/4504
1	N	0.25	0/3347	0.55	0/4504
2	a	0.25	0/1498	0.53	0/2020
2	b	0.26	0/1498	0.56	0/2020
2	c	0.27	0/1498	0.55	0/2020
2	d	0.25	0/1498	0.53	0/2020
2	e	0.24	0/1498	0.50	0/2020
2	f	0.25	0/1498	0.51	0/2020
2	g	0.24	0/1498	0.50	0/2020
2	h	0.24	0/1498	0.50	0/2020
2	i	0.26	0/1498	0.54	0/2020
2	j	0.24	0/1498	0.51	0/2020
All	All	0.25	0/56369	0.53	1/75909 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	162	LEU	CA-CB-CG	5.18	127.20	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/419 (57%)	225 (94%)	13 (6%)	0	100	100
1	B	417/419 (100%)	403 (97%)	14 (3%)	0	100	100
1	C	417/419 (100%)	410 (98%)	7 (2%)	0	100	100
1	D	417/419 (100%)	413 (99%)	4 (1%)	0	100	100
1	E	417/419 (100%)	407 (98%)	10 (2%)	0	100	100
1	F	417/419 (100%)	407 (98%)	10 (2%)	0	100	100
1	G	372/419 (89%)	362 (97%)	10 (3%)	0	100	100
1	I	376/419 (90%)	359 (96%)	17 (4%)	0	100	100
1	J	417/419 (100%)	410 (98%)	7 (2%)	0	100	100
1	K	417/419 (100%)	407 (98%)	10 (2%)	0	100	100
1	L	417/419 (100%)	407 (98%)	10 (2%)	0	100	100
1	M	417/419 (100%)	409 (98%)	8 (2%)	0	100	100
1	N	417/419 (100%)	402 (96%)	15 (4%)	0	100	100
2	a	185/190 (97%)	179 (97%)	6 (3%)	0	100	100
2	b	185/190 (97%)	182 (98%)	3 (2%)	0	100	100
2	c	185/190 (97%)	183 (99%)	2 (1%)	0	100	100
2	d	185/190 (97%)	183 (99%)	2 (1%)	0	100	100
2	e	185/190 (97%)	184 (100%)	1 (0%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	f	185/190 (97%)	185 (100%)	0	0	100	100
2	g	185/190 (97%)	183 (99%)	2 (1%)	0	100	100
2	h	185/190 (97%)	184 (100%)	1 (0%)	0	100	100
2	i	185/190 (97%)	182 (98%)	3 (2%)	0	100	100
2	j	185/190 (97%)	183 (99%)	2 (1%)	0	100	100
All	All	7006/7347 (95%)	6849 (98%)	157 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/359 (56%)	200 (98%)	3 (2%)	60	83
1	B	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	C	359/359 (100%)	357 (99%)	2 (1%)	84	93
1	D	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	E	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	F	359/359 (100%)	359 (100%)	0	100	100
1	G	319/359 (89%)	313 (98%)	6 (2%)	52	79
1	I	324/359 (90%)	321 (99%)	3 (1%)	75	89
1	J	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	K	359/359 (100%)	357 (99%)	2 (1%)	84	93
1	L	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	M	359/359 (100%)	358 (100%)	1 (0%)	91	96
1	N	359/359 (100%)	356 (99%)	3 (1%)	79	90
2	a	153/156 (98%)	151 (99%)	2 (1%)	65	85
2	b	153/156 (98%)	153 (100%)	0	100	100
2	c	153/156 (98%)	153 (100%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	d	153/156 (98%)	153 (100%)	0	100	100
2	e	153/156 (98%)	153 (100%)	0	100	100
2	f	153/156 (98%)	153 (100%)	0	100	100
2	g	153/156 (98%)	153 (100%)	0	100	100
2	h	153/156 (98%)	153 (100%)	0	100	100
2	i	153/156 (98%)	152 (99%)	1 (1%)	81	91
2	j	153/156 (98%)	153 (100%)	0	100	100
All	All	5966/6227 (96%)	5938 (100%)	28 (0%)	85	94

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	LYS
1	A	252	ARG
1	A	305	ARG
1	B	348	LYS
1	C	105	LYS
1	C	417	LYS
1	D	115	LYS
1	E	88	ARG
1	G	221	ARG
1	G	249	LYS
1	G	326	LYS
1	G	353	ARG
1	G	362	ARG
1	G	385	LYS
1	I	283	LYS
1	I	326	LYS
1	I	353	ARG
1	J	109	ARG
1	K	109	ARG
1	K	275	ASN
1	L	32	GLN
1	M	417	LYS
1	N	221	ARG
1	N	348	LYS
1	N	362	ARG
2	a	52	ARG
2	a	156	ARG
2	i	136	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (28) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	189	GLN
1	A	190	ASN
1	A	241	GLN
1	A	275	ASN
1	B	241	GLN
1	C	172	GLN
1	C	275	ASN
1	D	193	GLN
1	F	25	ASN
1	F	90	ASN
1	G	129	ASN
1	G	172	GLN
1	G	239	HIS
1	I	256	HIS
1	I	340	ASN
1	J	140	HIS
1	J	172	GLN
1	J	275	ASN
1	L	42	HIS
1	M	41	GLN
1	M	239	HIS
1	M	241	GLN
1	N	8	ASN
1	N	135	ASN
1	N	190	ASN
2	a	63	ASN
2	i	120	GLN
2	j	91	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 12 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AGS	F	501	4	26,33,33	0.70	1 (3%)	26,52,52	1.10	2 (7%)
3	AGS	D	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.05	2 (7%)
3	AGS	J	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.08	2 (7%)
3	AGS	M	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.05	2 (7%)
3	AGS	L	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.04	2 (7%)
3	AGS	C	501	4	26,33,33	0.70	1 (3%)	26,52,52	1.06	2 (7%)
3	AGS	G	501	4	26,33,33	0.70	1 (3%)	26,52,52	1.13	2 (7%)
3	AGS	K	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.07	2 (7%)
3	AGS	N	501	4	26,33,33	0.70	1 (3%)	26,52,52	1.11	3 (11%)
3	AGS	E	501	4	26,33,33	0.71	1 (3%)	26,52,52	1.07	2 (7%)
3	AGS	I	501	4	26,33,33	0.72	1 (3%)	26,52,52	1.28	3 (11%)
3	AGS	B	501	4	26,33,33	0.70	1 (3%)	26,52,52	1.13	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AGS	F	501	4	-	3/17/38/38	0/3/3/3
3	AGS	D	501	4	-	2/17/38/38	0/3/3/3
3	AGS	J	501	4	-	3/17/38/38	0/3/3/3
3	AGS	M	501	4	-	3/17/38/38	0/3/3/3

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AGS	L	501	4	-	2/17/38/38	0/3/3/3
3	AGS	C	501	4	-	3/17/38/38	0/3/3/3
3	AGS	G	501	4	-	1/17/38/38	0/3/3/3
3	AGS	K	501	4	-	2/17/38/38	0/3/3/3
3	AGS	N	501	4	-	1/17/38/38	0/3/3/3
3	AGS	E	501	4	-	3/17/38/38	0/3/3/3
3	AGS	I	501	4	-	2/17/38/38	0/3/3/3
3	AGS	B	501	4	-	1/17/38/38	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	I	501	AGS	PG-S1G	2.25	1.95	1.90
3	N	501	AGS	PG-S1G	2.14	1.95	1.90
3	G	501	AGS	PG-S1G	2.14	1.95	1.90
3	F	501	AGS	PG-S1G	2.13	1.95	1.90
3	B	501	AGS	PG-S1G	2.12	1.95	1.90
3	M	501	AGS	PG-S1G	2.11	1.95	1.90
3	J	501	AGS	PG-S1G	2.10	1.95	1.90
3	E	501	AGS	PG-S1G	2.10	1.95	1.90
3	D	501	AGS	PG-S1G	2.09	1.95	1.90
3	L	501	AGS	PG-S1G	2.09	1.95	1.90
3	K	501	AGS	PG-S1G	2.08	1.95	1.90
3	C	501	AGS	PG-S1G	2.08	1.95	1.90

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	I	501	AGS	PA-O3A-PB	-4.80	116.34	132.83
3	G	501	AGS	PA-O3A-PB	-4.09	118.80	132.83
3	F	501	AGS	PA-O3A-PB	-3.85	119.63	132.83
3	E	501	AGS	PA-O3A-PB	-3.83	119.69	132.83
3	J	501	AGS	PA-O3A-PB	-3.81	119.75	132.83
3	M	501	AGS	PA-O3A-PB	-3.80	119.77	132.83
3	K	501	AGS	PA-O3A-PB	-3.78	119.85	132.83
3	C	501	AGS	PA-O3A-PB	-3.74	119.99	132.83
3	D	501	AGS	PA-O3A-PB	-3.73	120.02	132.83
3	L	501	AGS	PA-O3A-PB	-3.72	120.05	132.83
3	B	501	AGS	PA-O3A-PB	-3.67	120.22	132.83
3	N	501	AGS	PA-O3A-PB	-3.62	120.39	132.83
3	B	501	AGS	C5-C6-N6	2.28	123.82	120.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	501	AGS	C5-C6-N6	2.28	123.82	120.35
3	G	501	AGS	C5-C6-N6	2.28	123.81	120.35
3	C	501	AGS	C5-C6-N6	2.28	123.81	120.35
3	F	501	AGS	C5-C6-N6	2.27	123.81	120.35
3	E	501	AGS	C5-C6-N6	2.27	123.80	120.35
3	J	501	AGS	C5-C6-N6	2.27	123.80	120.35
3	N	501	AGS	C5-C6-N6	2.27	123.80	120.35
3	K	501	AGS	C5-C6-N6	2.27	123.80	120.35
3	M	501	AGS	C5-C6-N6	2.26	123.79	120.35
3	I	501	AGS	C5-C6-N6	2.26	123.79	120.35
3	D	501	AGS	C5-C6-N6	2.26	123.78	120.35
3	I	501	AGS	C3'-C2'-C1'	2.15	104.21	100.98
3	N	501	AGS	C3'-C2'-C1'	2.14	104.19	100.98

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	501	AGS	C5'-O5'-PA-O3A
3	D	501	AGS	O4'-C4'-C5'-O5'
3	E	501	AGS	C5'-O5'-PA-O3A
3	F	501	AGS	C5'-O5'-PA-O3A
3	L	501	AGS	O4'-C4'-C5'-O5'
3	M	501	AGS	O4'-C4'-C5'-O5'
3	M	501	AGS	C5'-O5'-PA-O3A
3	B	501	AGS	PA-O3A-PB-O1B
3	J	501	AGS	PA-O3A-PB-O1B
3	N	501	AGS	PA-O3A-PB-O1B
3	C	501	AGS	C5'-O5'-PA-O1A
3	C	501	AGS	C5'-O5'-PA-O2A
3	E	501	AGS	C5'-O5'-PA-O1A
3	E	501	AGS	C5'-O5'-PA-O2A
3	F	501	AGS	C5'-O5'-PA-O1A
3	F	501	AGS	C5'-O5'-PA-O2A
3	M	501	AGS	PA-O3A-PB-O1B
3	L	501	AGS	C3'-C4'-C5'-O5'
3	J	501	AGS	PB-O3B-PG-O2G
3	I	501	AGS	PA-O3A-PB-O1B
3	D	501	AGS	C3'-C4'-C5'-O5'
3	I	501	AGS	PG-O3B-PB-O1B
3	K	501	AGS	O4'-C4'-C5'-O5'
3	G	501	AGS	C5'-O5'-PA-O1A

*Continued on next page...*


*Continued from previous page...*



Mol	Chain	Res	Type	Atoms
3	J	501	AGS	C5'-O5'-PA-O1A
3	K	501	AGS	C5'-O5'-PA-O1A



There are no ring outliers.

No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17641. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 224




Y Index: 224




Z Index: 224


### 6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 244




Y Index: 226




Z Index: 197


### 6.3.2 Raw map



X Index: 244



Y Index: 226




Z Index: 197

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.25. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 482 nm<sup>3</sup>; this corresponds to an approximate mass of 436 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of  $0.333 \text{ \AA}^{-1}$

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.333 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	2.95	3.37	3.02
Unmasked-calculated*	3.79	7.07	3.85

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.79 differs from the reported value 3.0 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17641 and PDB model 8PEY. Per-residue inclusion information can be found in section 3 on page 8.


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.25 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.25).



















































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 92% of all backbone atoms, 81% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.25) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8100	 0.4070
A	 0.6300	 0.3010
B	 0.7740	 0.3790
C	 0.8730	 0.4650
D	 0.8880	 0.4880
E	 0.8680	 0.4690
F	 0.7360	 0.3800
G	 0.5020	 0.2600
I	 0.6500	 0.3170
J	 0.8860	 0.4680
K	 0.9150	 0.5030
L	 0.9220	 0.5060
M	 0.8710	 0.4530
N	 0.6850	 0.3190
a	 0.6620	 0.2690
b	 0.7630	 0.3390
c	 0.8830	 0.3960
d	 0.9090	 0.4280
e	 0.9000	 0.4320
f	 0.9050	 0.4300
g	 0.9040	 0.4280
h	 0.9010	 0.4250
i	 0.8760	 0.3800
j	 0.8480	 0.3480

