



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 5, 2024 – 06:31 PM EDT

PDB ID : 2PGC  
Title : Crystal structure of a a marine metagenome protein (jcvi\_pep\_1096685590403) from uncultured marine organism at 2.53 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-04-09  
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6935 (2.54-2.50)
Clashscore	180529	7778 (2.54-2.50)
Ramachandran outliers	177936	7674 (2.54-2.50)
Sidechain outliers	177891	7676 (2.54-2.50)
RSRZ outliers	164620	6935 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	<div> <div>3%</div> <div>94%</div> <div>6%</div> </div>
1	B	207	<div> <div>%</div> <div>87%</div> <div>11%</div> <div>.</div> </div>
1	C	207	<div> <div>5%</div> <div>81%</div> <div>15%</div> <div>..</div> </div>
1	D	207	<div> <div>89%</div> <div>9%</div> <div>.</div> </div>
1	E	207	<div> <div>4%</div> <div>89%</div> <div>9%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	Se	0	1	1
			1602	1018	263	311	2	8			
1	B	205	Total	C	N	O	S	Se	0	0	0
			1573	999	256	309	2	7			
1	C	203	Total	C	N	O	S	Se	0	1	0
			1550	985	251	304	2	8			
1	D	205	Total	C	N	O	S	Se	0	1	0
			1580	1008	255	307	2	8			
1	E	205	Total	C	N	O	S	Se	0	1	0
			1578	1001	257	311	2	7			

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	2	Total	Cl	0	0
			2	2		
2	E	1	Total	Cl	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	26	Total	O	0	0
			26	26		
3	B	34	Total	O	0	0
			34	34		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	22	Total 22	O 22	0	0
3	D	27	Total 27	O 27	0	0
3	E	22	Total 22	O 22	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: uncharacterized protein




- Molecule 1: uncharacterized protein




- Molecule 1: uncharacterized protein




- Molecule 1: uncharacterized protein



- Molecule 1: uncharacterized protein





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	137.47Å 139.94Å 60.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.92 – 2.53 29.92 – 2.53	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.92-2.53) 99.7 (29.92-2.53)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.90 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.214 , 0.246 0.216 , 0.244	Depositor DCC
$R_{free}$ test set	2000 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	54.2	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 53.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8020	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.57	0/1624	0.64	0/2176
1	B	0.58	0/1592	0.68	0/2138
1	C	0.54	0/1570	0.63	0/2110
1	D	0.59	0/1603	0.65	0/2151
1	E	0.57	0/1602	0.65	0/2155
All	All	0.57	0/7991	0.65	0/10730

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1573	10	0
1	B	1573	0	1530	19	0
1	C	1550	0	1498	29	0
1	D	1580	0	1543	10	0
1	E	1578	0	1525	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	2	0	0	0	0
2	E	1	0	0	0	0
3	A	26	0	0	1	0
3	B	34	0	0	0	0
3	C	22	0	0	1	0
3	D	27	0	0	0	0
3	E	22	0	0	0	0
All	All	8020	0	7669	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:TYR:CZ	1:B:172:MSE:HE2	2.09	0.88
1:A:176:GLU:HB2	1:B:172:MSE:HE3	1.54	0.87
1:C:14:VAL:HG21	1:C:56:LEU:HD12	1.59	0.83
1:C:115:ILE:HD12	1:C:205:ILE:HD11	1.69	0.75
1:B:14:VAL:HG21	1:B:56:LEU:HD12	1.71	0.72
1:E:43:ARG:HD2	1:E:115:ILE:HD11	1.72	0.71
1:C:115:ILE:CD1	1:C:205:ILE:HD11	2.21	0.70
1:C:114:TYR:CE1	1:C:172:MSE:HE2	2.27	0.69
1:D:4:ILE:HD12	1:D:4:ILE:H	1.57	0.69
1:D:84:ILE:HG22	1:D:85:MSE:HE2	1.73	0.69
1:E:29:TYR:CE2	1:E:33:LEU:HD11	2.27	0.69
1:A:176:GLU:CB	1:B:172:MSE:HE3	2.22	0.68
1:C:69:GLN:HB2	1:E:172:MSE:HG3	1.79	0.64
1:B:131:LEU:O	1:B:135:ASN:ND2	2.31	0.64
1:C:28:SER:HB3	1:C:80:VAL:HG21	1.79	0.64
1:A:57:ILE:HG21	1:A:203:ILE:HD13	1.82	0.62
1:C:28:SER:CB	1:C:80:VAL:HG21	2.31	0.61
1:E:84:ILE:HG22	1:E:85:MSE:CE	2.31	0.60
1:C:8:ILE:HD12	1:C:97:ILE:HG23	1.83	0.60
1:E:14:VAL:HG21	1:E:56:LEU:HD12	1.84	0.60
1:A:81:PHE:CE1	1:A:85:MSE:HE3	2.37	0.59
1:B:114:TYR:CE1	1:B:172:MSE:HE2	2.37	0.59
1:C:29:TYR:CE2	1:C:33:LEU:HD11	2.38	0.59
1:C:4:ILE:CD1	1:C:4:ILE:N	2.66	0.59
1:A:81:PHE:CZ	1:A:85:MSE:HE3	2.38	0.58
1:D:114:TYR:CE1	1:D:172:MSE:HE2	2.39	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:ILE:HD12	1:A:8:ILE:N	2.20	0.57
1:E:144:ASN:HD22	1:E:144:ASN:N	2.03	0.56
1:E:84:ILE:HG22	1:E:85:MSE:HE2	1.88	0.55
1:D:57:ILE:HG21	1:D:203:ILE:HD13	1.90	0.54
1:D:29:TYR:CE2	1:D:33:LEU:HD11	2.43	0.53
1:C:122:ALA:HB3	1:C:163:TYR:CE1	2.44	0.52
1:C:191:LEU:HD12	1:C:194:PHE:CZ	2.44	0.52
1:B:8:ILE:N	1:B:8:ILE:CD1	2.73	0.51
1:B:149:LEU:C	1:B:149:LEU:HD23	2.30	0.51
1:C:8:ILE:HD13	1:C:8:ILE:N	2.26	0.51
1:C:149:LEU:C	1:C:149:LEU:HD23	2.32	0.50
1:B:112:PRO:HB3	1:B:147:LEU:HD12	1.94	0.50
1:C:8:ILE:N	1:C:8:ILE:CD1	2.75	0.49
1:B:8:ILE:N	1:B:8:ILE:HD12	2.26	0.49
1:D:29:TYR:OH	1:D:75:GLN:HG2	2.13	0.48
1:B:191:LEU:HD23	1:B:192:MSE:HE3	1.94	0.48
1:E:8:ILE:HD12	1:E:8:ILE:N	2.30	0.47
1:D:122:ALA:HB3	1:D:163:TYR:CE1	2.50	0.47
1:B:4:ILE:H	1:B:4:ILE:HD12	1.79	0.46
1:C:25:LEU:O	1:C:80:VAL:HG11	2.15	0.46
1:C:199:MSE:HE2	3:C:209:HOH:O	2.15	0.46
1:A:16:PHE:HA	1:A:19[A]:ARG:NE	2.31	0.46
1:A:176:GLU:HB2	1:B:172:MSE:CE	2.37	0.46
1:B:4:ILE:HD12	1:B:4:ILE:N	2.31	0.45
1:B:84:ILE:HG22	1:B:85:MSE:HE2	1.98	0.45
1:C:118:THR:HG21	2:C:207:CL:CL	2.54	0.45
1:E:191:LEU:HG	1:E:192:MSE:CE	2.47	0.45
1:B:84:ILE:HG22	1:B:85:MSE:CE	2.47	0.44
1:C:29:TYR:HB2	1:C:80:VAL:HG11	1.99	0.44
1:B:114:TYR:CZ	1:B:172:MSE:CE	2.93	0.44
1:C:4:ILE:N	1:C:4:ILE:HD12	2.32	0.44
1:A:160:VAL:O	1:A:160:VAL:HG13	2.18	0.43
1:D:191:LEU:HG	1:D:192:MSE:HE2	2.00	0.43
1:C:28:SER:OG	1:C:80:VAL:HG21	2.19	0.43
1:C:18:TYR:OH	1:C:87:SER:HB3	2.19	0.43
1:E:149:LEU:C	1:E:149:LEU:HD23	2.39	0.43
1:C:43:ARG:HD2	1:C:115:ILE:HD11	2.01	0.42
1:A:80:VAL:HG23	3:A:230:HOH:O	2.20	0.42
1:E:111[A]:HIS:CE1	1:E:205:ILE:HG21	2.55	0.42
1:B:29:TYR:CE2	1:B:33:LEU:HD11	2.54	0.42
1:C:115:ILE:HD11	1:C:205:ILE:HD11	2.00	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:107:GLN:OE1	1:D:148:THR:HA	2.20	0.41
1:E:29:TYR:HE2	1:E:33:LEU:HD11	1.81	0.41
1:D:112:PRO:HB3	1:D:147:LEU:HD12	2.02	0.41
1:C:131:LEU:O	1:C:135:ASN:ND2	2.54	0.41
1:C:5:ASN:HB2	1:C:100:SER:HB3	2.02	0.41
1:B:192:MSE:HA	1:B:192:MSE:HE2	2.02	0.41
1:C:81:PHE:CE2	1:C:85:MSE:HG3	2.56	0.41
1:E:122:ALA:HB3	1:E:163:TYR:CE1	2.56	0.41
1:C:183:LEU:HD23	1:C:188:TYR:CE2	2.56	0.40
1:C:114:TYR:CZ	1:C:172:MSE:HE2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/207 (100%)	204 (99%)	2 (1%)	0	100	100
1	B	203/207 (98%)	201 (99%)	2 (1%)	0	100	100
1	C	202/207 (98%)	199 (98%)	3 (2%)	0	100	100
1	D	204/207 (99%)	202 (99%)	2 (1%)	0	100	100
1	E	204/207 (99%)	202 (99%)	2 (1%)	0	100	100
All	All	1019/1035 (98%)	1008 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/169 (101%)	169 (99%)	2 (1%)	67	85
1	B	167/169 (99%)	159 (95%)	8 (5%)	21	41
1	C	163/169 (96%)	156 (96%)	7 (4%)	25	45
1	D	167/169 (99%)	164 (98%)	3 (2%)	54	77
1	E	169/169 (100%)	166 (98%)	3 (2%)	54	77
All	All	837/845 (99%)	814 (97%)	23 (3%)	38	65

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	130	PHE
1	B	8	ILE
1	B	28	SER
1	B	108	SER
1	B	128	ASP
1	B	130	PHE
1	B	139	SER
1	B	186	SER
1	B	199	MSE
1	C	4	ILE
1	C	8	ILE
1	C	30	SER
1	C	80	VAL
1	C	130	PHE
1	C	136	ASP
1	C	199	MSE
1	D	3	ASN
1	D	4	ILE
1	D	130	PHE
1	E	130	PHE
1	E	144	ASN
1	E	182	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	3	ASN
1	C	75	GLN
1	E	144	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/207 (96%)	0.22	6 (3%)	52	51	40, 66, 82, 88	1 (0%)
1	B	198/207 (95%)	0.40	2 (1%)	79	77	47, 66, 82, 90	0
1	C	196/207 (94%)	0.64	10 (5%)	34	33	45, 65, 81, 84	0
1	D	198/207 (95%)	0.17	1 (0%)	87	86	46, 66, 82, 88	0
1	E	198/207 (95%)	0.43	9 (4%)	39	37	41, 66, 81, 89	1 (0%)
All	All	989/1035 (95%)	0.37	28 (2%)	55	53	40, 66, 82, 90	2 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	109	TYR	3.7
1	E	183	LEU	3.3
1	A	184	ALA	3.2
1	C	206	LEU	3.1
1	C	86	ASP	3.0
1	E	184	ALA	2.9
1	C	80	VAL	2.5
1	E	140	CYS	2.5
1	C	29	TYR	2.4
1	C	123	ALA	2.4
1	C	134	ILE	2.4
1	B	79	SER	2.4
1	B	21	THR	2.4
1	A	80	VAL	2.3
1	E	109	TYR	2.2
1	A	111	HIS	2.2
1	E	136	ASP	2.2
1	C	34	ILE	2.2
1	C	84	ILE	2.2
1	E	188	TYR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	2.1
1	C	87	SER	2.1
1	E	182	LEU	2.1
1	A	0	GLY	2.1
1	C	105	PHE	2.1
1	E	186	SER	2.0
1	D	80	VAL	2.0
1	E	111[A]	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	CL	E	207	1/1	0.88	0.12	72,72,72,72	0
2	CL	D	207	1/1	0.91	0.11	78,78,78,78	0
2	CL	A	207	1/1	0.93	0.14	74,74,74,74	0
2	CL	B	207	1/1	0.93	0.08	69,69,69,69	0
2	CL	C	207	1/1	0.94	0.08	74,74,74,74	0
2	CL	D	208	1/1	0.97	0.10	67,67,67,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.