



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 11, 2024 – 02:19 AM EST

PDB ID : 1PL6  
Title : Human SDH/NADH/inhibitor complex  
Authors : Pauly, T.A.; Ekstrom, J.L.; Beebe, D.A.; Chrnyk, B.; Cunningham, D.; Griffor, M.; Kamath, A.; Lee, S.E.; Madura, R.; McGuire, D.; Subashi, T.; Wasilko, D.; Watts, P.; Mylari, B.L.; Oates, P.J.; Adams, P.D.; Rath, V.L.  
Deposited on : 2003-06-07  
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	356	 88% 11% •
1	B	356	 87% 12% •
1	C	356	 88% 11% •
1	D	356	 88% 11% •

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorbitol dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	356	Total	C	N	O	S	Se	0	5	0
			2716	1724	473	498	11	10			
1	B	356	Total	C	N	O	S	Se	0	5	0
			2716	1724	473	498	11	10			
1	C	356	Total	C	N	O	S	Se	0	5	0
			2716	1724	473	498	11	10			
1	D	356	Total	C	N	O	S	Se	0	5	0
			2716	1724	473	498	11	10			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	MSE	MET	modified residue	UNP Q00796
A	65	MSE	MET	modified residue	UNP Q00796
A	107	MSE	MET	modified residue	UNP Q00796
A	185	MSE	MET	modified residue	UNP Q00796
A	194	MSE	MET	modified residue	UNP Q00796
A	238	GLN	LEU	SEE REMARK 999	UNP Q00796
A	278	MSE	MET	modified residue	UNP Q00796
A	309	MSE	MET	modified residue	UNP Q00796
A	346	MSE	MET	modified residue	UNP Q00796
B	38	MSE	MET	modified residue	UNP Q00796
B	65	MSE	MET	modified residue	UNP Q00796
B	107	MSE	MET	modified residue	UNP Q00796
B	185	MSE	MET	modified residue	UNP Q00796
B	194	MSE	MET	modified residue	UNP Q00796
B	238	GLN	LEU	SEE REMARK 999	UNP Q00796
B	278	MSE	MET	modified residue	UNP Q00796
B	309	MSE	MET	modified residue	UNP Q00796
B	346	MSE	MET	modified residue	UNP Q00796
C	38	MSE	MET	modified residue	UNP Q00796
C	65	MSE	MET	modified residue	UNP Q00796
C	107	MSE	MET	modified residue	UNP Q00796

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	185	MSE	MET	modified residue	UNP Q00796
C	194	MSE	MET	modified residue	UNP Q00796
C	238	GLN	LEU	SEE REMARK 999	UNP Q00796
C	278	MSE	MET	modified residue	UNP Q00796
C	309	MSE	MET	modified residue	UNP Q00796
C	346	MSE	MET	modified residue	UNP Q00796
D	38	MSE	MET	modified residue	UNP Q00796
D	65	MSE	MET	modified residue	UNP Q00796
D	107	MSE	MET	modified residue	UNP Q00796
D	185	MSE	MET	modified residue	UNP Q00796
D	194	MSE	MET	modified residue	UNP Q00796
D	238	GLN	LEU	SEE REMARK 999	UNP Q00796
D	278	MSE	MET	modified residue	UNP Q00796
D	309	MSE	MET	modified residue	UNP Q00796
D	346	MSE	MET	modified residue	UNP Q00796

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is 4-[2-(HYDROXYMETHYL)PYRIMIDIN-4-YL]-N,N-DIMETHYLPIPERAZINE-1-SULFONAMIDE (three-letter code: 572) (formula: C<sub>11</sub>H<sub>19</sub>N<sub>5</sub>O<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 20	C 11	N 5	O 3	S 1	0	0
4	B	1	Total 20	C 11	N 5	O 3	S 1	0	0
4	C	1	Total 20	C 11	N 5	O 3	S 1	0	0
4	D	1	Total 20	C 11	N 5	O 3	S 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	317	Total 317	O 317	0	0
5	B	290	Total 290	O 290	0	0
5	C	283	Total 283	O 283	0	0
5	D	323	Total 323	O 323	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: Sorbitol dehydrogenase




#### • Molecule 1: Sorbitol dehydrogenase



#### • Molecule 1: Sorbitol dehydrogenase



#### • Molecule 1: Sorbitol dehydrogenase



S308	M309	V315	E335	K338	K339	G340	L343	K344	L347	P356
------	------	------	------	------	------	------	------	------	------	------



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 62	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.81Å 134.81Å 225.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	57.84 – 2.00	Depositor
% Data completeness (in resolution range)	92.2 (57.84-2.00)	Depositor
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.183 , 0.211	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	12337	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 572, NAD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2765	0.69	0/3732
1	B	0.38	0/2765	0.68	0/3732
1	C	0.38	0/2765	0.68	0/3732
1	D	0.40	0/2765	0.69	0/3732
All	All	0.39	0/11060	0.69	0/14928

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2716	0	2768	54	0
1	B	2716	0	2768	52	0
1	C	2716	0	2768	49	0
1	D	2716	0	2768	49	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	44	0	26	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	44	0	26	1	0
3	C	44	0	26	1	0
3	D	44	0	26	0	0
4	A	20	0	19	0	0
4	B	20	0	19	1	0
4	C	20	0	19	1	0
4	D	20	0	19	0	0
5	A	317	0	0	8	0
5	B	290	0	0	6	0
5	C	283	0	0	4	0
5	D	323	0	0	7	0
All	All	12337	0	11252	174	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 174 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:172:GLY:HA2	1:B:309[A]:MSE:HE3	1.26	1.17
1:C:309[A]:MSE:HE3	1:D:172:GLY:HA2	1.26	1.15
1:C:197:ALA:CB	1:D:309[A]:MSE:HE1	1.77	1.15
1:C:172:GLY:HA2	1:D:309[A]:MSE:HE3	1.18	1.14
1:A:309[A]:MSE:HE3	1:B:172:GLY:HA2	1.13	1.12

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	359/356 (101%)	351 (98%)	8 (2%)	0	<b>100</b> <b>100</b>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	359/356 (101%)	349 (97%)	10 (3%)	0	100	100
1	C	359/356 (101%)	349 (97%)	10 (3%)	0	100	100
1	D	359/356 (101%)	348 (97%)	11 (3%)	0	100	100
All	All	1436/1424 (101%)	1397 (97%)	39 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/282 (105%)	288 (98%)	7 (2%)	44	47
1	B	295/282 (105%)	288 (98%)	7 (2%)	44	47
1	C	295/282 (105%)	289 (98%)	6 (2%)	50	55
1	D	295/282 (105%)	290 (98%)	5 (2%)	56	61
All	All	1180/1128 (105%)	1155 (98%)	25 (2%)	48	53

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	60	ILE
1	C	274	LEU
1	D	299	TYR
1	C	127	ASN
1	C	283	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	127	ASN
1	C	355	ASN
1	C	127	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	238	GLN
1	C	238	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAD	B	406	-	42,48,48	2.45	11 (26%)	50,73,73	1.87	6 (12%)
4	572	B	407	2	20,21,21	4.80	9 (45%)	26,30,30	1.76	5 (19%)
3	NAD	A	404	-	42,48,48	2.47	11 (26%)	50,73,73	1.81	8 (16%)
4	572	D	411	2	20,21,21	4.85	9 (45%)	26,30,30	1.82	7 (26%)
4	572	A	405	2	20,21,21	4.83	9 (45%)	26,30,30	1.81	7 (26%)
3	NAD	D	410	-	42,48,48	2.47	11 (26%)	50,73,73	1.81	8 (16%)
4	572	C	409	2	20,21,21	4.78	10 (50%)	26,30,30	1.74	5 (19%)
3	NAD	C	408	-	42,48,48	2.43	10 (23%)	50,73,73	1.85	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	B	406	-	-	5/26/62/62	0/5/5/5
4	572	B	407	2	-	7/18/28/28	0/2/2/2
3	NAD	A	404	-	-	6/26/62/62	0/5/5/5
4	572	D	411	2	-	7/18/28/28	0/2/2/2
4	572	A	405	2	-	7/18/28/28	0/2/2/2
3	NAD	D	410	-	-	5/26/62/62	0/5/5/5
4	572	C	409	2	-	7/18/28/28	0/2/2/2
3	NAD	C	408	-	-	5/26/62/62	0/5/5/5

The worst 5 of 80 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	409	572	O25-S28	11.78	1.64	1.42
4	B	407	572	O25-S28	11.66	1.64	1.42
4	D	411	572	O25-S28	11.52	1.64	1.42
4	A	405	572	O25-S28	11.49	1.64	1.42
4	D	411	572	O12-S28	10.92	1.63	1.42

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	406	NAD	C5N-C4N-C3N	-8.37	112.14	120.36
3	C	408	NAD	C5N-C4N-C3N	-8.25	112.25	120.36
3	D	410	NAD	C5N-C4N-C3N	-7.99	112.51	120.36
3	A	404	NAD	C5N-C4N-C3N	-7.98	112.52	120.36
3	D	410	NAD	C3N-C7N-N7N	4.26	122.99	117.74

There are no chirality outliers.

5 of 49 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	404	NAD	O4D-C1D-N1N-C2N
3	A	404	NAD	O4D-C1D-N1N-C6N
3	A	404	NAD	C2D-C1D-N1N-C2N
3	B	406	NAD	O4D-C1D-N1N-C2N
3	B	406	NAD	O4D-C1D-N1N-C6N

There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	406	NAD	1	0
4	B	407	572	1	0
4	C	409	572	1	0
3	C	408	NAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## Ligand 572 B 407




## Ligand NAD A 404









## Ligand 572 D 411



## Ligand 572 A 405







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.