



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 21, 2024 – 12:30 AM EDT

PDB ID : 1Q16  
Title : Crystal structure of Nitrate Reductase A, NarGHI, from Escherichia coli  
Authors : Bertero, M.G.; Strynadka, N.C.J.  
Deposited on : 2003-07-18  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1247	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>15%</div> <div>.</div> </div> </div>
2	B	512	<div> <div>2%</div> <div> <div></div> <div>90%</div> <div>8%</div> <div>..</div> </div> </div>
3	C	225	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>20%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SF4	B	1403	-	-	X	-
9	3PH	B	1310	X	-	-	-

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 17199 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Respiratory nitrate reductase 1 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1244	Total	C	N	O	S	0	0	0
			9878	6239	1731	1860	48			

- Molecule 2 is a protein called Respiratory nitrate reductase 1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	509	Total	C	N	O	S	0	0	0
			4050	2562	701	755	32			


- Molecule 3 is a protein called Respiratory nitrate reductase 1 gamma chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	224	Total	C	N	O	S	0	0	0
			1790	1187	302	287	14			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	FME	MET	modified residue	UNP P11350

- Molecule 4 is PHOSPHORIC ACID 4-(2-AMINO-4-OXO-3,4,5,6,-TETRAHYDRO-PTE RIDIN-6-YL)-2-HYDROXY-3,4-DIMERCAPTO-BUT-3-EN-YL ESTER GUANYLATE ESTER (three-letter code: MD1) (formula: C<sub>20</sub>H<sub>26</sub>N<sub>10</sub>O<sub>13</sub>P<sub>2</sub>S<sub>2</sub>).




- | Mol | Chain | Residues | Atoms           | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 5   | A     | 1        | Total Mo<br>1 1 | 0       | 0       |

-


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 8	Fe 4	S 4	0	0
6	B	1	Total 8	Fe 4	S 4	0	0
6	B	1	Total 8	Fe 4	S 4	0	0
6	B	1	Total 8	Fe 4	S 4	0	0

- Molecule 7 is (1S)-2-[[[(2S)-2,3-DIHYDROXYPROPYL]OXY}(HYDROXY)PHOSPHORYL]OXY}-1-[(PENTANOYLOXY)METHYL]ETHYL OCTANOATE (three-letter code: AGA) (formula: C<sub>19</sub>H<sub>36</sub>O<sub>10</sub>P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	O	P	0	0
			30	19	10	1		

- Molecule 8 is FE3-S4 CLUSTER (three-letter code: F3S) (formula:  $\text{Fe}_3\text{S}_4$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is 1,2-DIACYL-GLYCEROL-3-SN-PHOSPHATE (three-letter code: 3PH) (formula:  $C_{39}H_{77}O_8P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	O	P	0	0
			18	9	8	1		

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
10	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	710	Total	O	0	0
			710	710		
11	B	427	Total	O	0	0
			427	427		
11	C	76	Total	O	0	0
			76	76		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Respiratory nitrate reductase 1 alpha chain






- Molecule 2: Respiratory nitrate reductase 1 beta chain





● Molecule 3: Respiratory nitrate reductase 1 gamma chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	154.18Å 241.38Å 139.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.68 – 1.90 29.68 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.68-1.90) 99.6 (29.68-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.58 (at 1.91Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.202 , 0.230 0.191 , 0.220	Depositor DCC
$R_{free}$ test set	10078 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.7	Xtriage
Anisotropy	0.732	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 48.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	17199	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, F3S, MD1, 3PH, AGA, 6MO, HEM, FME

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/10138	0.64	11/13762 (0.1%)
2	B	0.33	0/4146	0.61	0/5609
3	C	0.32	0/1831	0.53	0/2476
All	All	0.32	0/16115	0.62	11/21847 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1163	HIS	C-N-CA	13.34	155.04	121.70
1	A	502	TYR	CB-CG-CD1	-10.86	114.48	121.00
1	A	502	TYR	CB-CG-CD2	10.19	127.11	121.00
1	A	501	ALA	C-N-CA	10.11	146.96	121.70
1	A	1163	HIS	O-C-N	-8.48	109.13	122.70

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1163	HIS	Peptide
1	A	501	ALA	Peptide,Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9878	0	9534	133	0
2	B	4050	0	3973	31	0
3	C	1790	0	1826	31	0
4	A	94	0	42	5	0
5	A	1	0	0	0	0
6	A	8	0	0	0	0
6	B	24	0	0	2	0
7	A	30	0	36	0	0
8	B	7	0	0	0	0
9	B	18	0	10	0	0
10	C	86	0	60	0	0
11	A	710	0	0	10	1
11	B	427	0	0	6	0
11	C	76	0	0	2	0
All	All	17199	0	15481	192	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 192 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:ASN:HD21	1:A:382:GLU:HB2	1.18	1.00
1:A:378:ASN:ND2	1:A:382:GLU:HB2	1.93	0.84
3:C:73:PRO:HG2	3:C:76:MET:HG3	1.57	0.84
1:A:863:PRO:HG2	1:A:864:LEU:HD22	1.60	0.83
2:B:503:ASP:HB3	11:B:1830:HOH:O	1.82	0.80

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:A:1448:HOH:O	11:A:1448:HOH:O[3_354]	0.98	1.22

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1242/1247 (100%)	1171 (94%)	61 (5%)	10 (1%)	16	8
2	B	507/512 (99%)	495 (98%)	12 (2%)	0	100	100
3	C	220/225 (98%)	216 (98%)	4 (2%)	0	100	100
All	All	1969/1984 (99%)	1882 (96%)	77 (4%)	10 (0%)	25	17

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	361	VAL
1	A	1166	GLU
1	A	501	ALA
1	A	578	VAL
1	A	190	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1041/1044 (100%)	1022 (98%)	19 (2%)	54	52
2	B	436/439 (99%)	427 (98%)	9 (2%)	48	45
3	C	185/186 (100%)	183 (99%)	2 (1%)	70	71
All	All	1662/1669 (100%)	1632 (98%)	30 (2%)	54	52

5 of 30 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1072	GLU
2	B	503	ASP
1	A	1233	LEU
3	C	81	LEU
2	B	183	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	919	ASN
1	A	1076	GLN
1	A	946	ASN
1	A	1098	HIS
1	A	387	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FME	C	1	3	8,9,10	1.31	2 (25%)	8,9,11	1.64	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FME	C	1	3	-	4/7/9/11	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1	FME	CB-CG	2.26	1.60	1.51
3	C	1	FME	CB-CA	-2.12	1.48	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	FME	CA-N-CN	2.89	127.27	122.82
3	C	1	FME	CB-CA-N	2.34	114.79	110.52

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	FME	N-CA-CB-CG
3	C	1	FME	O1-CN-N-CA
3	C	1	FME	C-CA-CB-CG
3	C	1	FME	CB-CA-N-CN

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 1 is monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	SF4	B	1404	2	0,12,12	-	-	-		
6	SF4	A	1401	1	0,12,12	-	-	-		
8	F3S	B	1405	2	0,9,9	-	-	-		
4	MD1	A	1301	5	39,51,51	3.66	9 (23%)	40,78,78	2.31	12 (30%)
10	HEM	C	807	3	42,50,50	2.03	17 (40%)	46,82,82	1.46	8 (17%)
9	3PH	B	1310	-	17,17,47	0.87	0	18,21,52	2.03	5 (27%)
6	SF4	B	1403	2	0,12,12	-	-	-		
6	SF4	B	1402	2	0,12,12	-	-	-		
10	HEM	C	806	3	42,50,50	1.95	13 (30%)	46,82,82	1.37	7 (15%)
4	MD1	A	1300	5	39,51,51	3.79	10 (25%)	40,78,78	2.41	12 (30%)
7	AGA	A	1309	-	29,29,29	0.74	1 (3%)	32,35,35	1.48	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	SF4	B	1404	2	-	-	0/6/5/5
6	SF4	A	1401	1	-	-	0/6/5/5
8	F3S	B	1405	2	-	-	0/3/3/3
4	MD1	A	1301	5	-	2/18/59/59	0/5/5/5
10	HEM	C	807	3	-	5/12/54/54	-
6	SF4	B	1403	2	-	-	0/6/5/5
9	3PH	B	1310	-	1/1/3/4	5/18/18/49	-
6	SF4	B	1402	2	-	-	0/6/5/5
10	HEM	C	806	3	-	6/12/54/54	-
4	MD1	A	1300	5	-	6/18/59/59	0/5/5/5
7	AGA	A	1309	-	-	8/34/34/34	-

The worst 5 of 50 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1300	MD1	C7-N8	19.98	1.50	1.27
4	A	1301	MD1	C7-N8	18.93	1.49	1.27
4	A	1300	MD1	C15-N17	5.72	1.42	1.33
4	A	1301	MD1	C6-N1	5.49	1.42	1.33
4	A	1301	MD1	C15-N17	5.33	1.42	1.33

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1300	MD1	C5-C6-N1	-6.48	114.76	123.42
4	A	1300	MD1	C2-N1-C6	6.47	124.96	115.96
4	A	1301	MD1	N3-C2-N1	-6.08	119.48	127.21
4	A	1300	MD1	C17-N17-C15	5.81	124.05	115.96
4	A	1301	MD1	C5-C6-N1	-5.71	115.79	123.42

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
9	B	1310	3PH	C2

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1300	MD1	C5'-O5'-PA-O3B
4	A	1300	MD1	C5'-O5'-PA-O2A
7	A	1309	AGA	O1-C1-C2-C3
7	A	1309	AGA	C3-O5-P1-O3
7	A	1309	AGA	C3-O5-P1-O6


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1301	MD1	2	0
6	B	1403	SF4	2	0
4	A	1300	MD1	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand MD1 A 1301





## Ligand HEM C 807









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1244/1247 (99%)	0.33	68 (5%) 32 33	12, 24, 42, 59	0
2	B	509/512 (99%)	-0.27	9 (1%) 67 70	11, 18, 31, 51	0
3	C	223/225 (99%)	0.46	13 (5%) 30 31	13, 24, 41, 55	0
All	All	1976/1984 (99%)	0.19	90 (4%) 38 39	11, 22, 41, 59	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	PHE	10.2
1	A	362	ASP	6.8
3	C	78	GLU	6.7
1	A	9	TYR	6.7
3	C	75	TRP	5.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	FME	C	1	10/11	0.89	0.12	37,44,53,54	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	3PH	B	1310	18/48	0.71	0.21	42,55,60,62	0
7	AGA	A	1309	30/30	0.90	0.14	19,23,55,60	0
4	MD1	A	1300	47/47	0.93	0.09	14,22,35,39	0
4	MD1	A	1301	47/47	0.94	0.07	16,21,29,34	0
5	6MO	A	1302	1/1	0.94	0.15	40,40,40,40	0
10	HEM	C	807	43/43	0.95	0.11	20,25,37,45	0
10	HEM	C	806	43/43	0.97	0.09	11,16,19,30	0
8	F3S	B	1405	7/7	0.98	0.04	13,14,15,15	0
6	SF4	B	1402	8/8	0.98	0.05	19,21,21,23	0
6	SF4	B	1404	8/8	0.98	0.04	17,19,20,21	0
6	SF4	A	1401	8/8	0.98	0.06	19,22,22,24	0
6	SF4	B	1403	8/8	0.99	0.03	13,13,14,15	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 3PH B 1310:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around AGA A 1309:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MD1 A 1300:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around MD1 A 1301:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM C 807:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers ⓘ

There are no such residues in this entry.