



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 12, 2024 – 11:36 AM EST

PDB ID : 3Q3G
Title : Crystal Structure of A-domain in complex with antibody
Authors : Mahalingam, B.; Xiong, J.P.; Arnaout, M.A.
Deposited on : 2010-12-21
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	220	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>16%</div> <div>.</div> </div> </div>
1	C	220	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>15%</div> <div>.</div> </div> </div>
1	F	220	<div> <div>18%</div> <div> <div></div> <div>83%</div> <div>15%</div> <div>.</div> </div> </div>
1	J	220	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>16%</div> <div>.</div> </div> </div>
2	B	224	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	224	<div><div></div><div>%</div><div>83%</div><div>15%</div><div>•</div></div>
2	H	224	<div><div></div><div>2%</div><div>84%</div><div>15%</div><div>•</div></div>
2	K	224	<div><div></div><div>2%</div><div>87%</div><div>13%</div><div></div></div>
3	E	190	<div><div></div><div>%</div><div>87%</div><div>11%</div><div>••</div></div>
3	G	190	<div><div></div><div>2%</div><div>84%</div><div>15%</div><div>••</div></div>
3	I	190	<div><div></div><div>27%</div><div>85%</div><div>14%</div><div>••</div></div>
3	L	190	<div><div></div><div>21%</div><div>83%</div><div>14%</div><div>••</div></div>

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 20257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Antibody Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	220	Total	C	N	O	S	0	0	0
			1718	1074	283	353	8			
1	A	220	Total	C	N	O	S	0	0	0
			1718	1074	283	353	8			
1	F	220	Total	C	N	O	S	0	0	0
			1718	1074	283	353	8			
1	J	220	Total	C	N	O	S	0	0	0
			1718	1074	283	353	8			

- Molecule 2 is a protein called Antibody Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	224	Total	C	N	O	S	0	0	0
			1689	1062	277	342	8			
2	B	224	Total	C	N	O	S	0	0	0
			1689	1062	277	342	8			
2	H	224	Total	C	N	O	S	0	0	0
			1689	1062	277	342	8			
2	K	224	Total	C	N	O	S	0	0	0
			1689	1062	277	342	8			

- Molecule 3 is a protein called Integrin alpha-M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	189	Total	C	N	O	S	0	0	0
			1531	975	271	282	3			
3	E	189	Total	C	N	O	S	0	0	0
			1531	975	271	282	3			
3	I	189	Total	C	N	O	S	0	0	0
			1531	975	271	282	3			
3	L	189	Total	C	N	O	S	0	0	0
			1531	975	271	282	3			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	G	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	E	1	Total C O 4 2 2	0	0
4	F	1	Total C O 4 2 2	0	0
4	J	1	Total C O 4 2 2	0	0
4	J	1	Total C O 4 2 2	0	0
4	K	1	Total C O 4 2 2	0	0
4	K	1	Total C O 4 2 2	0	0
4	K	1	Total C O 4 2 2	0	0
4	K	1	Total C O 4 2 2	0	0
4	L	1	Total C O 4 2 2	0	0
4	L	1	Total C O 4 2 2	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0
5	G	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	J	1	Total C O 6 3 3	0	0
5	K	1	Total C O 6 3 3	0	0
5	K	1	Total C O 6 3 3	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	D	1	Total Na 1 1	0	0
6	H	2	Total Na 2 2	0	0


- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	G	1	Total Ca 1 1	0	0
7	E	1	Total Ca 1 1	0	0
7	I	1	Total Ca 1 1	0	0
7	L	1	Total Ca 1 1	0	0

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	1	Total	Cl	0	0
			1	1		
8	E	2	Total	Cl	0	0
			2	2		
8	J	1	Total	Cl	0	0
			1	1		
8	K	1	Total	Cl	0	0
			1	1		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	H	1	Total	C	O	0	0
			7	4	3		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	C	42	Total	O	0	0
			42	42		
10	D	42	Total	O	0	0
			42	42		
10	G	29	Total	O	0	0
			29	29		
10	A	31	Total	O	0	0
			31	31		
10	B	49	Total	O	0	0
			49	49		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	E	12	Total 12	O 12	0	0
10	F	6	Total 6	O 6	0	0
10	H	22	Total 22	O 22	0	0
10	I	6	Total 6	O 6	0	0
10	J	31	Total 31	O 31	0	0
10	K	40	Total 40	O 40	0	0
10	L	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Antibody Light Chain




• Molecule 1: Antibody Light Chain




• Molecule 1: Antibody Light Chain





• Molecule 1: Antibody Light Chain









• Molecule 2: Antibody Heavy chain




• Molecule 2: Antibody Heavy chain




• Molecule 2: Antibody Heavy chain




• Molecule 2: Antibody Heavy chain




• Molecule 3: Integrin alpha-M









• Molecule 3: Integrin alpha-M



• Molecule 3: Integrin alpha-M



• Molecule 3: Integrin alpha-M



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	81.94Å 157.22Å 232.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.76 – 2.70 47.76 – 2.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.76-2.70) 97.2 (47.76-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.211 , 0.244 0.221 , 0.254	Depositor DCC
R_{free} test set	4048 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	47.3	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	20257	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO, GOL, NA, CL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1757	0.71	1/2384 (0.0%)
1	C	0.44	0/1757	0.69	0/2384
1	F	0.40	0/1757	0.68	0/2384
1	J	0.44	0/1757	0.74	0/2384
2	B	0.44	0/1732	0.74	0/2368
2	D	0.45	0/1732	0.75	0/2368
2	H	0.44	0/1732	0.74	0/2368
2	K	0.45	0/1732	0.74	0/2368
3	E	0.44	0/1560	0.71	1/2099 (0.0%)
3	G	0.46	0/1560	0.70	0/2099
3	I	0.45	0/1560	0.67	0/2099
3	L	0.47	0/1560	0.70	0/2099
All	All	0.44	0/20196	0.71	2/27404 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	GLU	C-N-CA	5.51	135.47	121.70
3	E	167	SER	N-CA-C	-5.27	96.76	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1718	0	1647	14	0
1	C	1718	0	1647	14	0
1	F	1718	0	1647	14	0
1	J	1718	0	1647	14	0
2	B	1689	0	1638	21	0
2	D	1689	0	1638	21	0
2	H	1689	0	1638	20	0
2	K	1689	0	1638	18	0
3	E	1531	0	1545	6	0
3	G	1531	0	1545	14	0
3	I	1531	0	1545	10	0
3	L	1531	0	1545	16	0
4	A	8	0	12	0	0
4	B	16	0	24	3	0
4	C	8	0	12	0	0
4	D	20	0	30	3	0
4	E	8	0	12	1	0
4	F	4	0	6	0	0
4	G	4	0	6	0	0
4	J	8	0	12	0	0
4	K	16	0	24	0	0
4	L	8	0	12	0	0
5	A	6	0	8	1	0
5	B	12	0	16	1	0
5	C	6	0	8	0	0
5	D	18	0	24	1	0
5	G	6	0	8	1	0
5	J	6	0	8	0	0
5	K	12	0	16	1	0
6	D	1	0	0	0	0
6	H	2	0	0	0	0
7	E	1	0	0	0	0
7	G	1	0	0	0	0
7	I	1	0	0	0	0
7	L	1	0	0	0	0
8	E	2	0	0	0	0
8	G	1	0	0	0	0
8	J	1	0	0	0	0
8	K	1	0	0	0	0
9	H	7	0	10	0	0
10	A	31	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	B	49	0	0	0	0
10	C	42	0	0	1	0
10	D	42	0	0	0	0
10	E	12	0	0	0	0
10	F	6	0	0	1	0
10	G	29	0	0	0	0
10	H	22	0	0	0	0
10	I	6	0	0	0	0
10	J	31	0	0	0	0
10	K	40	0	0	0	0
10	L	10	0	0	1	0
All	All	20257	0	19568	174	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 174 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:199:ARG:HG3	2:B:199:ARG:HH11	1.00	1.12
2:D:199:ARG:HH11	2:D:199:ARG:HG3	1.09	1.10
1:C:13:VAL:HG22	1:C:19:VAL:HG11	1.41	1.02
2:K:199:ARG:HH11	2:K:199:ARG:HG3	1.23	1.00
2:H:199:ARG:HG2	2:H:200:PRO:HA	1.43	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	218/220 (99%)	212 (97%)	4 (2%)	2 (1%)	14 35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	218/220 (99%)	212 (97%)	5 (2%)	1 (0%)	25	49
1	F	218/220 (99%)	211 (97%)	7 (3%)	0	100	100
1	J	218/220 (99%)	211 (97%)	6 (3%)	1 (0%)	25	49
2	B	222/224 (99%)	210 (95%)	10 (4%)	2 (1%)	14	35
2	D	222/224 (99%)	212 (96%)	6 (3%)	4 (2%)	7	18
2	H	222/224 (99%)	211 (95%)	10 (4%)	1 (0%)	25	49
2	K	222/224 (99%)	213 (96%)	8 (4%)	1 (0%)	25	49
3	E	187/190 (98%)	178 (95%)	7 (4%)	2 (1%)	12	30
3	G	187/190 (98%)	179 (96%)	7 (4%)	1 (0%)	25	49
3	I	187/190 (98%)	180 (96%)	5 (3%)	2 (1%)	12	30
3	L	187/190 (98%)	178 (95%)	8 (4%)	1 (0%)	25	49
All	All	2508/2536 (99%)	2407 (96%)	83 (3%)	18 (1%)	19	42

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	225	ASP
3	E	168	LYS
2	H	141	ALA
1	J	46	PRO
2	B	142	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/197 (100%)	180 (91%)	17 (9%)	8	21
1	C	197/197 (100%)	183 (93%)	14 (7%)	12	30
1	F	197/197 (100%)	179 (91%)	18 (9%)	7	19
1	J	197/197 (100%)	175 (89%)	22 (11%)	5	12
2	B	192/192 (100%)	181 (94%)	11 (6%)	17	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	D	192/192 (100%)	181 (94%)	11 (6%)	17	40
2	H	192/192 (100%)	184 (96%)	8 (4%)	25	53
2	K	192/192 (100%)	183 (95%)	9 (5%)	22	49
3	E	169/169 (100%)	156 (92%)	13 (8%)	10	26
3	G	169/169 (100%)	163 (96%)	6 (4%)	30	59
3	I	169/169 (100%)	159 (94%)	10 (6%)	16	38
3	L	169/169 (100%)	160 (95%)	9 (5%)	19	43
All	All	2232/2232 (100%)	2084 (93%)	148 (7%)	14	33

5 of 148 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	39	LEU
3	L	242	ASP
1	J	84	VAL
2	K	30	ASN
2	B	13	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 47 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	144	ASN
3	I	311	GLN
2	H	175	HIS
3	I	224	ASN
1	J	35	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 49 ligands modelled in this entry, 12 are monoatomic - leaving 37 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	K	611	-	3,3,3	0.52	0	2,2,2	0.22	0
4	EDO	A	618	-	3,3,3	0.46	0	2,2,2	0.39	0
4	EDO	J	603	-	3,3,3	0.62	0	2,2,2	0.10	0
5	GOL	G	703	-	5,5,5	1.02	0	5,5,5	1.21	0
5	GOL	C	710	-	5,5,5	1.35	1 (20%)	5,5,5	1.42	1 (20%)
4	EDO	J	622	-	3,3,3	0.54	0	2,2,2	0.13	0
4	EDO	L	615	-	3,3,3	0.43	0	2,2,2	0.41	0
4	EDO	D	619	-	3,3,3	0.47	0	2,2,2	0.38	0
4	EDO	G	620	-	3,3,3	0.42	0	2,2,2	0.43	0
4	EDO	F	621	-	3,3,3	0.55	0	2,2,2	0.24	0
5	GOL	D	702	-	5,5,5	0.77	0	5,5,5	1.03	1 (20%)
4	EDO	D	604	-	3,3,3	0.62	0	2,2,2	0.07	0
5	GOL	J	709	-	5,5,5	1.08	0	5,5,5	0.97	0
4	EDO	B	607	-	3,3,3	0.59	0	2,2,2	0.11	0
4	EDO	D	613	-	3,3,3	0.52	0	2,2,2	0.18	0
4	EDO	K	608	-	3,3,3	0.48	0	2,2,2	0.32	0
4	EDO	E	600	-	3,3,3	0.48	0	2,2,2	0.29	0
5	GOL	B	704	-	5,5,5	0.80	0	5,5,5	0.45	0
4	EDO	D	614	-	3,3,3	0.41	0	2,2,2	0.56	0
5	GOL	K	706	-	5,5,5	0.77	0	5,5,5	0.86	0
5	GOL	D	705	-	5,5,5	1.08	0	5,5,5	0.69	0
5	GOL	K	707	-	5,5,5	0.62	0	5,5,5	0.85	0
5	GOL	B	701	-	5,5,5	0.66	0	5,5,5	0.63	0
4	EDO	B	616	-	3,3,3	0.50	0	2,2,2	0.15	0
9	PEG	H	1000	-	6,6,6	0.46	0	5,5,5	0.57	0
4	EDO	B	606	-	3,3,3	0.46	0	2,2,2	0.43	0
4	EDO	C	609	-	3,3,3	0.55	0	2,2,2	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	625	-	3,3,3	0.42	0	2,2,2	0.35	0
4	EDO	K	612	-	3,3,3	0.57	0	2,2,2	0.27	0
4	EDO	D	605	-	3,3,3	0.50	0	2,2,2	0.24	0
4	EDO	L	610	-	3,3,3	0.50	0	2,2,2	0.32	0
5	GOL	D	708	-	5,5,5	1.21	0	5,5,5	0.77	0
5	GOL	A	700	-	5,5,5	0.80	0	5,5,5	0.80	0
4	EDO	C	628	-	3,3,3	0.46	0	2,2,2	0.51	0
4	EDO	K	601	-	3,3,3	0.48	0	2,2,2	0.39	0
4	EDO	E	627	-	3,3,3	0.47	0	2,2,2	0.34	0
4	EDO	B	624	-	3,3,3	0.47	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	K	611	-	-	1/1/1/1	-
4	EDO	A	618	-	-	0/1/1/1	-
4	EDO	J	603	-	-	1/1/1/1	-
5	GOL	G	703	-	-	3/4/4/4	-
5	GOL	C	710	-	-	0/4/4/4	-
4	EDO	J	622	-	-	0/1/1/1	-
4	EDO	L	615	-	-	1/1/1/1	-
4	EDO	D	619	-	-	1/1/1/1	-
4	EDO	G	620	-	-	1/1/1/1	-
4	EDO	F	621	-	-	1/1/1/1	-
5	GOL	D	702	-	-	2/4/4/4	-
4	EDO	D	604	-	-	1/1/1/1	-
5	GOL	J	709	-	-	2/4/4/4	-
4	EDO	B	607	-	-	1/1/1/1	-
4	EDO	D	613	-	-	0/1/1/1	-
4	EDO	K	608	-	-	0/1/1/1	-
4	EDO	E	600	-	-	0/1/1/1	-
5	GOL	B	704	-	-	0/4/4/4	-
4	EDO	D	614	-	-	0/1/1/1	-
5	GOL	K	706	-	-	2/4/4/4	-
5	GOL	D	705	-	-	4/4/4/4	-
5	GOL	K	707	-	-	2/4/4/4	-
5	GOL	B	701	-	-	0/4/4/4	-
4	EDO	B	616	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PEG	H	1000	-	-	2/4/4/4	-
4	EDO	B	606	-	-	0/1/1/1	-
4	EDO	C	609	-	-	1/1/1/1	-
4	EDO	A	625	-	-	0/1/1/1	-
4	EDO	K	612	-	-	0/1/1/1	-
4	EDO	D	605	-	-	1/1/1/1	-
4	EDO	L	610	-	-	0/1/1/1	-
5	GOL	D	708	-	-	2/4/4/4	-
5	GOL	A	700	-	-	0/4/4/4	-
4	EDO	C	628	-	-	0/1/1/1	-
4	EDO	K	601	-	-	1/1/1/1	-
4	EDO	E	627	-	-	0/1/1/1	-
4	EDO	B	624	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	710	GOL	O2-C2	2.03	1.49	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	710	GOL	O2-C2-C3	2.78	120.69	109.18
5	D	702	GOL	O1-C1-C2	2.20	120.30	110.38

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	702	GOL	O1-C1-C2-C3
5	D	705	GOL	O1-C1-C2-C3
5	D	705	GOL	C1-C2-C3-O3
5	D	708	GOL	O1-C1-C2-C3
5	J	709	GOL	O1-C1-C2-C3

There are no ring outliers.

9 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	703	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	702	GOL	1	0
4	D	604	EDO	2	0
4	B	607	EDO	3	0
4	E	600	EDO	1	0
5	B	704	GOL	1	0
4	D	614	EDO	1	0
5	K	707	GOL	1	0
5	A	700	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	220/220 (100%)	-0.04	2 (0%) 81 80	25, 42, 66, 94	0
1	C	220/220 (100%)	0.04	5 (2%) 61 60	26, 41, 80, 113	0
1	F	220/220 (100%)	1.26	40 (18%) 4 4	38, 74, 97, 112	0
1	J	220/220 (100%)	0.07	2 (0%) 81 80	26, 46, 66, 110	0
2	B	224/224 (100%)	-0.12	4 (1%) 67 67	24, 39, 73, 105	0
2	D	224/224 (100%)	-0.21	2 (0%) 81 80	21, 37, 70, 113	0
2	H	224/224 (100%)	0.08	4 (1%) 67 67	28, 47, 77, 111	0
2	K	224/224 (100%)	0.07	4 (1%) 67 67	24, 41, 74, 111	0
3	E	189/190 (99%)	0.08	2 (1%) 77 77	28, 53, 76, 115	0
3	G	189/190 (99%)	-0.09	3 (1%) 70 70	29, 44, 73, 104	0
3	I	189/190 (99%)	1.49	52 (27%) 2 2	42, 87, 116, 125	0
3	L	189/190 (99%)	1.19	39 (20%) 3 3	30, 73, 112, 122	0
All	All	2532/2536 (99%)	0.30	159 (6%) 27 25	21, 48, 98, 125	0

The worst 5 of 159 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	K	4	VAL	4.7
3	L	297	PHE	4.0
1	F	86	ALA	3.9
3	L	307	THR	3.9
3	I	319	ILE	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	B	616	4/4	0.65	0.25	63,64,64,65	0
4	EDO	D	614	4/4	0.69	0.18	49,49,50,51	0
4	EDO	A	618	4/4	0.71	0.18	57,59,62,62	0
5	GOL	C	710	6/6	0.74	0.17	53,56,57,58	0
5	GOL	D	702	6/6	0.74	0.19	61,63,64,64	0
4	EDO	K	608	4/4	0.75	0.14	52,56,56,60	0
4	EDO	L	610	4/4	0.75	0.16	57,58,58,65	0
4	EDO	K	611	4/4	0.77	0.18	60,62,62,65	0
5	GOL	D	705	6/6	0.77	0.17	59,61,61,62	0
5	GOL	A	700	6/6	0.78	0.17	68,69,70,71	0
5	GOL	B	701	6/6	0.78	0.14	71,71,72,73	0
8	CL	J	903	1/1	0.78	0.21	70,70,70,70	0
5	GOL	D	708	6/6	0.79	0.15	50,50,51,51	0
4	EDO	J	622	4/4	0.79	0.18	55,56,57,66	0
4	EDO	F	621	4/4	0.80	0.18	56,57,58,62	0
5	GOL	K	707	6/6	0.80	0.15	66,68,68,69	0
4	EDO	D	619	4/4	0.80	0.21	53,55,57,60	0
9	PEG	H	1000	7/7	0.80	0.16	59,59,62,63	0
5	GOL	K	706	6/6	0.81	0.15	60,62,62,62	0
4	EDO	L	615	4/4	0.81	0.11	51,54,55,55	0
6	NA	H	801	1/1	0.82	0.17	37,37,37,37	0
4	EDO	C	609	4/4	0.82	0.18	54,54,56,57	0
5	GOL	J	709	6/6	0.82	0.14	59,60,61,62	0
4	EDO	B	607	4/4	0.83	0.17	47,50,51,55	0
4	EDO	K	612	4/4	0.83	0.12	39,40,41,42	0
4	EDO	D	605	4/4	0.84	0.16	49,50,50,55	0
5	GOL	B	704	6/6	0.84	0.19	54,56,56,57	0
4	EDO	A	625	4/4	0.85	0.13	41,43,45,45	0
4	EDO	D	604	4/4	0.86	0.14	37,40,41,48	0
4	EDO	B	624	4/4	0.86	0.14	55,55,56,64	0
8	CL	G	907	1/1	0.86	0.11	74,74,74,74	0
4	EDO	E	600	4/4	0.86	0.17	42,43,48,49	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	E	627	4/4	0.86	0.14	54,57,60,62	0
4	EDO	B	606	4/4	0.87	0.11	43,43,47,48	0
5	GOL	G	703	6/6	0.87	0.11	48,48,49,49	0
4	EDO	C	628	4/4	0.87	0.13	50,50,52,56	0
4	EDO	J	603	4/4	0.88	0.12	40,42,45,45	0
4	EDO	D	613	4/4	0.88	0.13	51,51,55,58	0
8	CL	E	902	1/1	0.91	0.06	60,60,60,60	0
4	EDO	G	620	4/4	0.91	0.11	38,38,39,46	0
8	CL	K	904	1/1	0.91	0.08	63,63,63,63	0
4	EDO	K	601	4/4	0.91	0.11	40,40,41,46	0
6	NA	D	800	1/1	0.92	0.17	41,41,41,41	0
6	NA	H	802	1/1	0.93	0.27	58,58,58,58	0
8	CL	E	906	1/1	0.97	0.07	71,71,71,71	0
7	CA	L	500	1/1	0.97	0.04	40,40,40,40	0
7	CA	E	500	1/1	0.97	0.03	40,40,40,40	0
7	CA	I	500	1/1	0.97	0.04	57,57,57,57	0
7	CA	G	500	1/1	0.99	0.03	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.