



Full wwPDB EM Validation Report ⓘ

Apr 1, 2025 – 09:24 pm BST

PDB ID : 6Q6I / pdb_00006q6i
EMDB ID : EMD-4468
Title : Lysine decarboxylase A from *Pseudomonas aeruginosa*
Authors : Kandiah, E.; Gutsche, I.
Deposited on : 2018-12-11
Resolution : 3.70 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev117
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	751	<div> <div>24%</div> <div>78%</div> <div>21%</div> </div>

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 5720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Biodegradative arginine decarboxylase.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	749	5705	3632	998	1052	23	0	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
2	A	1	15	8	1	5	1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Biodegradative arginine decarboxylase



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, D5	Depositor
Number of particles used	66193	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	40	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.042	Depositor
Minimum map value	-0.025	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.007	Depositor
Map size (Å)	303.369, 303.369, 303.369	wwPDB
Map dimensions	372, 372, 372	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.815508, 0.815508, 0.815508	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/5840	0.66	4/7942 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	522	LEU	CA-CB-CG	8.22	134.21	115.30
1	A	476	LEU	CA-CB-CG	6.58	130.42	115.30
1	A	371	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	392	HIS	CA-C-N	-5.62	104.83	117.20

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	127	ASP	Peptide
1	A	163	ASN	Peptide
1	A	203	VAL	Peptide
1	A	221	ASP	Peptide
1	A	392	HIS	Mainchain
1	A	395	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5705	0	5561	91	0
2	A	15	0	6	2	0
All	All	5720	0	5567	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:HIS:HB3	1:A:18:ALA:HB2	1.76	0.68
1:A:392:HIS:NE2	2:A:801:PLP:O3P	2.30	0.64
1:A:264:HIS:ND1	1:A:265:LYS:N	2.46	0.63
1:A:365:HIS:HB2	1:A:455:ILE:HD11	1.79	0.62
1:A:323:LEU:H	1:A:351:VAL:HG23	1.67	0.59
1:A:674:ILE:HD11	1:A:744:TYR:HB2	1.82	0.59
1:A:533:PRO:HD2	1:A:546:ILE:HD11	1.86	0.58
1:A:247:TRP:HA	1:A:323:LEU:HD21	1.85	0.57
1:A:87:ARG:NH1	1:A:91:PRO:O	2.38	0.56
1:A:128:THR:O	1:A:132:LEU:N	2.34	0.56
1:A:144:LEU:HD23	1:A:147:LEU:HD21	1.86	0.56
1:A:222:ARG:NH2	1:A:447:GLU:OE1	2.38	0.55
1:A:337:ASN:HB3	1:A:340:LEU:HB2	1.89	0.54
1:A:174:GLY:O	1:A:178:ARG:NH1	2.40	0.54
1:A:378:ARG:HD3	1:A:407:GLN:HB2	1.91	0.53
1:A:390:SER:O	1:A:390:SER:OG	2.26	0.53
1:A:91:PRO:HG2	1:A:92:GLN:HE21	1.74	0.53
1:A:642:ALA:HA	1:A:645:ARG:HB2	1.90	0.52
1:A:729:ASP:OD1	1:A:729:ASP:N	2.37	0.52
1:A:275:GLY:HA2	1:A:658:SER:HB2	1.91	0.52
1:A:555:LEU:HA	1:A:558:ARG:HB2	1.92	0.52
1:A:206:LEU:HD23	1:A:432:GLN:HB3	1.92	0.52
1:A:45:SER:OG	1:A:46:SER:N	2.43	0.52
1:A:550:ILE:HG23	1:A:607:LEU:HD11	1.92	0.52
1:A:98:LEU:HD23	1:A:122:LEU:HB2	1.91	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:HIS:CG	1:A:265:LYS:H	2.29	0.51
1:A:256:LEU:HB3	1:A:279:LEU:HD11	1.91	0.51
1:A:197:SER:O	1:A:197:SER:OG	2.25	0.50
1:A:337:ASN:HA	1:A:371:ARG:HD3	1.92	0.50
1:A:207:GLY:O	1:A:432:GLN:NE2	2.45	0.50
1:A:244:LYS:O	1:A:248:HIS:HB2	2.12	0.49
1:A:329:SER:O	1:A:329:SER:OG	2.28	0.49
1:A:572:VAL:HG21	1:A:586:LEU:HD21	1.94	0.49
1:A:20:THR:O	1:A:27:ARG:NH2	2.35	0.49
1:A:187:HIS:HE1	1:A:192:GLU:HG3	1.78	0.48
1:A:467:ARG:HB2	1:A:493:GLY:HA2	1.96	0.48
1:A:520:TYR:HB3	1:A:521:VAL:HG22	1.95	0.48
1:A:681:ILE:HG12	1:A:701:ARG:HB2	1.94	0.48
1:A:347:ASP:N	1:A:347:ASP:OD1	2.46	0.48
1:A:627:SER:O	1:A:631:HIS:N	2.40	0.48
1:A:264:HIS:CG	1:A:265:LYS:N	2.82	0.48
1:A:545:GLY:O	1:A:627:SER:OG	2.32	0.48
1:A:501:ASP:OD1	1:A:501:ASP:N	2.45	0.47
1:A:285:ARG:HG2	1:A:292:GLY:H	1.80	0.47
1:A:320:LYS:NZ	1:A:347:ASP:O	2.39	0.47
1:A:368:TYR:OH	1:A:524:ASP:OD1	2.27	0.47
1:A:226:ASN:HD21	1:A:447:GLU:HG2	1.80	0.47
1:A:308:ILE:HD13	1:A:318:GLU:HA	1.96	0.47
1:A:352:LEU:HB3	1:A:385:VAL:HA	1.96	0.47
1:A:431:PRO:HB2	1:A:432:GLN:HE21	1.80	0.46
1:A:240:SER:N	2:A:801:PLP:O1P	2.38	0.46
1:A:356:GLU:HB3	1:A:359:TYR:HB3	1.96	0.46
1:A:653:VAL:HG22	1:A:697:MET:HE1	1.98	0.46
1:A:434:GLY:O	1:A:438:SER:N	2.37	0.46
1:A:313:LEU:HD23	1:A:656:ARG:HG3	1.98	0.46
1:A:446:MET:HE1	1:A:454:LEU:HD23	1.98	0.45
1:A:695:LEU:HD21	1:A:709:ILE:HG21	1.98	0.45
1:A:695:LEU:HD23	1:A:696:ILE:HG23	1.97	0.45
1:A:210:LEU:HD23	1:A:424:MET:HG2	1.97	0.45
1:A:671:ALA:HB1	1:A:745:THR:HG21	1.98	0.45
1:A:738:GLY:HA3	1:A:739:PRO:HD3	1.86	0.45
1:A:236:ILE:HB	1:A:427:ILE:HD11	1.99	0.45
1:A:171:HIS:HD2	1:A:174:GLY:H	1.65	0.44
1:A:168:THR:HG23	1:A:169:PRO:HD3	1.99	0.44
1:A:686:LEU:HD12	1:A:696:ILE:HD11	2.00	0.44
1:A:17:LYS:HE3	1:A:17:LYS:HB2	1.84	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:401:ALA:HB1	1:A:439:LEU:HD13	2.01	0.43
1:A:340:LEU:O	1:A:344:THR:OG1	2.32	0.43
1:A:141:ARG:HA	1:A:144:LEU:HB2	2.00	0.43
1:A:548:ALA:O	1:A:564:LYS:NZ	2.44	0.43
1:A:548:ALA:HB1	1:A:564:LYS:HD2	2.02	0.42
1:A:8:PRO:HG2	1:A:59:LEU:HD23	2.01	0.42
1:A:541:LEU:HD23	1:A:631:HIS:CE1	2.54	0.42
1:A:628:ASP:HA	1:A:631:HIS:HB3	2.01	0.42
1:A:687:VAL:HG22	1:A:694:PRO:HA	2.00	0.42
1:A:14:ARG:O	1:A:18:ALA:N	2.52	0.42
1:A:151:PHE:HB2	1:A:219:ALA:HB2	2.01	0.42
1:A:235:VAL:HG22	1:A:423:PHE:HE1	1.84	0.42
1:A:212:HIS:NE2	1:A:420:ASN:OD1	2.53	0.42
1:A:239:THR:HA	1:A:242:ALA:HB3	2.01	0.42
1:A:246:VAL:HG13	1:A:404:ILE:HD13	2.03	0.41
1:A:645:ARG:HD3	1:A:645:ARG:HA	1.88	0.41
1:A:26:VAL:HA	1:A:29:ILE:HG12	2.03	0.41
1:A:542:SER:O	1:A:631:HIS:NE2	2.36	0.41
1:A:750:LYS:HA	1:A:750:LYS:HD3	1.90	0.41
1:A:267:ILE:O	1:A:271:ILE:HG13	2.21	0.41
1:A:337:ASN:O	1:A:341:ILE:HG12	2.21	0.41
1:A:33:LEU:HD23	1:A:33:LEU:HA	1.94	0.40
1:A:213:THR:OG1	1:A:214:GLY:N	2.55	0.40
1:A:311:SER:HA	1:A:312:PRO:HD3	1.95	0.40
1:A:483:PHE:CD1	1:A:532:THR:HG22	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	747/751 (100%)	620 (83%)	125 (17%)	2 (0%)	37 67

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	PRO
1	A	490	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	581/612 (95%)	571 (98%)	10 (2%)	56	73

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	259	VAL
1	A	351	VAL
1	A	361	TYR
1	A	393	LYS
1	A	398	PHE
1	A	521	VAL
1	A	546	ILE
1	A	555	LEU
1	A	584	SER
1	A	641	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	GLN
1	A	343	GLN
1	A	365	HIS
1	A	456	GLN
1	A	544	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	A	801	1	15,15,16	2.10	4 (26%)	20,22,23	1.63	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	801	1	-	0/6/6/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	PLP	P-O4P	5.66	1.78	1.60
2	A	801	PLP	O4P-C5A	-3.43	1.32	1.45
2	A	801	PLP	C5-C4	-2.91	1.37	1.40
2	A	801	PLP	C6-C5	2.00	1.41	1.37

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	PLP	O3P-P-O4P	-3.52	97.37	106.73
2	A	801	PLP	O4P-P-O1P	-2.35	99.88	106.47
2	A	801	PLP	O3P-P-O2P	2.33	116.55	107.64
2	A	801	PLP	O2P-P-O1P	2.19	119.25	110.68
2	A	801	PLP	O3-C3-C4	2.15	123.76	118.10
2	A	801	PLP	O2P-P-O4P	-2.08	101.19	106.73
2	A	801	PLP	C5A-C5-C6	-2.06	115.98	119.37

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	PLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-4468. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 186



Y Index: 186



Z Index: 186


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 164



Y Index: 229




Z Index: 215


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.007. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 252 nm^3 ; this corresponds to an approximate mass of 228 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.270 Å⁻¹

8 Fourier-Shell correlation ⓘ


This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-4468 and PDB model 6Q6I. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlays

9.1.1 Map-model overlay [i](#)




9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.007 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.007).

9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.007) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.5840	<div></div> 0.4280
A	<div></div> 0.5840	<div></div> 0.4280

