



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 03:21 AM EDT

PDB ID : 4QF1  
Title : Crystal structure of unliganded CH59UA, the inferred unmutated ancestor of the RV144 anti-HIV antibody lineage producing CH59  
Authors : Wiehe, K.; Easterhoff, D.; Luo, K.; Nicely, N.I.; Bradley, T.; Jaeger, F.H.; Denison, S.M.; Zhang, R.; Lloyd, K.E.; Stolarчук, C.; Parks, R.; Sutherland, L.L.; Searce, R.M.; Morris, L.; Kaewkungwal, J.; Nitayaphan, S.; Pitisutthum, P.; Rerks-Ngarm, S.; Michael, N.; Kim, J.; Kelsoe, G.; Montefiori, D.C.; Tomaras, G.; Bonsignori, M.; Santra, S.; Kepler, T.B.; Alam, S.M.; Moody, M.A.; Liao, H.-X.; Haynes, B.F.  
Deposited on : 2014-05-19  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	 4% 84% 10% 6%
1	H	225	 4% 82% 12% 6%
2	B	214	 3% 90% 8% .

*Continued on next page...*

Ideal geometry (proteins) : Engh & Huber (2001)  
 Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
 Validation Pipeline (wwPDB-VP) : 2.37.1

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	L	214	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	B	303	-	-	X	-

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6642 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CH59UA Fab fragment of heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	212	Total	C	N	O	S	0	0	0
			1598	1010	271	311	6			
1	A	212	Total	C	N	O	S	0	0	0
			1598	1010	271	311	6			

- Molecule 2 is a protein called Inferred unmutated ancestor (UA) of anti-HIV antibody CH59.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	1	0
			1594	998	266	325	5			
2	B	210	Total	C	N	O	S	0	0	0
			1579	989	262	323	5			

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	C	O	0	0
			10	6	4		
4	B	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	1	Total	Cl	0	0
			1	1		
5	B	1	Total	Cl	0	0
			1	1		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	48	Total	O	0	0
			48	48		
7	L	48	Total	O	0	0
			48	48		
7	A	61	Total	O	0	0
			61	61		
7	B	44	Total	O	0	0
			44	44		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CH59UA Fab fragment of heavy chain




- Molecule 1: CH59UA Fab fragment of heavy chain



- Molecule 2: Inferred unmutated ancestor (UA) of anti-HIV antibody CH59



- Molecule 2: Inferred unmutated ancestor (UA) of anti-HIV antibody CH59



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.33Å 70.80Å 107.31Å 90.00° 96.84° 90.00°	Depositor
Resolution (Å)	25.80 – 2.40 25.87 – 2.40	Depositor EDS
% Data completeness (in resolution range)	93.4 (25.80-2.40) 92.9 (25.87-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.06 (at 2.39Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928)	Depositor
R, $R_{free}$	0.182 , 0.229 0.175 , 0.221	Depositor DCC
$R_{free}$ test set	2000 reflections (4.48%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.0	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6642	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, CL, MES, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.36	0/1638	0.55	0/2232
1	H	0.37	0/1638	0.52	0/2232
2	B	0.34	0/1619	0.51	0/2209
2	L	0.35	0/1637	0.53	0/2234
All	All	0.36	0/6532	0.53	0/8907

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1598	0	1555	14	0
1	H	1598	0	1555	12	0
2	B	1579	0	1528	11	0
2	L	1594	0	1548	7	0
3	A	12	0	12	0	0
3	H	12	0	12	0	0
4	B	10	0	14	3	0
4	L	10	0	14	1	0
5	B	1	0	0	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	1	0	0	1	0
6	A	13	0	18	1	0
6	B	13	0	18	3	0
7	A	61	0	0	2	0
7	B	44	0	0	4	0
7	H	48	0	0	1	0
7	L	48	0	0	1	0
All	All	6642	0	6274	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:303:CL:CL	7:B:444:HOH:O	2.31	0.83
5:B:303:CL:CL	7:B:442:HOH:O	2.41	0.76
2:B:41:GLY:H	4:B:302:PGE:H62	1.53	0.74
2:L:31:LYS:HE2	4:L:301:PGE:H62	1.70	0.72
5:L:302:CL:CL	7:L:444:HOH:O	2.44	0.72
2:B:53:LYS:NZ	7:B:437:HOH:O	2.27	0.67
2:L:30:LYS:NZ	1:A:100:GLU:OE2	2.32	0.63
2:L:2:TYR:HD2	2:L:97:VAL:HG11	1.67	0.60
2:L:132:LEU:HD12	2:L:178:LEU:HD23	1.85	0.58
1:A:66:ARG:NH2	1:A:86:ASP:OD2	2.38	0.55
2:B:28:LEU:O	5:B:303:CL:CL	2.63	0.54
1:A:97:PRO:O	7:A:442:HOH:O	2.19	0.52
1:H:33:ALA:HB1	6:B:301:PG4:H71	1.92	0.52
6:A:301:PG4:H72	2:B:96:ARG:HG3	1.93	0.50
1:H:32:TYR:HE2	1:H:97:PRO:HD3	1.77	0.50
2:L:117:LEU:HD12	2:L:133:VAL:O	2.13	0.49
1:A:184:VAL:HG11	1:A:194:TYR:CE1	2.47	0.49
1:H:159:LEU:HD21	1:H:182:VAL:HG21	1.94	0.49
1:H:170:LEU:HD13	1:H:176:TYR:CZ	2.48	0.48
1:H:139:GLY:HA2	1:H:154:TRP:CH2	2.49	0.48
1:A:66:ARG:HD3	7:A:408:HOH:O	2.14	0.47
2:B:110:LYS:HG2	2:B:141:PRO:HD3	1.96	0.47
1:A:30:ASP:O	1:A:52(A):TRP:HB2	2.14	0.47
2:B:95(A):ASN:HD21	6:B:301:PG4:H72	1.78	0.47
4:B:302:PGE:H6	7:B:439:HOH:O	2.14	0.47
2:B:47:VAL:HG12	2:B:58:ILE:HD12	1.97	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:H:432:HOH:O	6:B:301:PG4:H81	2.16	0.45
1:A:135:THR:HG23	1:A:186:SER:H	1.81	0.45
1:H:13:GLN:H	1:H:16:ARG:HD3	1.81	0.45
1:H:30:ASP:O	1:H:52(A):TRP:HB2	2.16	0.45
1:A:22:CYS:HB3	1:A:78:LEU:HB3	1.98	0.45
2:L:54[B]:ARG:NH1	2:L:58:ILE:O	2.48	0.45
2:B:2:TYR:HD2	2:B:97:VAL:HG11	1.82	0.45
2:B:41:GLY:N	4:B:302:PGE:H62	2.28	0.44
2:B:119:PRO:HA	2:B:132:LEU:HD23	1.98	0.44
1:A:96:SER:HB3	1:A:100(A):LEU:HB2	2.00	0.44
1:A:14:PRO:HD2	1:A:113:SER:HB3	2.00	0.43
1:H:34:MET:HB3	1:H:78:LEU:HD22	1.99	0.43
1:H:12:VAL:HG11	1:H:18:LEU:HD13	2.00	0.43
1:A:64:LYS:HB3	1:A:64:LYS:HE3	1.81	0.43
1:H:121:VAL:HG12	1:H:209:LYS:HG3	2.02	0.42
1:A:167:PRO:HG2	2:B:165:SER:OG	2.20	0.42
1:A:119:PRO:HB3	1:A:145:TYR:HB3	2.01	0.41
1:H:96:SER:HB3	1:H:100(A):LEU:HB2	2.02	0.41
1:H:18:LEU:HB3	1:H:82:MET:CE	2.50	0.41
2:L:153:SER:HA	2:L:154:PRO:HD3	1.95	0.40
1:A:154:TRP:CH2	1:A:196:CYS:HB3	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/225 (92%)	202 (97%)	6 (3%)	0	100	100
1	H	208/225 (92%)	203 (98%)	5 (2%)	0	100	100
2	B	208/214 (97%)	204 (98%)	4 (2%)	0	100	100
2	L	210/214 (98%)	203 (97%)	6 (3%)	1 (0%)	29	41

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	834/878 (95%)	812 (97%)	21 (2%)	1 (0%)	51 68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	151	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	177/188 (94%)	174 (98%)	3 (2%)	60 78
1	H	177/188 (94%)	171 (97%)	6 (3%)	37 56
2	B	177/181 (98%)	174 (98%)	3 (2%)	60 78
2	L	179/181 (99%)	175 (98%)	4 (2%)	52 71
All	All	710/738 (96%)	694 (98%)	16 (2%)	50 70

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	19	ARG
1	H	30	ASP
1	H	64	LYS
1	H	135	THR
1	H	164	HIS
1	H	193	THR
2	L	31	LYS
2	L	97	VAL
2	L	144	VAL
2	L	205	THR
1	A	30	ASP
1	A	171	GLN
1	A	181	VAL
2	B	31	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	144	VAL
2	B	190	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	PG4	B	301	-	12,12,12	0.74	0	11,11,11	1.46	0
6	PG4	A	301	-	12,12,12	0.77	0	11,11,11	1.35	0
3	MES	A	302	-	12,12,12	2.21	1 (8%)	14,16,16	1.96	5 (35%)
3	MES	H	301	-	12,12,12	2.26	1 (8%)	14,16,16	1.91	3 (21%)
4	PGE	L	301	-	9,9,9	0.70	0	8,8,8	1.33	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	PGE	B	302	-	9,9,9	0.62	0	8,8,8	1.65	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PG4	B	301	-	-	7/10/10/10	-
6	PG4	A	301	-	-	6/10/10/10	-
3	MES	A	302	-	-	4/6/14/14	0/1/1/1
3	MES	H	301	-	-	4/6/14/14	0/1/1/1
4	PGE	L	301	-	-	5/7/7/7	-
4	PGE	B	302	-	-	2/7/7/7	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	301	MES	C8-S	-7.60	1.66	1.77
3	A	302	MES	C8-S	-7.41	1.67	1.77

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	301	MES	C5-N4-C3	5.10	120.31	108.83
3	A	302	MES	C5-N4-C3	3.85	117.48	108.83
3	A	302	MES	C6-C5-N4	-2.94	105.65	110.10
3	A	302	MES	O3S-S-C8	2.69	110.11	105.77
3	A	302	MES	C7-N4-C5	2.60	117.89	111.23
3	H	301	MES	O3S-S-C8	2.43	109.70	105.77
3	A	302	MES	O1S-S-C8	2.32	109.71	106.92
3	H	301	MES	C7-N4-C5	2.27	117.04	111.23
4	B	302	PGE	O2-C2-C1	2.15	119.51	110.07

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	H	301	MES	C8-C7-N4-C5
3	A	302	MES	C8-C7-N4-C5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	302	MES	C7-C8-S-O1S
3	A	302	MES	C7-C8-S-O2S
3	A	302	MES	C7-C8-S-O3S
4	L	301	PGE	O2-C3-C4-O3
6	B	301	PG4	O4-C7-C8-O5
6	B	301	PG4	O2-C3-C4-O3
4	L	301	PGE	O1-C1-C2-O2
6	A	301	PG4	O2-C3-C4-O3
4	L	301	PGE	C3-C4-O3-C5
4	B	302	PGE	O2-C3-C4-O3
3	H	301	MES	C7-C8-S-O3S
6	B	301	PG4	C4-C3-O2-C2
4	B	302	PGE	O1-C1-C2-O2
6	A	301	PG4	C8-C7-O4-C6
6	B	301	PG4	C6-C5-O3-C4
6	A	301	PG4	C4-C3-O2-C2
6	A	301	PG4	O4-C7-C8-O5
6	B	301	PG4	C5-C6-O4-C7
6	A	301	PG4	O1-C1-C2-O2
6	B	301	PG4	O1-C1-C2-O2
6	B	301	PG4	O3-C5-C6-O4
6	A	301	PG4	C5-C6-O4-C7
4	L	301	PGE	C6-C5-O3-C4
3	H	301	MES	C7-C8-S-O1S
3	H	301	MES	C7-C8-S-O2S
4	L	301	PGE	C4-C3-O2-C2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	301	PG4	3	0
6	A	301	PG4	1	0
4	L	301	PGE	1	0
4	B	302	PGE	3	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	212/225 (94%)	-0.14	8 (3%) 40 39	13, 33, 68, 88	0
1	H	212/225 (94%)	-0.25	8 (3%) 40 39	13, 32, 64, 86	0
2	B	210/214 (98%)	-0.31	6 (2%) 51 50	16, 30, 69, 89	0
2	L	211/214 (98%)	-0.38	4 (1%) 66 64	14, 30, 63, 82	0
All	All	845/878 (96%)	-0.27	26 (3%) 49 47	13, 31, 67, 89	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	THR	4.6
1	A	158	ALA	4.0
1	H	1	GLU	4.0
2	B	1	SER	3.6
2	L	1	SER	3.4
2	B	183	GLU	3.3
1	H	114	ALA	3.2
1	A	115	SER	3.2
1	A	172	SER	3.1
2	L	168	SER	3.0
2	L	189	ARG	2.8
1	H	187	SER	2.8
2	B	187	SER	2.8
1	A	114	ALA	2.7
1	H	191	THR	2.7
2	B	189	ARG	2.7
2	B	168	SER	2.7
1	A	1	GLU	2.4
1	H	186	SER	2.3
2	B	182	PRO	2.2
1	H	190	GLY	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	L	156	LYS	2.1
1	A	157	GLY	2.1
1	A	188	SER	2.1
1	H	158	ALA	2.0
1	H	201	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	CL	L	302	1/1	0.89	0.14	48,48,48,48	0
4	PGE	B	302	10/10	0.92	0.12	32,39,42,42	0
4	PGE	L	301	10/10	0.92	0.27	50,56,57,59	0
3	MES	A	302	12/12	0.93	0.15	55,59,69,72	0
3	MES	H	301	12/12	0.93	0.17	57,61,65,66	0
5	CL	B	303	1/1	0.94	0.08	38,38,38,38	0
6	PG4	A	301	13/13	0.94	0.22	31,36,40,42	0
6	PG4	B	301	13/13	0.94	0.22	29,37,48,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.