



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 02:31 AM EDT

PDB ID : 6QFJ
Title : MamB CTD magnetosome protein [Desulfamplus magnetovallimortis BW-1]
Authors : Zeytuni, N.Z.; Keren, N.K.; Zarivach, R.Z.
Deposited on : 2019-01-10
Resolution : 2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	84	<div> <div>2%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	B	84	<div> <div>%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>
1	C	84	<div> <div>13%</div> <div>76%</div> <div>19%</div> <div>..</div> </div>
1	D	84	<div> <div>11%</div> <div>73%</div> <div>23%</div> <div>..</div> </div>
1	E	84	<div> <div>30%</div> <div>76%</div> <div>17%</div> <div>..</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	84	<div><div></div><div>19%</div><div>74%</div><div>19%</div><div></div><div></div></div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	84	Total	C	N	O	S	0	3	0
			694	440	111	142	1			
1	B	83	Total	C	N	O	S	0	1	0
			674	430	110	133	1			
1	C	83	Total	C	N	O	S	0	0	0
			668	425	109	133	1			
1	D	81	Total	C	N	O		0	0	0
			650	414	105	131				
1	E	81	Total	C	N	O		0	0	0
			650	414	105	131				
1	F	81	Total	C	N	O		0	0	0
			650	414	105	131				

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	187	SER	-	expression tag	UNP L0R6P9
A	188	HIS	-	expression tag	UNP L0R6P9
A	189	MET	-	expression tag	UNP L0R6P9
B	187	SER	-	expression tag	UNP L0R6P9
B	188	HIS	-	expression tag	UNP L0R6P9
B	189	MET	-	expression tag	UNP L0R6P9
C	187	SER	-	expression tag	UNP L0R6P9
C	188	HIS	-	expression tag	UNP L0R6P9
C	189	MET	-	expression tag	UNP L0R6P9
D	187	SER	-	expression tag	UNP L0R6P9
D	188	HIS	-	expression tag	UNP L0R6P9
D	189	MET	-	expression tag	UNP L0R6P9
E	187	SER	-	expression tag	UNP L0R6P9
E	188	HIS	-	expression tag	UNP L0R6P9
E	189	MET	-	expression tag	UNP L0R6P9
F	187	SER	-	expression tag	UNP L0R6P9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	188	HIS	-	expression tag	UNP L0R6P9
F	189	MET	-	expression tag	UNP L0R6P9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	48	Total 48	O 48	0	0
2	B	47	Total 47	O 47	0	0
2	C	24	Total 24	O 24	0	0
2	D	24	Total 24	O 24	0	0
2	E	7	Total 7	O 7	0	0
2	F	6	Total 6	O 6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family




- Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family



- Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family




- Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family




- Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family





● Molecule 1: Magnetosome membrane protein MamB, putative Co/Zn/Cd cation transporter. Cation diffusion facilitator family



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.97Å 47.07Å 74.63Å 74.19° 84.86° 84.21°	Depositor
Resolution (Å)	23.91 – 2.13 23.90 – 2.13	Depositor EDS
% Data completeness (in resolution range)	97.0 (23.91-2.13) 97.2 (23.90-2.13)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 2.13Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.200 , 0.235 0.206 , 0.241	Depositor DCC
R_{free} test set	1683 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	38.2	Xtriage
Anisotropy	0.074	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 44.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4142	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/707	0.91	0/954
1	B	0.99	1/684 (0.1%)	0.94	0/922
1	C	0.89	1/675 (0.1%)	1.05	3/911 (0.3%)
1	D	0.87	1/656 (0.2%)	0.99	4/886 (0.5%)
1	E	0.79	2/656 (0.3%)	0.98	3/886 (0.3%)
1	F	0.73	0/656	0.96	2/886 (0.2%)
All	All	0.88	5/4034 (0.1%)	0.97	12/5445 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	239	GLU	CD-OE1	6.96	1.33	1.25
1	C	207	SER	CB-OG	-5.97	1.34	1.42
1	D	207	SER	CB-OG	-5.78	1.34	1.42
1	E	232	ASP	CB-CG	5.60	1.63	1.51
1	B	244	GLU	CG-CD	5.33	1.59	1.51

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	211	ASP	CB-CG-OD2	7.68	125.21	118.30
1	E	211	ASP	CB-CG-OD1	-7.50	111.55	118.30
1	D	240	ARG	NE-CZ-NH2	7.34	123.97	120.30
1	F	254	ASP	CB-CG-OD1	7.27	124.84	118.30
1	F	232	ASP	CB-CG-OD1	6.91	124.52	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	240	ARG	NE-CZ-NH2	6.36	123.48	120.30
1	E	232	ASP	CB-CG-OD1	5.78	123.50	118.30
1	D	191	ASP	CB-CG-OD1	5.59	123.33	118.30
1	C	191	ASP	CB-CG-OD2	5.38	123.14	118.30
1	D	211	ASP	CB-CG-OD1	5.31	123.08	118.30
1	D	240	ARG	NE-CZ-NH1	-5.27	117.67	120.30
1	C	270	LEU	CA-CB-CG	5.13	127.11	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	190	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	694	0	710	5	0
1	B	674	0	703	3	0
1	C	668	0	690	21	0
1	D	650	0	674	8	0
1	E	650	0	674	4	0
1	F	650	0	674	11	0
2	A	48	0	0	2	0
2	B	47	0	0	0	0
2	C	24	0	0	1	0
2	D	24	0	0	1	0
2	E	7	0	0	1	0
2	F	6	0	0	0	0
All	All	4142	0	4125	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:216:GLU:OE1	1:E:255:ARG:NH2	1.69	1.23
1:C:255:ARG:HH11	1:C:255:ARG:HB3	1.21	1.06
1:F:219:GLN:HA	1:F:255:ARG:HH21	1.41	0.85
1:A:187:SER:O	1:A:190[B]:GLU:HG3	1.79	0.83
1:C:219:GLN:O	1:C:255:ARG:NH2	2.11	0.83
1:C:255:ARG:HH11	1:C:255:ARG:CB	1.92	0.81
1:F:219:GLN:HA	1:F:255:ARG:NH2	2.00	0.75
1:C:189:MET:SD	1:C:221:LEU:HD21	2.25	0.75
1:F:219:GLN:CA	1:F:255:ARG:HH21	1.98	0.75
1:C:189:MET:HA	1:C:189:MET:HE3	1.71	0.72
1:C:255:ARG:HB3	1:C:255:ARG:NH1	2.02	0.70
1:A:201:GLU:OE1	2:A:301:HOH:O	2.09	0.70
1:C:188:HIS:N	1:C:191:ASP:OD1	2.28	0.67
1:C:189:MET:HE1	1:C:192:TYR:CD2	2.32	0.65
1:C:221:LEU:CD1	1:C:255:ARG:HD3	2.27	0.65
1:A:189:MET:HE3	2:A:320:HOH:O	1.97	0.64
1:C:189:MET:HE1	1:C:192:TYR:HD2	1.63	0.64
1:F:215:ARG:NH1	1:F:224:GLU:OE1	2.31	0.63
1:C:221:LEU:HD12	1:C:255:ARG:HD3	1.81	0.62
1:F:219:GLN:C	1:F:255:ARG:HH21	2.04	0.60
1:C:189:MET:CE	1:C:192:TYR:HD2	2.14	0.59
1:C:189:MET:SD	1:C:221:LEU:CD2	2.91	0.58
1:F:196:ILE:HD13	1:F:249:ILE:HD12	1.87	0.57
1:D:216:GLU:OE2	1:D:255:ARG:NH1	2.32	0.56
1:D:201:GLU:OE2	1:D:201:GLU:HA	2.06	0.55
1:C:188:HIS:O	1:C:191:ASP:HB2	2.06	0.55
1:A:187:SER:O	1:A:190[B]:GLU:CG	2.52	0.54
1:E:267:GLN:O	2:E:301:HOH:O	2.18	0.54
1:C:189:MET:HA	1:C:189:MET:CE	2.33	0.52
1:F:190:GLU:HG3	1:F:193:ILE:HB	1.91	0.51
1:B:246:ASN:O	1:B:250:LYS:HG2	2.13	0.49
1:F:255:ARG:HD3	1:F:256:LYS:O	2.14	0.48
1:A:246:ASN:O	1:A:250:LYS:HG2	2.13	0.47
1:F:246:ASN:O	1:F:250:LYS:HG2	2.15	0.46
1:C:246:ASN:O	1:C:250:LYS:HG2	2.17	0.45
1:E:215:ARG:HG3	1:E:222:GLU:HB2	1.98	0.45
1:E:215:ARG:NH2	1:E:224:GLU:OE2	2.50	0.45
1:F:219:GLN:C	1:F:255:ARG:HE	2.20	0.44
1:D:246:ASN:O	1:D:250:LYS:HG2	2.17	0.44
1:D:217:LEU:HB2	1:D:222:GLU:OE1	2.17	0.44
1:D:197:ALA:O	1:D:201:GLU:HG2	2.17	0.44
1:D:267:GLN:NE2	2:D:303:HOH:O	2.44	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:243[B]:LYS:HB3	1:B:243[B]:LYS:HE3	1.74	0.43
1:C:204:PRO:HD2	2:C:307:HOH:O	2.18	0.43
1:C:219:GLN:C	1:C:255:ARG:HH22	2.12	0.42
1:D:219:GLN:C	1:D:255:ARG:HH21	2.23	0.42
1:C:254:ASP:OD1	1:C:254:ASP:N	2.53	0.41
1:C:217:LEU:HD12	1:C:222:GLU:OE1	2.20	0.41
1:D:221:LEU:HG	1:D:255:ARG:CZ	2.49	0.41
1:C:188:HIS:CA	1:C:191:ASP:OD1	2.68	0.41
1:B:222:GLU:HG2	1:B:258:THR:OG1	2.19	0.41
1:F:196:ILE:CD1	1:F:249:ILE:HD12	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/84 (101%)	85 (100%)	0	0	100	100
1	B	82/84 (98%)	82 (100%)	0	0	100	100
1	C	81/84 (96%)	80 (99%)	1 (1%)	0	100	100
1	D	79/84 (94%)	79 (100%)	0	0	100	100
1	E	79/84 (94%)	79 (100%)	0	0	100	100
1	F	79/84 (94%)	79 (100%)	0	0	100	100
All	All	485/504 (96%)	484 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/77 (104%)	74 (92%)	6 (8%)	13	8
1	B	77/77 (100%)	71 (92%)	6 (8%)	12	7
1	C	76/77 (99%)	71 (93%)	5 (7%)	16	11
1	D	74/77 (96%)	69 (93%)	5 (7%)	16	10
1	E	74/77 (96%)	63 (85%)	11 (15%)	3	1
1	F	74/77 (96%)	65 (88%)	9 (12%)	5	2
All	All	455/462 (98%)	413 (91%)	42 (9%)	9	4

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	189	MET
1	A	201	GLU
1	A	254[A]	ASP
1	A	254[B]	ASP
1	A	255	ARG
1	A	258	THR
1	B	188	HIS
1	B	189	MET
1	B	201	GLU
1	B	253	VAL
1	B	254	ASP
1	B	255	ARG
1	C	225	ILE
1	C	253	VAL
1	C	254	ASP
1	C	255	ARG
1	C	258	THR
1	D	201	GLU
1	D	225	ILE
1	D	247	GLN
1	D	254	ASP
1	D	258	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	194	GLU
1	E	198	ASN
1	E	201	GLU
1	E	215	ARG
1	E	225	ILE
1	E	232	ASP
1	E	246	ASN
1	E	247	GLN
1	E	254	ASP
1	E	255	ARG
1	E	258	THR
1	F	201	GLU
1	F	225	ILE
1	F	232	ASP
1	F	243	LYS
1	F	246	ASN
1	F	247	GLN
1	F	253	VAL
1	F	255	ARG
1	F	258	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	246	ASN
1	C	246	ASN
1	C	247	GLN
1	D	198	ASN
1	D	267	GLN
1	E	247	GLN
1	F	198	ASN
1	F	247	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	84/84 (100%)	0.13	2 (2%) 59 65	23, 37, 64, 96	1 (1%)
1	B	83/84 (98%)	0.10	1 (1%) 79 83	24, 38, 68, 93	0
1	C	83/84 (98%)	0.60	11 (13%) 3 4	25, 60, 111, 127	0
1	D	81/84 (96%)	0.61	9 (11%) 5 6	25, 57, 94, 110	0
1	E	81/84 (96%)	1.56	25 (30%) 0 0	45, 84, 159, 184	0
1	F	81/84 (96%)	1.08	16 (19%) 1 1	40, 75, 138, 162	0
All	All	493/504 (97%)	0.67	64 (12%) 3 4	23, 56, 127, 184	1 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	217	LEU	10.6
1	E	220	VAL	8.8
1	E	217	LEU	8.2
1	E	252	ILE	8.1
1	F	218	GLY	7.6
1	D	219	GLN	7.3
1	E	258	THR	7.0
1	E	218	GLY	6.2
1	F	219	GLN	5.6
1	E	219	GLN	5.4
1	E	254	ASP	5.2
1	F	221	LEU	4.9
1	C	217	LEU	4.8
1	C	218	GLY	4.7
1	E	216	GLU	4.6
1	C	219	GLN	4.5
1	F	254	ASP	4.2
1	E	213	ILE	4.2
1	E	192	TYR	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	188	HIS	3.8
1	D	214	ALA	3.8
1	F	214	ALA	3.8
1	E	256	LYS	3.7
1	F	192	TYR	3.6
1	C	221	LEU	3.5
1	A	188	HIS	3.5
1	E	202	LYS	3.4
1	E	214	ALA	3.4
1	F	191	ASP	3.3
1	D	217	LEU	3.2
1	F	215	ARG	3.2
1	C	188	HIS	3.2
1	C	189	MET	3.2
1	D	256	LYS	3.1
1	E	251	GLU	3.1
1	E	230	PRO	3.0
1	E	222	GLU	3.0
1	F	249	ILE	2.9
1	D	204	PRO	2.9
1	D	220	VAL	2.8
1	E	195	ALA	2.7
1	F	204	PRO	2.7
1	E	221	LEU	2.6
1	E	215	ARG	2.6
1	F	258	THR	2.5
1	C	255	ARG	2.5
1	C	204	PRO	2.5
1	F	222	GLU	2.4
1	E	232	ASP	2.4
1	D	218	GLY	2.4
1	C	220	VAL	2.3
1	D	215	ARG	2.3
1	F	194	GLU	2.3
1	E	257	SER	2.3
1	C	216	GLU	2.3
1	F	232	ASP	2.3
1	F	195	ALA	2.2
1	E	255	ARG	2.2
1	A	190[A]	GLU	2.2
1	E	244	GLU	2.2
1	C	245	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	191	ASP	2.1
1	D	192	TYR	2.0
1	E	253	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.