



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 7, 2025 – 07:08 pm BST

PDB ID : 9QM1 / pdb_00009qm1
Title : W-formate dehydrogenase U192C from *Desulfovibrio vulgaris* - Soaked with Dithionite
Authors : Vilela-Alves, G.; Oliveira, A.R.; Pereira, I.C.; Romao, M.J.; Mota, C.
Deposited on : 2025-03-21
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1013	<div> <div>21%</div> <div>85%</div> <div>10%</div> <div>5%</div> </div>
2	B	214	<div> <div>5%</div> <div>97%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	SO2	A	1209	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 9828 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Formate dehydrogenase, alpha subunit, selenocysteine-containing.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	967	Total	C	N	O	S	0	1	0
			7579	4832	1321	1384	42			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	192	CYS	SEC	engineered mutation	UNP Q72EJ1
A	1006	TRP	-	expression tag	UNP Q72EJ1
A	1007	SER	-	expression tag	UNP Q72EJ1
A	1008	HIS	-	expression tag	UNP Q72EJ1
A	1009	PRO	-	expression tag	UNP Q72EJ1
A	1010	GLN	-	expression tag	UNP Q72EJ1
A	1011	PHE	-	expression tag	UNP Q72EJ1
A	1012	GLU	-	expression tag	UNP Q72EJ1
A	1013	LYS	-	expression tag	UNP Q72EJ1

- Molecule 2 is a protein called Formate dehydrogenase, beta subunit, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	214	Total	C	N	O	S	0	0	0
			1664	1041	291	316	16			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is SULFUR DIOXIDE (CCD ID: SO2) (formula: O_2S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			3	2	1		

- Molecule 7 is HYDROSULFURIC ACID (CCD ID: H2S) (formula: H₂S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	S	0	0
			1	1		

- Molecule 8 is TUNGSTEN ION (CCD ID: W) (formula: W) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total W 1 1	0	0

- Molecule 9 is 2-AMINO-5,6-DIMERCAPTO-7-METHYL-3,7,8A,9-TETRAHYDRO-8-OXA-1,3,9,10-TETRAAZA-ANTHRACEN-4-ONE GUANOSINE DINUCLEOTIDE (CCD ID: MGD) (formula: $C_{20}H_{26}N_{10}O_{13}P_2S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
9	A	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0
9	A	1	Total 47	C 20	N 10	O 13	P 2	S 2	0	0


- Molecule 10 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	Fe	S	0	0
			8	4	4		
10	B	1	Total	Fe	S	0	0
			8	4	4		
10	B	1	Total	Fe	S	0	0
			8	4	4		
10	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	296	Total	O	0	0
			296	296		
11	B	119	Total	O	0	0
			119	119		



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	61.80Å 122.51Å 146.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.93 – 2.00 62.93 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.5 (62.93-2.00) 99.3 (62.93-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.214 , 0.267 0.221 , 0.272	Depositor DCC
R_{free} test set	3696 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9828	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: W, PEG, H2S, SF4, MGD, SO2, GOL, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.99	0/7790	1.37	0/10566
2	B	0.99	0/1699	1.38	0/2302
All	All	0.99	0/9489	1.37	0/12868

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7579	0	7425	71	0
2	B	1664	0	1633	4	0
3	A	12	0	16	1	0
4	A	20	0	30	0	0
5	A	7	0	10	0	0
6	A	3	0	0	3	0
7	A	1	0	0	0	0
8	A	1	0	0	0	0
9	A	94	0	44	5	0
10	A	8	0	0	0	0
10	B	24	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
11	A	296	0	0	1	0
11	B	119	0	0	0	0
All	All	9828	0	9158	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:GLU:HA	1:A:690:MET:HE1	1.51	0.91
1:A:171:TRP:CB	1:A:690:MET:HE2	2.11	0.79
1:A:171:TRP:HB3	1:A:690:MET:HE2	1.73	0.69
1:A:171:TRP:HB3	1:A:690:MET:CE	2.25	0.66
1:A:775:ARG:O	3:A:1201:GOL:O2	2.14	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	964/1013 (95%)	919 (95%)	39 (4%)	6 (1%)	22	17
2	B	212/214 (99%)	204 (96%)	8 (4%)	0	100	100
All	All	1176/1227 (96%)	1123 (96%)	47 (4%)	6 (0%)	25	21

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	GLN
1	A	592	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	678	HIS
1	A	636	ALA
1	A	533	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	788/820 (96%)	769 (98%)	19 (2%)	44	47
2	B	185/185 (100%)	185 (100%)	0	100	100
All	All	973/1005 (97%)	954 (98%)	19 (2%)	50	55

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	696	LYS
1	A	795	LYS
1	A	969	MET
1	A	754	ARG
1	A	633	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	ASN
1	A	704	GLN
2	B	24	GLN
2	B	194	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 1 is modelled with single atom and 1 is monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
10	SF4	B	302	2	0,12,12	-	-	-		
4	EDO	A	1204	-	3,3,3	0.09	0	2,2,2	0.20	0
10	SF4	B	301	2	0,12,12	-	-	-		
5	PEG	A	1206	-	6,6,6	0.17	0	5,5,5	0.11	0
6	SO2	A	1209	-	2,2,2	0.34	0	1,1,1	0.06	0
9	MGD	A	1212	8	41,52,52	0.77	1 (2%)	40,81,81	1.11	3 (7%)
4	EDO	A	1207	-	3,3,3	0.07	0	2,2,2	0.20	0
10	SF4	A	1214	1	0,12,12	-	-	-		
3	GOL	A	1201	-	5,5,5	0.09	0	5,5,5	0.22	0
10	SF4	B	303	2	0,12,12	-	-	-		
9	MGD	A	1213	8	41,52,52	1.00	2 (4%)	40,81,81	1.16	4 (10%)
4	EDO	A	1208	-	3,3,3	0.07	0	2,2,2	0.16	0
3	GOL	A	1203	-	5,5,5	0.11	0	5,5,5	0.28	0
4	EDO	A	1205	-	3,3,3	0.07	0	2,2,2	0.20	0
4	EDO	A	1202	-	3,3,3	0.05	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	SF4	B	302	2	-	-	0/6/5/5
4	EDO	A	1204	-	-	1/1/1/1	-
10	SF4	B	301	2	-	-	0/6/5/5
5	PEG	A	1206	-	-	2/4/4/4	-
9	MGD	A	1212	8	-	1/18/66/66	0/6/6/6
4	EDO	A	1207	-	-	1/1/1/1	-
10	SF4	A	1214	1	-	-	0/6/5/5
3	GOL	A	1201	-	-	2/4/4/4	-
10	SF4	B	303	2	-	-	0/6/5/5
9	MGD	A	1213	8	-	5/18/66/66	0/6/6/6
4	EDO	A	1208	-	-	0/1/1/1	-
3	GOL	A	1203	-	-	2/4/4/4	-
4	EDO	A	1205	-	-	1/1/1/1	-
4	EDO	A	1202	-	-	1/1/1/1	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	1213	MGD	C23-C14	4.12	1.56	1.53
9	A	1212	MGD	C5-C6	-2.62	1.42	1.47
9	A	1213	MGD	C5-C6	-2.58	1.42	1.47

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	1213	MGD	O11-C23-C14	3.59	111.36	108.96
9	A	1212	MGD	C19-N20-C21	2.92	118.69	113.43
9	A	1213	MGD	C19-N20-C21	2.81	118.50	113.43
9	A	1212	MGD	O4'-C1'-C2'	-2.33	103.53	106.93
9	A	1213	MGD	O4'-C1'-C2'	-2.28	103.60	106.93

There are no chirality outliers.

5 of 16 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1201	GOL	O1-C1-C2-O2
3	A	1201	GOL	O1-C1-C2-C3
4	A	1204	EDO	O1-C1-C2-O2
9	A	1213	MGD	C4'-C5'-O5'-PB
9	A	1213	MGD	O4'-C4'-C5'-O5'

There are no ring outliers.


4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1209	SO2	3	0
9	A	1212	MGD	4	0
3	A	1201	GOL	1	0
9	A	1213	MGD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand SF4 B 301




Bond lengths





Bond angles




Torsions




Rings







Ligand SF4 B 303




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	967/1013 (95%)	1.11	212 (21%) 3 2	10, 31, 62, 100	1 (0%)
2	B	214/214 (100%)	0.40	10 (4%) 37 35	14, 25, 52, 83	0
All	All	1181/1227 (96%)	0.98	222 (18%) 4 3	10, 30, 60, 100	1 (0%)

The worst 5 of 222 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	107	PRO	8.5
1	A	117	TRP	7.6
1	A	122	TRP	7.5
1	A	672	ILE	6.1
1	A	109	TYR	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	EDO	A	1202	4/4	0.67	0.19	41,49,57,62	0
4	EDO	A	1208	4/4	0.70	0.19	33,41,62,93	0
4	EDO	A	1207	4/4	0.72	0.15	30,37,45,48	0
5	PEG	A	1206	7/7	0.75	0.20	29,49,66,78	0
3	GOL	A	1203	6/6	0.76	0.19	31,58,59,64	0
3	GOL	A	1201	6/6	0.82	0.20	31,49,61,63	0
6	SO2	A	1209	3/3	0.83	0.18	21,21,30,36	3
4	EDO	A	1205	4/4	0.88	0.13	32,38,54,61	0
4	EDO	A	1204	4/4	0.92	0.09	23,23,23,39	0
9	MGD	A	1212	47/47	0.96	0.06	10,16,22,29	0
9	MGD	A	1213	47/47	0.96	0.07	13,22,36,47	0
7	H2S	A	1210	1/1	0.98	0.12	27,27,27,27	0
10	SF4	A	1214	8/8	0.99	0.03	15,17,19,20	0
10	SF4	B	301	8/8	0.99	0.03	13,15,17,17	0
10	SF4	B	302	8/8	0.99	0.04	18,19,21,23	0
10	SF4	B	303	8/8	0.99	0.03	21,21,22,24	0
8	W	A	1211	1/1	1.00	0.02	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SO2 A 1209:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGD A 1212:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around MGD A 1213:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H2S A 1210:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 A 1214:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 B 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around SF4 B 303:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around W A 1211:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.