



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2025 – 06:52 PM EDT

PDB ID : 4QX8  
Title : Crystal structure of histone demethylase kdm2a-h3k36me3 complex with alpha-kg  
Authors : Cheng, Z.J.; Patel, D.J.  
Deposited on : 2014-07-19  
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	<div> <div>3%</div> <div>88%</div> <div>10%</div> <div>.</div> </div>
1	C	329	<div> <div>5%</div> <div>91%</div> <div>9%</div> </div>
2	B	68	<div> <div>%</div> <div>88%</div> <div>12%</div> </div>
2	D	68	<div> <div>4%</div> <div>94%</div> <div>.</div> <div>.</div> </div>
3	E	15	<div> <div>7%</div> <div>53%</div> <div>20%</div> <div>27%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	15	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	M3L	F	36	-	-	X	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	329	Total	C	N	O	S	0	0	0
			2743	1758	458	506	21			
1	C	329	Total	C	N	O	S	0	3	0
			2762	1769	463	509	21			

- Molecule 2 is a protein called Lysine-specific demethylase 2A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	68	Total	C	N	O	S	0	2	0
			538	348	87	101	2			
2	D	68	Total	C	N	O	S	0	3	0
			542	350	86	103	3			

- Molecule 3 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	11	Total	C	N	O	0	0	0
			77	50	15	12			
3	F	10	Total	C	N	O	0	0	0
			76	48	17	11			

- Molecule 4 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ni	0	0
			1	1		
4	C	1	Total	Ni	0	0
			1	1		

- Molecule 5 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula: C<sub>5</sub>H<sub>6</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	5	5		
5	C	1	Total	C	O	0	0
			10	5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	300	Total	O	0	0
			300	300		
6	B	76	Total	O	0	0
			76	76		
6	C	286	Total	O	0	0
			286	286		
6	D	61	Total	O	0	0
			61	61		
6	E	5	Total	O	0	0
			5	5		
6	F	6	Total	O	0	0
			6	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lysine-specific demethylase 2A




- Molecule 1: Lysine-specific demethylase 2A




- Molecule 2: Lysine-specific demethylase 2A




- Molecule 2: Lysine-specific demethylase 2A



- Molecule 3: Histone H3.2



● Molecule 3: Histone H3.2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.80Å 84.61Å 170.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.04 – 1.65 85.04 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.2 (85.04-1.65) 99.2 (85.04-1.65)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 1.65Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.188 , 0.227 0.197 , 0.234	Depositor DCC
$R_{free}$ test set	4648 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtriage
Anisotropy	0.096	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 34.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7494	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.84 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5482e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: M3L, NI, AKG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/2819	0.62	0/3817
1	C	0.40	0/2844	0.64	0/3850
2	B	0.36	0/555	0.60	0/756
2	D	0.35	0/561	0.68	1/763 (0.1%)
3	E	0.34	0/67	0.49	0/91
3	F	0.36	0/65	0.53	0/86
All	All	0.39	0/6911	0.63	1/9363 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	462	ARG	NE-CZ-NH2	6.00	123.30	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	0	2639	24	0
1	C	2762	0	2662	22	0
2	B	538	0	558	5	0
2	D	542	0	566	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	77	0	83	4	0
3	F	76	0	84	10	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	10	0	4	3	0
5	C	10	0	4	1	0
6	A	300	0	0	4	0
6	B	76	0	0	1	0
6	C	286	0	0	3	0
6	D	61	0	0	0	0
6	E	5	0	0	0	0
6	F	6	0	0	0	0
All	All	7494	0	6600	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:222:TYR:OH	3:F:36:M3L:HM11	1.86	0.76
1:A:69:ARG:NH2	6:A:907:HOH:O	2.27	0.66
2:D:462:ARG:HG2	2:D:462:ARG:HH21	1.64	0.62
1:C:222:TYR:OH	3:F:36:M3L:CM1	2.47	0.61
1:C:222:TYR:CZ	3:F:36:M3L:HM11	2.36	0.61
1:A:290:THR:O	1:A:292:THR:HG23	2.03	0.57
2:B:453:LEU:CD2	2:B:513:ILE:HG13	2.34	0.57
1:A:226:GLN:O	1:A:292:THR:HB	2.05	0.56
1:C:298:ASN:HD21	3:F:36:M3L:HE2	1.71	0.55
1:C:298:ASN:HD21	3:F:36:M3L:HM12	1.71	0.55
1:C:150:HIS:HE1	1:C:193:TYR:OH	1.90	0.54
1:A:290:THR:O	1:A:292:THR:CG2	2.56	0.54
2:B:450:GLN:NE2	6:B:674:HOH:O	2.40	0.54
1:C:122:THR:H	1:C:125:GLN:HE21	1.54	0.53
1:C:253:GLN:HE22	3:F:40:ARG:HB2	1.73	0.53
1:C:128:ARG:O	1:C:132:THR:HG23	2.10	0.52
1:A:144:ILE:HD12	5:A:602:AKG:O2	2.10	0.52
1:A:150:HIS:HE1	1:A:193:TYR:OH	1.93	0.52
1:A:150:HIS:CE1	1:A:181:GLN:HG2	2.46	0.51
1:A:122:THR:H	1:A:125:GLN:HE21	1.57	0.51
5:A:602:AKG:C1	3:E:36:M3L:HM22	2.41	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:150:HIS:CE1	1:C:181:GLN:HG2	2.48	0.48
1:A:150:HIS:HD2	6:A:779:HOH:O	1.97	0.48
1:C:37:THR:HB	1:C:255:ASP:HA	1.96	0.47
2:D:462:ARG:HH21	2:D:462:ARG:CG	2.28	0.47
5:C:602:AKG:O2	3:F:36:M3L:HM22	2.14	0.47
1:A:228:GLY:H	1:A:292:THR:HG21	1.80	0.47
1:C:301:HIS:CD2	1:C:303:PHE:H	2.32	0.47
6:C:802:HOH:O	3:F:36:M3L:HE3	2.14	0.46
1:A:116:GLN:NE2	1:A:210:ASP:H	2.14	0.46
1:C:132:THR:HG21	6:C:886:HOH:O	2.14	0.46
1:C:298:ASN:HD21	3:F:36:M3L:CE	2.28	0.45
5:A:602:AKG:O2	3:E:36:M3L:HM22	2.16	0.45
1:A:172:MET:HA	1:A:339:ARG:HG3	1.99	0.45
1:A:47:ASN:C	1:A:47:ASN:HD22	2.19	0.44
1:C:191:MET:HG2	1:C:193:TYR:O	2.17	0.44
2:B:465:VAL:CG2	2:B:492:VAL:HG11	2.47	0.44
1:A:234:ILE:HB	1:A:283:ILE:HB	2.00	0.44
1:A:298:ASN:HD21	3:E:36:M3L:HE2	1.83	0.43
1:A:301:HIS:HE1	6:A:734:HOH:O	2.00	0.43
1:A:201:LEU:HD23	1:A:201:LEU:N	2.34	0.43
1:A:129:TYR:CE1	1:A:137:ARG:HD2	2.54	0.43
1:C:128:ARG:O	1:C:132:THR:CG2	2.66	0.43
1:C:298:ASN:ND2	3:F:36:M3L:HE2	2.33	0.43
1:C:113:VAL:HG11	1:C:140:LEU:HD23	2.01	0.43
1:A:106:ARG:HD2	6:A:728:HOH:O	2.19	0.43
1:C:234:ILE:HB	1:C:283:ILE:HB	2.01	0.42
1:C:304:ASN:OD1	6:C:805:HOH:O	2.22	0.42
1:A:129:TYR:CZ	1:A:137:ARG:HD2	2.55	0.42
1:A:344:ILE:HG21	2:B:495:LEU:HD11	2.02	0.41
1:C:201:LEU:HB2	1:C:294:VAL:HB	2.02	0.41
3:E:29:ALA:N	3:E:30:PRO:HD3	2.36	0.41
1:A:288:THR:HG22	1:A:290:THR:O	2.21	0.41
1:A:191:MET:HG2	1:A:193:TYR:O	2.20	0.41
2:B:453:LEU:HD22	2:B:513:ILE:HG13	2.02	0.41
1:C:150:HIS:CE1	1:C:193:TYR:OH	2.72	0.40
1:A:301:HIS:CD2	1:A:303:PHE:H	2.38	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/329 (99%)	323 (99%)	4 (1%)	0	100	100
1	C	330/329 (100%)	326 (99%)	4 (1%)	0	100	100
2	B	68/68 (100%)	67 (98%)	0	1 (2%)	8	1
2	D	69/68 (102%)	66 (96%)	3 (4%)	0	100	100
3	E	8/15 (53%)	8 (100%)	0	0	100	100
3	F	7/15 (47%)	6 (86%)	1 (14%)	0	100	100
All	All	809/824 (98%)	796 (98%)	12 (2%)	1 (0%)	48	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	483	GLU

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/305 (100%)	291 (95%)	14 (5%)	23	5
1	C	308/305 (101%)	299 (97%)	9 (3%)	37	14
2	B	63/61 (103%)	62 (98%)	1 (2%)	58	37
2	D	64/61 (105%)	59 (92%)	5 (8%)	10	1
3	E	6/10 (60%)	6 (100%)	0	100	100
3	F	6/10 (60%)	6 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	752/752 (100%)	723 (96%)	29 (4%)	28 7

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	THR
1	A	47	ASN
1	A	50	ASN
1	A	60	LYS
1	A	69	ARG
1	A	117	LYS
1	A	142	ASN
1	A	153	LEU
1	A	201	LEU
1	A	214	ASP
1	A	215	PHE
1	A	243	LEU
1	A	292	THR
1	A	293	LEU
2	B	511	VAL
1	C	40	LEU
1	C	45	GLN
1	C	50	ASN
1	C	132	THR
1	C	142	ASN
1	C	215	PHE
1	C	243	LEU
1	C	258	LEU
1	C	319	ARG
2	D	462	ARG
2	D	468	LEU
2	D	480[A]	THR
2	D	480[B]	THR
2	D	511	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	50	ASN
1	A	114	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	116	GLN
1	A	125	GLN
1	A	142	ASN
1	A	150	HIS
1	A	298	ASN
1	A	301	HIS
1	A	346	ASN
2	B	450	GLN
1	C	50	ASN
1	C	116	GLN
1	C	125	GLN
1	C	142	ASN
1	C	150	HIS
1	C	253	GLN
1	C	298	ASN
1	C	301	HIS
2	D	455	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	M3L	E	36	3	10,11,12	0.75	0	9,14,16	0.80	0
3	M3L	F	36	3	10,11,12	0.67	0	9,14,16	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	M3L	E	36	3	-	1/9/10/12	-
3	M3L	F	36	3	-	4/9/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	36	M3L	CE-CD-CG-CB
3	F	36	M3L	CD-CE-NZ-CM2
3	F	36	M3L	CD-CE-NZ-CM1
3	F	36	M3L	CD-CE-NZ-CM3
3	E	36	M3L	CD-CE-NZ-CM3

There are no ring outliers.

2 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	36	M3L	3	0
3	F	36	M3L	9	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	AKG	C	602	4	9,9,9	1.96	2 (22%)	11,11,11	1.78	3 (27%)
5	AKG	A	602	4	9,9,9	1.92	2 (22%)	11,11,11	1.44	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AKG	C	602	4	-	2/9/9/9	-
5	AKG	A	602	4	-	3/9/9/9	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	602	AKG	C2-C1	-4.59	1.46	1.53
5	A	602	AKG	C2-C1	-4.48	1.46	1.53
5	A	602	AKG	O2-C1	-2.18	1.24	1.30
5	C	602	AKG	O2-C1	-2.15	1.24	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	602	AKG	C3-C2-C1	3.25	121.38	115.86
5	A	602	AKG	C3-C2-C1	2.51	120.12	115.86
5	C	602	AKG	C4-C3-C2	-2.28	108.64	112.91
5	C	602	AKG	O4-C5-C4	2.22	121.03	114.00
5	A	602	AKG	O4-C5-C4	2.05	120.47	114.00

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	602	AKG	C1-C2-C3-C4
5	C	602	AKG	C1-C2-C3-C4
5	A	602	AKG	C3-C4-C5-O4
5	A	602	AKG	C3-C4-C5-O3
5	C	602	AKG	C3-C4-C5-O4

There are no ring outliers.



2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	602	AKG	1	0
5	A	602	AKG	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	329/329 (100%)	0.35	10 (3%) 52 56	17, 26, 49, 59	0
1	C	329/329 (100%)	0.27	15 (4%) 38 41	13, 24, 44, 62	3 (0%)
2	B	68/68 (100%)	0.32	1 (1%) 71 76	12, 27, 36, 50	2 (2%)
2	D	68/68 (100%)	0.53	3 (4%) 39 43	13, 28, 46, 51	3 (4%)
3	E	10/15 (66%)	0.77	1 (10%) 14 15	25, 30, 41, 46	0
3	F	9/15 (60%)	1.76	4 (44%) 1 0	37, 44, 60, 72	0
All	All	813/824 (98%)	0.35	34 (4%) 41 44	12, 26, 46, 72	8 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	31	ALA	3.3
1	C	251	GLY	3.1
3	E	29	ALA	3.0
1	A	363	MET	3.0
1	C	37	THR	3.0
1	A	254	GLY	3.0
1	C	189	LEU	2.9
1	C	255	ASP	2.9
3	F	35	VAL	2.8
1	C	322	ASN	2.7
2	D	473	LEU	2.7
1	C	254	GLY	2.7
1	C	38	PHE	2.6
1	C	250	SER	2.6
1	C	132	THR	2.6
1	C	321	PRO	2.6
2	D	500	ALA	2.5
1	A	37	THR	2.5
2	D	499	LEU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	252	LYS	2.4
1	A	113	VAL	2.4
1	A	69	ARG	2.4
1	A	40	LEU	2.3
1	C	188	ILE	2.3
1	A	322	ASN	2.2
1	C	190	GLU	2.2
1	A	252	LYS	2.2
1	A	189	LEU	2.2
3	F	40	ARG	2.1
2	B	450	GLN	2.1
3	F	39	HIS	2.1
1	C	138	GLU	2.1
1	C	222	TYR	2.1
1	A	38	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	M3L	F	36	12/13	0.80	0.13	38,39,41,43	0
3	M3L	E	36	12/13	0.93	0.07	25,28,31,31	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	AKG	C	602	10/10	0.91	0.11	25,27,28,32	2

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	AKG	A	602	10/10	0.92	0.10	29,31,33,33	2
4	NI	C	601	1/1	0.98	0.07	13,13,13,13	1
4	NI	A	601	1/1	1.00	0.05	16,16,16,16	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.