



Full wwPDB X-ray Structure Validation Report ⓘ

May 7, 2025 – 07:17 pm BST

PDB ID : 9QZ0 / pdb_00009qz0
Title : MINPP1 from Bacteroides thetaiotaomicron E325N mutant complex with myo-inositol hexakisphosphate
Authors : Li, A.W.H.; Shang, X.Y.; Hemmings, A.M.
Deposited on : 2025-04-22
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0rc1
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AA	426	<div> <div></div> <div>84%</div> <div>8%</div> <div>8%</div> </div>
1	BA	426	<div> <div>2%</div> <div>81%</div> <div>11%</div> <div>8%</div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7282 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called multiple inositol polyphosphate histidine phosphatase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AA	393	Total	C	N	O	S	0	6	0
			3249	2102	554	576	17			
1	BA	394	Total	C	N	O	S	0	10	0
			3268	2115	558	577	18			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AA	0	MET	-	initiating methionine	UNP Q89YI8
AA	1	GLY	-	expression tag	UNP Q89YI8
AA	2	SER	-	expression tag	UNP Q89YI8
AA	3	SER	-	expression tag	UNP Q89YI8
AA	4	HIS	-	expression tag	UNP Q89YI8
AA	5	HIS	-	expression tag	UNP Q89YI8
AA	6	HIS	-	expression tag	UNP Q89YI8
AA	7	HIS	-	expression tag	UNP Q89YI8
AA	8	HIS	-	expression tag	UNP Q89YI8
AA	9	HIS	-	expression tag	UNP Q89YI8
AA	10	SER	-	expression tag	UNP Q89YI8
AA	11	SER	-	expression tag	UNP Q89YI8
AA	12	GLY	-	expression tag	UNP Q89YI8
AA	13	LEU	-	expression tag	UNP Q89YI8
AA	14	VAL	-	expression tag	UNP Q89YI8
AA	15	PRO	-	expression tag	UNP Q89YI8
AA	16	ARG	-	expression tag	UNP Q89YI8
AA	17	GLY	-	expression tag	UNP Q89YI8
AA	18	SER	-	expression tag	UNP Q89YI8
AA	19	HIS	-	expression tag	UNP Q89YI8
AA	20	MET	-	expression tag	UNP Q89YI8
AA	325	ASN	GLU	engineered mutation	UNP Q89YI8
BA	0	MET	-	initiating methionine	UNP Q89YI8
BA	1	GLY	-	expression tag	UNP Q89YI8
BA	2	SER	-	expression tag	UNP Q89YI8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
BA	3	SER	-	expression tag	UNP Q89YI8
BA	4	HIS	-	expression tag	UNP Q89YI8
BA	5	HIS	-	expression tag	UNP Q89YI8
BA	6	HIS	-	expression tag	UNP Q89YI8
BA	7	HIS	-	expression tag	UNP Q89YI8
BA	8	HIS	-	expression tag	UNP Q89YI8
BA	9	HIS	-	expression tag	UNP Q89YI8
BA	10	SER	-	expression tag	UNP Q89YI8
BA	11	SER	-	expression tag	UNP Q89YI8
BA	12	GLY	-	expression tag	UNP Q89YI8
BA	13	LEU	-	expression tag	UNP Q89YI8
BA	14	VAL	-	expression tag	UNP Q89YI8
BA	15	PRO	-	expression tag	UNP Q89YI8
BA	16	ARG	-	expression tag	UNP Q89YI8
BA	17	GLY	-	expression tag	UNP Q89YI8
BA	18	SER	-	expression tag	UNP Q89YI8
BA	19	HIS	-	expression tag	UNP Q89YI8
BA	20	MET	-	expression tag	UNP Q89YI8
BA	325	ASN	GLU	engineered mutation	UNP Q89YI8

- Molecule 2 is INOSITOL HEXAKISPHOSPHATE (CCD ID: IHP) (formula: $C_6H_{18}O_{24}P_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AA	1	Total	C	O	P	0	0
			36	6	24	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AA	1	Total	C	O	P	0	0
			36	6	24	6		
2	BA	1	Total	C	O	P	0	0
			36	6	24	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AA	362	Total	O	0	0
			362	362		
3	BA	295	Total	O	0	0
			295	295		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: multiple inositol polyphosphate histidine phosphatase 1



- Molecule 1: multiple inositol polyphosphate histidine phosphatase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	52.83Å 120.41Å 75.94Å 90.00° 108.04° 90.00°	Depositor
Resolution (Å)	26.40 – 1.86 26.40 – 1.86	Depositor EDS
% Data completeness (in resolution range)	97.3 (26.40-1.86) 93.5 (26.40-1.86)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.87Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.165 , 0.193 0.165 , 0.193	Depositor DCC
R_{free} test set	5471 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.568	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7282	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: IHP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AA	0.33	0/3353	0.52	0/4537
1	BA	0.32	0/3384	0.53	0/4579
All	All	0.33	0/6737	0.53	0/9116

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AA	3249	0	3260	27	0
1	BA	3268	0	3297	34	0
2	AA	72	0	12	4	0
2	BA	36	0	6	2	0
3	AA	362	0	0	12	1
3	BA	295	0	0	8	1
All	All	7282	0	6575	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:AA:501:IHP:O24	3:AA:601:HOH:O	1.90	0.88
2:BA:501:IHP:O31	3:BA:601:HOH:O	1.93	0.85
1:BA:152:ASN:ND2	3:BA:603:HOH:O	2.12	0.76
1:AA:403:GLU:OE1	1:AA:406:ARG:NH2	2.19	0.76
1:AA:110:LYS:HG2	3:AA:929:HOH:O	1.88	0.74
1:BA:223:LEU:O	3:BA:602:HOH:O	2.05	0.73
1:BA:255:ASN:OD1	1:BA:257:GLU:HG2	1.88	0.72
1:AA:88:SER:HA	1:AA:91:MET:HE2	1.72	0.70
1:BA:177[B]:GLN:H	1:BA:177[B]:GLN:CD	2.02	0.66
1:AA:342:GLN:OE1	3:AA:603:HOH:O	2.14	0.65
2:AA:501:IHP:O31	3:AA:602:HOH:O	2.14	0.64
1:AA:89:GLU:OE1	1:AA:220:LYS:HE3	2.01	0.61
1:AA:36:ASN:HD22	1:AA:37:ARG:N	1.97	0.61
1:BA:411:GLN:O	1:BA:415:MET:HG3	2.01	0.61
1:AA:312:LYS:HE2	3:AA:799:HOH:O	2.04	0.58
1:AA:224:ASN:ND2	1:AA:226:GLU:OE1	2.37	0.58
1:BA:372:ASP:OD2	1:BA:376:ARG:NH1	2.37	0.58
1:BA:219:LYS:HG2	1:BA:225:PRO:HG3	1.87	0.57
1:AA:22:THR:OG1	1:AA:25:GLN:HG3	2.06	0.56
1:BA:218:MET:HE3	1:BA:236:PHE:CD2	2.41	0.56
1:AA:226:GLU:H	1:AA:226:GLU:CD	2.14	0.55
1:AA:98:ARG:NE	3:AA:609:HOH:O	2.35	0.55
1:BA:177[B]:GLN:NE2	3:BA:613:HOH:O	2.41	0.52
1:AA:274:LEU:O	1:AA:278:MET:HG3	2.09	0.52
1:BA:274:LEU:HG	1:BA:278:MET:HE2	1.92	0.52
1:BA:23:LYS:H	1:BA:23:LYS:HE2	1.74	0.51
2:AA:501:IHP:O45	3:AA:605:HOH:O	2.20	0.51
1:AA:209[B]:HIS:HE1	3:AA:888:HOH:O	1.94	0.51
1:BA:344:CYS:HB2	3:BA:851:HOH:O	2.11	0.50
1:BA:124:GLY:O	1:BA:128:ARG:HG3	2.12	0.49
1:BA:337:GLU:O	1:BA:338:LYS:HB3	2.11	0.49
1:BA:143:ILE:HA	1:BA:173:SER:O	2.13	0.49
1:BA:209[B]:HIS:ND1	1:BA:234:GLU:OE1	2.44	0.49
1:BA:96:MET:SD	1:BA:254:LEU:HD13	2.53	0.48
1:BA:120:GLU:HG2	1:BA:160[B]:CYS:SG	2.54	0.48
1:AA:137:SER:N	3:AA:606:HOH:O	2.20	0.47
1:AA:96:MET:HA	1:AA:96:MET:HE2	1.96	0.47
1:BA:23:LYS:H	1:BA:23:LYS:CE	2.29	0.46
1:AA:163:ARG:NH2	3:AA:618:HOH:O	2.49	0.45
1:AA:48:MET:HE2	1:AA:48:MET:HB2	1.79	0.45
2:AA:502:IHP:O21	1:BA:354:LYS:HE2	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BA:53:ILE:HG13	1:BA:133:LEU:HD21	1.99	0.44
1:BA:397:PHE:CD1	1:BA:398:PRO:HA	2.53	0.44
1:BA:397:PHE:CG	1:BA:398:PRO:HA	2.52	0.44
1:AA:36:ASN:HD22	1:AA:36:ASN:C	2.25	0.44
1:BA:48:MET:N	3:BA:621:HOH:O	2.51	0.44
1:BA:295:TRP:CG	1:BA:296:PRO:HD3	2.53	0.44
1:AA:22:THR:OG1	1:AA:24:ILE:HG22	2.18	0.43
1:BA:98:ARG:NH1	3:BA:619:HOH:O	2.48	0.43
1:BA:218:MET:HE2	1:BA:222:LEU:HD12	2.01	0.43
1:AA:137:SER:HA	1:AA:167:ALA:O	2.17	0.43
1:BA:53:ILE:HG13	1:BA:133:LEU:CD2	2.49	0.43
1:BA:334:MET:HE1	1:BA:381[A]:ILE:HG21	2.02	0.42
1:BA:68:LYS:HA	1:BA:68:LYS:HD3	1.71	0.42
1:AA:337:GLU:O	1:AA:338:LYS:HB2	2.20	0.42
1:BA:231:LYS:H	1:BA:231:LYS:HG2	1.46	0.41
1:AA:106:GLY:O	1:AA:110:LYS:HE3	2.20	0.41
1:BA:334:MET:HE2	1:BA:334:MET:HB3	1.87	0.41
1:AA:36:ASN:ND2	1:AA:38:THR:H	2.17	0.41
1:BA:23:LYS:HD3	1:BA:23:LYS:N	2.35	0.41
1:AA:219:LYS:HA	1:AA:225:PRO:HG3	2.01	0.41
1:AA:180:HIS:HB2	3:AA:821:HOH:O	2.20	0.41
1:AA:397:PHE:CG	1:AA:398:PRO:HA	2.56	0.41
2:BA:501:IHP:O24	3:BA:604:HOH:O	2.22	0.40
1:AA:117:THR:HG22	3:AA:868:HOH:O	2.21	0.40
1:BA:159:SER:HB3	1:BA:163:ARG:NH1	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:AA:916:HOH:O	3:BA:767:HOH:O[2_647]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AA	395/426 (93%)	389 (98%)	4 (1%)	2 (0%)	25	13
1	BA	400/426 (94%)	389 (97%)	9 (2%)	2 (0%)	25	13
All	All	795/852 (93%)	778 (98%)	13 (2%)	4 (0%)	25	13

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AA	338	LYS
1	BA	322	ALA
1	BA	338	LYS
1	AA	322	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AA	353/376 (94%)	352 (100%)	1 (0%)	91	90
1	BA	358/376 (95%)	357 (100%)	1 (0%)	91	90
All	All	711/752 (94%)	709 (100%)	2 (0%)	91	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AA	22	THR
1	BA	66	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	AA	36	ASN
1	AA	107	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	AA	152	ASN
1	AA	224	ASN
1	AA	227	GLN
1	AA	270	GLN
1	AA	342	GLN
1	BA	318	ASN
1	BA	375	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	IHP	AA	502	-	36,36,36	1.43	6 (16%)	54,60,60	0.68	1 (1%)
2	IHP	AA	501	-	36,36,36	1.50	6 (16%)	54,60,60	0.86	2 (3%)
2	IHP	BA	501	-	36,36,36	1.50	6 (16%)	54,60,60	0.94	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	IHP	AA	502	-	-	2/30/54/54	0/1/1/1
2	IHP	AA	501	-	-	3/30/54/54	0/1/1/1
2	IHP	BA	501	-	-	2/30/54/54	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	AA	501	IHP	P5-O15	3.66	1.66	1.59
2	AA	501	IHP	P6-O16	3.49	1.65	1.59
2	BA	501	IHP	P4-O14	3.48	1.65	1.59
2	BA	501	IHP	P5-O15	3.44	1.65	1.59
2	AA	502	IHP	P2-O12	3.41	1.65	1.59
2	AA	501	IHP	P2-O12	3.31	1.65	1.59
2	AA	501	IHP	P4-O14	3.30	1.65	1.59
2	BA	501	IHP	P6-O16	3.27	1.65	1.59
2	AA	502	IHP	P5-O15	3.17	1.65	1.59
2	AA	502	IHP	P3-O13	3.11	1.65	1.59
2	BA	501	IHP	P1-O11	3.09	1.65	1.59
2	AA	502	IHP	P4-O14	3.05	1.65	1.59
2	AA	501	IHP	P3-O13	3.03	1.65	1.59
2	BA	501	IHP	P3-O13	3.01	1.65	1.59
2	BA	501	IHP	P2-O12	2.88	1.64	1.59
2	AA	502	IHP	P6-O16	2.82	1.64	1.59
2	AA	502	IHP	P1-O11	2.60	1.64	1.59
2	AA	501	IHP	P1-O11	2.39	1.63	1.59

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BA	501	IHP	C3-C2-C1	2.43	115.74	110.41
2	AA	501	IHP	C4-C3-C2	2.36	115.57	110.41
2	BA	501	IHP	O15-C5-C4	2.08	113.59	108.69
2	BA	501	IHP	O13-C3-C2	2.07	113.57	108.69
2	AA	501	IHP	C6-C1-C2	2.05	114.90	110.41
2	AA	502	IHP	C3-C2-C1	-2.02	105.99	110.41

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	AA	501	IHP	C1-O11-P1-O21
2	AA	501	IHP	C3-O13-P3-O33
2	AA	501	IHP	C6-O16-P6-O46
2	AA	502	IHP	C2-O12-P2-O22
2	BA	501	IHP	C3-O13-P3-O33
2	AA	502	IHP	C2-O12-P2-O32
2	BA	501	IHP	C2-O12-P2-O32

There are no ring outliers.


3 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AA	502	IHP	1	0
2	AA	501	IHP	3	0
2	BA	501	IHP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand IHP AA 501





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AA	393/426 (92%)	-0.34	3 (0%) 82 85	12, 22, 41, 55	6 (1%)
1	BA	394/426 (92%)	-0.24	7 (1%) 67 70	11, 23, 46, 66	10 (2%)
All	All	787/852 (92%)	-0.29	10 (1%) 74 78	11, 22, 43, 66	16 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AA	38	THR	3.4
1	AA	378[A]	TRP	3.2
1	BA	38	THR	3.0
1	BA	48	MET	2.7
1	BA	228	TYR	2.7
1	BA	230	ASP	2.4
1	AA	82	GLN	2.1
1	BA	226	GLU	2.1
1	BA	338	LYS	2.1
1	BA	224	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	IHP	BA	501	36/36	0.83	0.15	16,49,74,83	36
2	IHP	AA	502	36/36	0.85	0.13	37,59,92,96	36
2	IHP	AA	501	36/36	0.87	0.13	20,48,68,72	36

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around IHP BA 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around IHP AA 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around IHP AA 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.