



# Full wwPDB NMR Structure Validation Report ⓘ

Jun 26, 2024 – 01:32 AM EDT

PDB ID : 6R9Z  
BMRB ID : 34391  
Title : 3D NMR solution structure of ligand peptide (Ac)EVNPPVP of Pro-Pro  
          endopeptidase-1  
Authors : Diaz, D.  
Deposited on : 2019-04-04

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLUTION NMR*

The overall completeness of chemical shifts assignment is 40%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	9	

## 2 Ensemble composition and analysis ⓘ

This entry contains 20 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms ( 4) was below the domain threshold value ( 8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 112 atoms, of which 56 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2.


Mol	Chain	Residues	Atoms					Trace
1	A	9	Total	C	H	N	O	1
			112	36	56	9	11	

## 4 Residue-property plots

### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2




### 4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


#### 4.2.1 Score per residue for model 1

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2



#### 4.2.2 Score per residue for model 2

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2



### 4.2.3 Score per residue for model 3


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  100%

There are no outlier residues in this chain.

### 4.2.4 Score per residue for model 4

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



### 4.2.5 Score per residue for model 5


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  100%

There are no outlier residues in this chain.

### 4.2.6 Score per residue for model 6

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



### 4.2.7 Score per residue for model 7


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  100%

There are no outlier residues in this chain.

#### 4.2.8 Score per residue for model 8

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  78% 22%



#### 4.2.9 Score per residue for model 9


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  100%

There are no outlier residues in this chain.

#### 4.2.10 Score per residue for model 10

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  89% 11%



#### 4.2.11 Score per residue for model 11


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  89% 11%



#### 4.2.12 Score per residue for model 12


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



#### 4.2.13 Score per residue for model 13

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  89% 11%



#### 4.2.14 Score per residue for model 14

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



#### 4.2.15 Score per residue for model 15

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  100%

There are no outlier residues in this chain.

#### 4.2.16 Score per residue for model 16


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  89% 11%



#### 4.2.17 Score per residue for model 17

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2


Chain A:  89% 11%





#### 4.2.18 Score per residue for model 18


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



#### 4.2.19 Score per residue for model 19


- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



#### 4.2.20 Score per residue for model 20

- Molecule 1: ACE-GLU-VAL-ASN-PRO-PRO-VAL-PRO-NH2

Chain A:  89% 11%



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	74
Number of shifts mapped to atoms	36
Number of unparsed shifts	0
Number of shifts with mapping errors	38
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	40%

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	1120	1120	1080	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

### 6.3 Torsion angles [i](#)

#### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	7/9 (78%)	5±1 (71±10%)	2±1 (28±11%)	0±0 (1±3%)	26	73
All	All	140/180 (78%)	100 (71%)	39 (28%)	1 (1%)	26	73

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	3	VAL	1

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	7/7 (100%)	6±0 (89±6%)	1±0 (11±6%)	10	54
All	All	140/140 (100%)	125 (89%)	15 (11%)	10	54

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	4	ASN	15

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 40% for the well-defined parts and 40% for the entire structure.

### 7.1 Chemical shift list 1

File name: `working_cs.cif`

Chemical shift list name: *starch\_output*

#### 7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	74
Number of shifts mapped to atoms	36
Number of unparsed shifts	0
Number of shifts with mapping errors	38
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

The following assigned chemical shifts were not mapped to the molecules present in the coordinate file.

- No matching atom found in the structure. All 38 occurrences are reported below.

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	2	GLU	2HB	1.987	0.020	2
1	A	2	GLU	3HB	1.884	0.020	2
1	A	2	GLU	2HG	2.236	0.020	1
1	A	2	GLU	3HG	2.236	0.020	1
1	A	3	VAL	1HG1	0.914	0.020	1
1	A	3	VAL	2HG1	0.914	0.020	1
1	A	3	VAL	3HG1	0.914	0.020	1
1	A	3	VAL	1HG2	0.914	0.020	1
1	A	3	VAL	2HG2	0.914	0.020	1
1	A	3	VAL	3HG2	0.914	0.020	1
1	A	4	ASN	2HB	2.777	0.020	2
1	A	4	ASN	3HB	2.628	0.020	2
1	A	4	ASN	1HD2	6.942	0.020	1
1	A	4	ASN	2HD2	7.6	0.020	1

*Continued on next page...*

*Continued from previous page...*

List ID	Chain	Res	Type	Atom	Shift Data		
					Value	Uncertainty	Ambiguity
1	A	5	PRO	2HB	2.324	0.020	2
1	A	5	PRO	3HB	1.925	0.020	2
1	A	5	PRO	2HG	2.033	0.020	1
1	A	5	PRO	3HG	2.033	0.020	1
1	A	5	PRO	2HD	3.793	0.020	2
1	A	5	PRO	3HD	3.691	0.020	2
1	A	6	PRO	2HB	2.265	0.020	2
1	A	6	PRO	3HB	1.867	0.020	2
1	A	6	PRO	2HG	1.997	0.020	2
1	A	6	PRO	3HG	1.888	0.020	2
1	A	6	PRO	2HD	3.805	0.020	2
1	A	6	PRO	3HD	3.649	0.020	2
1	A	7	VAL	1HG1	0.967	0.020	1
1	A	7	VAL	2HG1	0.967	0.020	1
1	A	7	VAL	3HG1	0.967	0.020	1
1	A	7	VAL	1HG2	0.967	0.020	1
1	A	7	VAL	2HG2	0.967	0.020	1
1	A	7	VAL	3HG2	0.967	0.020	1
1	A	8	PRO	2HB	2.305	0.020	2
1	A	8	PRO	3HB	1.923	0.020	2
1	A	8	PRO	2HG	2.051	0.020	1
1	A	8	PRO	3HG	2.051	0.020	1
1	A	8	PRO	2HD	3.852	0.020	2
1	A	8	PRO	3HD	3.693	0.020	2

### 7.1.2 Chemical shift referencing ⓘ

No chemical shift referencing corrections were calculated (not enough data).

### 7.1.3 Completeness of resonance assignments ⓘ

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 40%, i.e. 36 atoms were assigned a chemical shift out of a possible 90. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	<sup>1</sup> H	<sup>13</sup> C	<sup>15</sup> N
Backbone	18/29 (62%)	11/11 (100%)	7/14 (50%)	0/4 (0%)
Sidechain	18/61 (30%)	2/40 (5%)	16/20 (80%)	0/1 (0%)
Overall	36/90 (40%)	13/51 (25%)	23/34 (68%)	0/5 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 40%, i.e. 36 atoms were assigned a chemical shift out of a possible 90. 0 out of 2 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^1\text{H}$	$^{13}\text{C}$	$^{15}\text{N}$
Backbone	18/29 (62%)	11/11 (100%)	7/14 (50%)	0/4 (0%)
Sidechain	18/61 (30%)	2/40 (5%)	16/20 (80%)	0/1 (0%)
Overall	36/90 (40%)	13/51 (25%)	23/34 (68%)	0/5 (0%)

#### 7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

#### 7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

