



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 12, 2024 – 10:04 PM EDT

PDB ID : 1RBO  
Title : SPINACH RUBISCO IN COMPLEX WITH THE INHIBITOR 2-CARBOX  
YARABINITOL-1,5-DIPHOSPHATE  
Authors : Taylor, T.C.; Andersson, I.  
Deposited on : 1996-10-31  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	475	<div> <div></div> <div>86%</div> <div>11%</div> <div></div> </div>
1	E	475	<div> <div></div> <div>87%</div> <div>11%</div> <div></div> </div>
1	H	475	<div> <div></div> <div>87%</div> <div>11%</div> <div></div> </div>
1	L	475	<div> <div></div> <div>87%</div> <div>11%</div> <div></div> </div>
2	C	123	<div> <div></div> <div>78%</div> <div>18%</div> <div></div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	123	
2	I	123	
2	S	123	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	CAP	B	476	X	-	-	-
3	CAP	E	476	X	-	-	-
3	CAP	H	476	X	-	-	-
3	CAP	L	476	X	-	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 19900 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGEN ASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	467	Total	C	N	O	S	0	0	0
			3649	2315	640	676	18			
1	B	467	Total	C	N	O	S	0	0	0
			3649	2315	640	676	18			
1	E	467	Total	C	N	O	S	0	0	0
			3649	2315	640	676	18			
1	H	467	Total	C	N	O	S	0	0	0
			3649	2315	640	676	18			

- Molecule 2 is a protein called RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGEN ASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	S	123	Total	C	N	O	S	0	0	0
			1033	673	167	186	7			
2	C	123	Total	C	N	O	S	0	0	0
			1033	673	167	186	7			
2	F	123	Total	C	N	O	S	0	0	0
			1033	673	167	186	7			
2	I	123	Total	C	N	O	S	0	0	0
			1033	673	167	186	7			

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
S	2	GLN	LYS	conflict	UNP P00870
S	6	ILE	THR	conflict	UNP P00870
S	7	LEU	GLN	conflict	UNP P00870
S	9	LEU	MET	conflict	UNP P00870
S	11	LYS	ARG	conflict	UNP P00870
S	109	GLU	GLN	conflict	UNP P00870
S	113	ILE	VAL	conflict	UNP P00870

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	2	GLN	LYS	conflict	UNP P00870
C	6	ILE	THR	conflict	UNP P00870
C	7	LEU	GLN	conflict	UNP P00870
C	9	LEU	MET	conflict	UNP P00870
C	11	LYS	ARG	conflict	UNP P00870
C	109	GLU	GLN	conflict	UNP P00870
C	113	ILE	VAL	conflict	UNP P00870
F	2	GLN	LYS	conflict	UNP P00870
F	6	ILE	THR	conflict	UNP P00870
F	7	LEU	GLN	conflict	UNP P00870
F	9	LEU	MET	conflict	UNP P00870
F	11	LYS	ARG	conflict	UNP P00870
F	109	GLU	GLN	conflict	UNP P00870
F	113	ILE	VAL	conflict	UNP P00870
I	2	GLN	LYS	conflict	UNP P00870
I	6	ILE	THR	conflict	UNP P00870
I	7	LEU	GLN	conflict	UNP P00870
I	9	LEU	MET	conflict	UNP P00870
I	11	LYS	ARG	conflict	UNP P00870
I	109	GLU	GLN	conflict	UNP P00870
I	113	ILE	VAL	conflict	UNP P00870

- Molecule 3 is 2-CARBOXYARABINITOL-1,5-DIPHOSPHATE (three-letter code: CAP) (formula:  $C_6H_{14}O_{13}P_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total C O P 21 6 13 2	0	0
3	B	1	Total C O P 21 6 13 2	0	0
3	E	1	Total C O P 21 6 13 2	0	0
3	H	1	Total C O P 21 6 13 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	229	Total O 229 229	0	0
4	S	44	Total O 44 44	0	0
4	B	227	Total O 227 227	0	0
4	C	43	Total O 43 43	0	0
4	E	229	Total O 229 229	0	0
4	F	45	Total O 45 45	0	0
4	H	227	Total O 227 227	0	0
4	I	44	Total O 44 44	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE




#### • Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE



#### • Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE





#### • Molecule 1: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE









- Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE



- Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE



- Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE



- Molecule 2: RIBULOSE BISPHOSPHATE CARBOXYLASE/OXYGENASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.20Å 157.20Å 201.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.30 7.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.1 (7.00-2.30) 95.4 (7.00-2.30)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	12.33 (at 2.30Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.154 , 0.171 0.144 , 0.141	Depositor DCC
$R_{free}$ test set	5279 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	10.9	Xtriage
Anisotropy	0.184	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 59.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.038 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	19900	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 31.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1607e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.52	0/3739	0.75	2/5071 (0.0%)
1	E	0.52	0/3739	0.75	2/5071 (0.0%)
1	H	0.52	0/3739	0.75	2/5071 (0.0%)
1	L	0.52	0/3739	0.75	2/5071 (0.0%)
2	C	0.52	0/1068	0.73	1/1453 (0.1%)
2	F	0.52	0/1068	0.72	1/1453 (0.1%)
2	I	0.52	0/1068	0.73	1/1453 (0.1%)
2	S	0.52	0/1068	0.72	1/1453 (0.1%)
All	All	0.52	0/19228	0.74	12/26096 (0.0%)

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	33	LEU	CA-CB-CG	7.30	132.09	115.30
2	C	33	LEU	CA-CB-CG	7.30	132.08	115.30
2	S	33	LEU	CA-CB-CG	7.29	132.06	115.30
2	I	33	LEU	CA-CB-CG	7.28	132.05	115.30
1	B	325	HIS	CB-CA-C	-5.46	99.49	110.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3649	0	3565	35	1
1	E	3649	0	3565	34	1
1	H	3649	0	3565	35	2
1	L	3649	0	3565	35	1
2	C	1033	0	990	26	0
2	F	1033	0	990	27	0
2	I	1033	0	990	26	2
2	S	1033	0	990	26	0
3	B	21	0	9	0	0
3	E	21	0	9	0	0
3	H	21	0	9	0	0
3	L	21	0	9	0	0
4	B	227	0	0	3	15
4	C	43	0	0	1	1
4	E	229	0	0	3	22
4	F	45	0	0	1	1
4	H	227	0	0	3	16
4	I	44	0	0	1	1
4	L	229	0	0	3	24
4	S	44	0	0	1	1
All	All	19900	0	18256	222	57

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 222 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:89:GLU:HG3	2:I:92:LYS:NZ	1.85	0.92
2:F:89:GLU:HG3	2:F:92:LYS:NZ	1.85	0.91
2:C:89:GLU:HG3	2:C:92:LYS:NZ	1.85	0.91
2:S:89:GLU:HG3	2:S:92:LYS:NZ	1.85	0.89
1:H:267:HIS:HD2	1:H:277:ASN:HD22	1.23	0.87

The worst 5 of 57 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:645:HOH:O	4:H:646:HOH:O[3_555]	0.69	1.51
4:L:642:HOH:O	4:E:646:HOH:O[3_555]	0.71	1.49
4:L:643:HOH:O	4:E:645:HOH:O[3_555]	0.72	1.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:523:HOH:O	4:B:523:HOH:O[3_555]	0.75	1.45
4:B:644:HOH:O	4:B:645:HOH:O[3_555]	0.75	1.45

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	465/475 (98%)	452 (97%)	13 (3%)	0	100	100
1	E	465/475 (98%)	452 (97%)	13 (3%)	0	100	100
1	H	465/475 (98%)	452 (97%)	13 (3%)	0	100	100
1	L	465/475 (98%)	452 (97%)	13 (3%)	0	100	100
2	C	121/123 (98%)	117 (97%)	4 (3%)	0	100	100
2	F	121/123 (98%)	117 (97%)	4 (3%)	0	100	100
2	I	121/123 (98%)	117 (97%)	4 (3%)	0	100	100
2	S	121/123 (98%)	117 (97%)	4 (3%)	0	100	100
All	All	2344/2392 (98%)	2276 (97%)	68 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	378/386 (98%)	366 (97%)	12 (3%)	34	50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	378/386 (98%)	366 (97%)	12 (3%)	34	50
1	H	378/386 (98%)	366 (97%)	12 (3%)	34	50
1	L	378/386 (98%)	366 (97%)	12 (3%)	34	50
2	C	112/112 (100%)	104 (93%)	8 (7%)	12	17
2	F	112/112 (100%)	104 (93%)	8 (7%)	12	17
2	I	112/112 (100%)	104 (93%)	8 (7%)	12	17
2	S	112/112 (100%)	104 (93%)	8 (7%)	12	17
All	All	1960/1992 (98%)	1880 (96%)	80 (4%)	26	39

5 of 80 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	68	THR
1	H	466	LYS
2	F	92	LYS
1	H	94	GLU
2	I	47	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 68 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	238	HIS
1	H	267	HIS
1	H	432	ASN
1	B	241	ASN
1	B	238	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	CAP	H	476	-	18,20,20	1.71	2 (11%)	23,31,31	1.60	4 (17%)
3	CAP	E	476	-	18,20,20	1.72	2 (11%)	23,31,31	1.61	4 (17%)
3	CAP	B	476	-	18,20,20	1.72	2 (11%)	23,31,31	1.61	4 (17%)
3	CAP	L	476	-	18,20,20	1.71	2 (11%)	23,31,31	1.61	4 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CAP	H	476	-	1/1/6/6	3/29/29/29	-
3	CAP	E	476	-	1/1/6/6	3/29/29/29	-
3	CAP	B	476	-	1/1/6/6	3/29/29/29	-
3	CAP	L	476	-	1/1/6/6	3/29/29/29	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	476	CAP	C4-C3	4.27	1.58	1.54
3	E	476	CAP	C4-C3	4.25	1.58	1.54
3	L	476	CAP	C4-C3	4.21	1.58	1.54
3	H	476	CAP	C4-C3	4.15	1.57	1.54
3	H	476	CAP	C2-C	3.53	1.57	1.53

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	476	CAP	O7-C-C2	4.40	121.42	114.06
3	B	476	CAP	O7-C-C2	4.38	121.39	114.06
3	L	476	CAP	O7-C-C2	4.37	121.38	114.06
3	H	476	CAP	O7-C-C2	4.37	121.38	114.06
3	B	476	CAP	O6-C-C2	-3.25	116.27	122.32

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	L	476	CAP	C3
3	B	476	CAP	C3
3	E	476	CAP	C3
3	H	476	CAP	C3


5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	L	476	CAP	O6-C-C2-C3
3	B	476	CAP	O6-C-C2-C3
3	E	476	CAP	O6-C-C2-C3
3	H	476	CAP	O6-C-C2-C3
3	L	476	CAP	O2-C2-C3-C4


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	467/475 (98%)	-1.00	4 (0%) 81 81	2, 6, 24, 52	0
1	E	467/475 (98%)	-1.01	4 (0%) 81 81	2, 6, 24, 52	0
1	H	467/475 (98%)	-0.98	5 (1%) 77 78	2, 6, 24, 52	0
1	L	467/475 (98%)	-1.01	6 (1%) 74 75	2, 6, 24, 52	0
2	C	123/123 (100%)	-0.62	0 100 100	3, 15, 30, 38	0
2	F	123/123 (100%)	-0.71	0 100 100	3, 15, 30, 38	0
2	I	123/123 (100%)	-0.64	0 100 100	3, 15, 30, 38	0
2	S	123/123 (100%)	-0.67	0 100 100	3, 15, 30, 38	0
All	All	2360/2392 (98%)	-0.93	19 (0%) 82 83	2, 7, 28, 52	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	9	ALA	5.0
1	L	9	ALA	4.7
1	E	10	SER	4.2
1	H	9	ALA	4.0
1	H	92	GLY	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CAP	B	476	21/21	0.98	0.06	3,9,15,20	0
3	CAP	E	476	21/21	0.98	0.05	3,9,15,20	0
3	CAP	H	476	21/21	0.98	0.05	3,9,15,20	0
3	CAP	L	476	21/21	0.99	0.05	3,9,15,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







**Electron density around CAP E 476:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around CAP H 476:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.