



wwPDB EM Validation Summary Report ⓘ

May 18, 2024 – 03:59 pm BST

PDB ID : 6RBD
EMDB ID : EMD-4792
Title : State 1 of yeast Tsr1-TAP Rps20-Deltaloop pre-40S particles
Authors : Shayan, R.; Mitterer, V.; Ferreira-Cerca, S.; Murat, G.; Enne, T.; Rinaldi, D.; Weigl, S.; Omanic, H.; Gleizes, P.E.; Kressler, D.; Pertschy, B.; Plisson-Chastang, C.
Deposited on : 2019-04-10
Resolution : 3.47 Å (reported)
Based on initial models : 6FAI, 6EML, ?

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























| Metric | Whole archive (#Entries) | EM structures (#Entries) |
|-----------------------|-----------------------------|-----------------------------|
| Ramachandran outliers | 154571 | 4023 |
| Sidechain outliers | 154315 | 3826 |
| RNA backbone | 4643 | 859 |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | b | 82 | 98% .. |
| 2 | e | 63 | 8% 75% 24% . |
| 3 | 2 | 1801 | 6% 60% 34% . . |
| 4 | A | 252 | 81% 18% . |
| 5 | B | 255 | 84% 16% . |
| 6 | C | 254 | 84% 15% . |
| 7 | E | 261 | 98% . |
| 8 | G | 236 | 98% . |



Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 9 | H | 190 |  |
| 10 | I | 200 |  |
| 11 | J | 197 |  |
| 12 | L | 156 |  |
| 13 | N | 151 |  |
| 14 | V | 87 |  |
| 15 | W | 130 |  |
| 16 | X | 145 |  |
| 17 | Y | 135 |  |
| 18 | c | 67 |  |
| 19 | d | 56 |  |
| 20 | h | 274 |  |
| 21 | i | 483 |  |
| 22 | k | 788 |  |
| 23 | l | 425 |  |
| 24 | y | 318 |  |
| 25 | F | 225 |  |
| 26 | M | 143 |  |
| 27 | O | 137 |  |
| 28 | P | 142 |  |
| 29 | Q | 143 |  |
| 30 | R | 136 |  |
| 31 | S | 146 |  |
| 32 | T | 144 |  |
| 33 | Z | 108 |  |

Continued on next page...

Continued from previous page...

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|--|
| 34 | D | 240 |  |
| 35 | U | 121 |  |

2 Entry composition

There are 35 unique types of molecules in this entry. The entry contains 81048 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 40S ribosomal protein S27-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 1 | b | 81 | Total | C | N | O | S | 0 | 0 |
| | | | 610 | 382 | 110 | 113 | 5 | | |

- Molecule 2 is a protein called 40S ribosomal protein S30-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 2 | e | 48 | Total | C | N | O | S | 0 | 0 |
| | | | 384 | 242 | 81 | 59 | 2 | | |

- Molecule 3 is a RNA chain called 20S ribosomal RNA.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-------|------|-------|------|---------|-------|
| 3 | 2 | 1777 | Total | C | N | O | P | 0 | 0 |
| | | | 37868 | 16929 | 6704 | 12458 | 1777 | | |

- Molecule 4 is a protein called 40S ribosomal protein S0-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 4 | A | 206 | Total | C | N | O | S | 0 | 0 |
| | | | 1611 | 1036 | 285 | 288 | 2 | | |

There is a discrepancy between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment | Reference |
|-------|---------|----------|--------|----------|------------|
| A | 206 | PHE | ASP | conflict | UNP P32905 |

- Molecule 5 is a protein called 40S ribosomal protein S1-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 5 | B | 214 | Total | C | N | O | S | 0 | 0 |
| | | | 1709 | 1084 | 310 | 311 | 4 | | |

- Molecule 6 is a protein called 40S ribosomal protein S2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 6 | C | 217 | Total | C | N | O | S | 0 | 0 |
| | | | 1635 | 1047 | 289 | 297 | 2 | | |

- Molecule 7 is a protein called 40S ribosomal protein S4-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 7 | E | 260 | Total | C | N | O | S | 0 | 0 |
| | | | 2068 | 1316 | 389 | 360 | 3 | | |

- Molecule 8 is a protein called 40S ribosomal protein S6-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 8 | G | 232 | Total | C | N | O | S | 0 | 0 |
| | | | 1873 | 1172 | 366 | 332 | 3 | | |

- Molecule 9 is a protein called 40S ribosomal protein S7-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 9 | H | 184 | Total | C | N | O | S | 0 | 0 |
| | | | 1481 | 951 | 265 | 265 | | | |

- Molecule 10 is a protein called 40S ribosomal protein S8-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 10 | I | 188 | Total | C | N | O | S | 0 | 0 |
| | | | 1489 | 925 | 298 | 264 | 2 | | |

- Molecule 11 is a protein called 40S ribosomal protein S9-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 11 | J | 185 | Total | C | N | O | S | 0 | 0 |
| | | | 1494 | 943 | 289 | 261 | 1 | | |

- Molecule 12 is a protein called 40S ribosomal protein S11-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 12 | L | 140 | Total | C | N | O | S | 0 | 0 |
| | | | 1129 | 724 | 215 | 187 | 3 | | |

- Molecule 13 is a protein called 40S ribosomal protein S13.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 13 | N | 150 | Total | C | N | O | S | 0 | 0 |
| | | | 1192 | 759 | 224 | 207 | 2 | | |

- Molecule 14 is a protein called 40S ribosomal protein S21-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 14 | V | 87 | Total | C | N | O | S | 0 | 0 |
| | | | 684 | 420 | 125 | 137 | 2 | | |

- Molecule 15 is a protein called 40S ribosomal protein S22-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 15 | W | 129 | Total | C | N | O | S | 0 | 0 |
| | | | 1021 | 650 | 188 | 180 | 3 | | |

- Molecule 16 is a protein called 40S ribosomal protein S23-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 16 | X | 144 | Total | C | N | O | S | 0 | 0 |
| | | | 1121 | 708 | 220 | 191 | 2 | | |

- Molecule 17 is a protein called 40S ribosomal protein S24-A.

| Mol | Chain | Residues | Atoms | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| 17 | Y | 134 | Total | C | N | O | 0 | 0 |
| | | | 1073 | 676 | 208 | 189 | | |

- Molecule 18 is a protein called 40S ribosomal protein S28-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 18 | c | 63 | Total | C | N | O | S | 0 | 0 |
| | | | 497 | 306 | 99 | 91 | 1 | | |

- Molecule 19 is a protein called 40S ribosomal protein S29-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 19 | d | 37 | Total | C | N | O | S | 0 | 0 |
| | | | 302 | 186 | 62 | 50 | 4 | | |

- Molecule 20 is a protein called Pre-rRNA-processing protein PNO1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 20 | h | 181 | Total | C | N | O | S | 0 | 0 |
| | | | 1436 | 917 | 261 | 254 | 4 | | |

- Molecule 21 is a protein called Essential nuclear protein 1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 21 | i | 139 | Total | C | N | O | S | 0 | 0 |
| | | | 1116 | 731 | 181 | 202 | 2 | | |

- Molecule 22 is a protein called Ribosome biogenesis protein TSR1.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 22 | k | 605 | Total | C | N | O | S | 0 | 0 |
| | | | 4860 | 3112 | 849 | 885 | 14 | | |

- Molecule 23 is a protein called Serine/threonine-protein kinase RIO2.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 23 | l | 271 | Total | C | N | O | S | 0 | 0 |
| | | | 2199 | 1390 | 385 | 407 | 17 | | |

- Molecule 24 is a protein called Dimethyladenosine transferase.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 24 | y | 154 | Total | C | N | O | S | 0 | 0 |
| | | | 1245 | 788 | 223 | 225 | 9 | | |

- Molecule 25 is a protein called 40S ribosomal protein S5.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|---|---------|-------|
| 25 | F | 206 | Total | C | N | O | S | 0 | 0 |
| | | | 1609 | 1007 | 300 | 299 | 3 | | |

- Molecule 26 is a protein called 40S ribosomal protein S12.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 26 | M | 125 | Total | C | N | O | S | 0 | 0 |
| | | | 941 | 591 | 166 | 182 | 2 | | |

- Molecule 27 is a protein called 40S ribosomal protein S14-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 27 | O | 127 | Total | C | N | O | S | 0 | 0 |
| | | | 926 | 569 | 185 | 169 | 3 | | |

- Molecule 28 is a protein called 40S ribosomal protein S15.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 28 | P | 114 | Total | C | N | O | S | 0 | 0 |
| | | | 902 | 574 | 166 | 155 | 7 | | |

- Molecule 29 is a protein called 40S ribosomal protein S16-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 29 | Q | 127 | Total | C | N | O | S | 0 | 0 |
| | | | 993 | 640 | 177 | 176 | | | |

- Molecule 30 is a protein called 40S ribosomal protein S17-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 30 | R | 125 | Total | C | N | O | S | 0 | 0 |
| | | | 1000 | 625 | 188 | 185 | 2 | | |

- Molecule 31 is a protein called 40S ribosomal protein S18-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 31 | S | 104 | Total | C | N | O | S | 0 | 0 |
| | | | 840 | 528 | 154 | 156 | 2 | | |

- Molecule 32 is a protein called 40S ribosomal protein S19-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 32 | T | 143 | Total | C | N | O | S | 0 | 0 |
| | | | 1112 | 694 | 208 | 208 | 2 | | |

- Molecule 33 is a protein called 40S ribosomal protein S25-A.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|----|----|---|---------|-------|
| 33 | Z | 63 | Total | C | N | O | S | 0 | 0 |
| | | | 512 | 328 | 94 | 90 | | | |

- Molecule 34 is a protein called 40S ribosomal protein S3.

| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 34 | D | 185 | Total | C | N | O | S | 0 | 0 |
| | | | 1442 | 911 | 266 | 259 | 6 | | |

- Molecule 35 is a protein called 40S ribosomal protein S20.


| Mol | Chain | Residues | Atoms | | | | | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 35 | U | 82 | Total | C | N | O | S | 0 | 0 |
| | | | 674 | 431 | 123 | 119 | 1 | | |

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 40S ribosomal protein S27-A

Chain b:  98%




- Molecule 2: 40S ribosomal protein S30-A

Chain e:  8% 75% 24%





- Molecule 3: 20S ribosomal RNA

Chain 2:  6% 60% 34%







Chain A:  81% 18%




- Molecule 5: 40S ribosomal protein S1-A

Chain B:  84% 16%



- Molecule 6: 40S ribosomal protein S2

Chain C:  84% 15%




- Molecule 7: 40S ribosomal protein S4-A

Chain E:  98%




- Molecule 8: 40S ribosomal protein S6-A

Chain G:  98%




- Molecule 9: 40S ribosomal protein S7-A

Chain H:  97%




- Molecule 10: 40S ribosomal protein S8-A

Chain I:  94% 6%





- Molecule 11: 40S ribosomal protein S9-A

Chain J:  93% • 6%




- Molecule 12: 40S ribosomal protein S11-A

Chain L:  88% • 10%



- Molecule 13: 40S ribosomal protein S13

Chain N:  99% •



- Molecule 14: 40S ribosomal protein S21-A

Chain V:  100%

There are no outlier residues recorded for this chain.


- Molecule 15: 40S ribosomal protein S22-A

Chain W:  96% • •



- Molecule 16: 40S ribosomal protein S23-A

Chain X:  6% 98% • •



- Molecule 17: 40S ribosomal protein S24-A

Chain Y:  99% •




-
- Diagram illustrating the structure of the 12S ribosomal subunit, showing various proteins (MET, ASP, ASN, LYS, T5, G17, R18, L33, E34, D35, T36, S37, R64, R65, L66, R67) and their interactions, marked by red diamonds.


- MET**
- ALA**
- HIS**
- GLU**
- ASN**
- VAL**
- TRP**
- PHE**
- SER**
- HIS**
- PRO**
- ARG**
- ARG**
- TYR**
- GLY**
- LYS**
- GLY**
- ARG**
- R20**
- C21**
- R22**
- V23**
- T28**
- G35**
- L36**
- N37**
- A47**
- N48**
- D49**
- I50**
- G51**
- F52**
- N53**
- R56**

- | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|------|------|------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| GLY | SER | ARG | ALA | LYS | THR | HIS | GLU | SER | LYS | THR | VAL | VAL | VAL | ASP | Gln | GLY | LYS | PRO | ARG | PHE | THR | SER | ALA | ALA | SER | LYS | THR | Gln | ASN | GLY | ASN | LYS | Ile | K93 | R221 | E272 | R273 | Tyr | | | | | | | | | |
| MET | VAL | ALA | PRO | THR | ALA | LEU | LYS | ALA | THR | VAL | THR | PRO | VAL | SER | GLY | Gln | ASP | GLY | ILE | GLY | ILE | ASN | ASN | THR | GLU | SER | ILE | ASP | GLU | ASP | ASP | ASP | ASP | VAL | LEU | ASP | SER | ASN | ASN | THR | ALA | LYS | GLU | VAL | GLU | GLY | GLU |

- [illegible]


- 

Chain k: 




• Molecule 23: Serine/threonine-protein kinase RIO2


Chain l: 




• Molecule 24: Dimethyladenosine transferase

Chain y: 








- Molecule 25: 40S ribosomal protein S5



- Molecule 26: 40S ribosomal protein S12





- Molecule 27: 40S ribosomal protein S14-A




- Molecule 28: 40S ribosomal protein S15




- Molecule 29: 40S ribosomal protein S16-A




- Molecule 30: 40S ribosomal protein S17-A







- Molecule 31: 40S ribosomal protein S18-A





- Molecule 32: 40S ribosomal protein S19-A




- Molecule 33: 40S ribosomal protein S25-A



- Molecule 34: 40S ribosomal protein S3



Chain U:



4 Experimental information

| Property | Value | Source |
|--------------------------------------|---|-----------|
| EM reconstruction method | SINGLE PARTICLE | Depositor |
| Imposed symmetry | POINT, C1 | Depositor |
| Number of particles used | 54130 | Depositor |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor |
| CTF correction method | PHASE FLIPPING AND AMPLITUDE CORRECTION | Depositor |
| Microscope | FEI TITAN KRIOS | Depositor |
| Voltage (kV) | 300 | Depositor |
| Electron dose ($e^-/\text{\AA}^2$) | 32.4 | Depositor |
| Minimum defocus (nm) | 800 | Depositor |
| Maximum defocus (nm) | 3000 | Depositor |
| Magnification | Not provided | |
| Image detector | GATAN K2 SUMMIT (4k x 4k) | Depositor |
| Maximum map value | 0.528 | Depositor |
| Minimum map value | -0.170 | Depositor |
| Average map value | 0.003 | Depositor |
| Map value standard deviation | 0.020 | Depositor |
| Recommended contour level | 0.075 | Depositor |
| Map size (Å) | 409.72803, 409.72803, 409.72803 | wwPDB |
| Map dimensions | 384, 384, 384 | wwPDB |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB |
| Pixel spacing (Å) | 1.067, 1.067, 1.067 | Depositor |

5 Model quality ⓘ

5.1 Standard geometry ⓘ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|------------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 1 | b | 0.32 | 0/620 | 0.55 | 0/838 |
| 2 | e | 0.37 | 0/390 | 0.62 | 0/517 |
| 3 | 2 | 0.58 | 0/42353 | 1.19 | 375/65990 (0.6%) |
| 4 | A | 0.36 | 0/1653 | 0.56 | 1/2261 (0.0%) |
| 5 | B | 0.33 | 0/1735 | 0.59 | 0/2335 |
| 6 | C | 0.36 | 0/1665 | 0.57 | 0/2263 |
| 7 | E | 0.40 | 0/2109 | 0.59 | 0/2839 |
| 8 | G | 0.35 | 0/1897 | 0.52 | 0/2532 |
| 9 | H | 0.31 | 0/1506 | 0.53 | 0/2028 |
| 10 | I | 0.40 | 0/1514 | 0.56 | 0/2021 |
| 11 | J | 0.37 | 0/1519 | 0.57 | 1/2035 (0.0%) |
| 12 | L | 0.43 | 0/1155 | 0.56 | 0/1557 |
| 13 | N | 0.33 | 0/1215 | 0.55 | 0/1638 |
| 14 | V | 0.39 | 0/693 | 0.53 | 0/935 |
| 15 | W | 0.43 | 0/1038 | 0.63 | 0/1395 |
| 16 | X | 0.40 | 0/1139 | 0.59 | 0/1518 |
| 17 | Y | 0.39 | 0/1087 | 0.54 | 0/1449 |
| 18 | c | 0.27 | 0/499 | 0.57 | 0/670 |
| 19 | d | 0.27 | 0/305 | 0.51 | 0/401 |
| 20 | h | 0.30 | 0/1462 | 0.53 | 0/1969 |
| 21 | i | 0.29 | 0/1145 | 0.54 | 0/1556 |
| 22 | k | 0.31 | 0/4970 | 0.53 | 0/6718 |
| 23 | l | 0.29 | 0/2243 | 0.56 | 0/3014 |
| 24 | y | 0.28 | 0/1265 | 0.58 | 0/1702 |
| 25 | F | 0.28 | 0/1629 | 0.55 | 0/2202 |
| 26 | M | 0.28 | 0/949 | 0.55 | 0/1284 |
| 27 | O | 0.31 | 0/937 | 0.53 | 0/1261 |
| 28 | P | 0.28 | 0/921 | 0.51 | 0/1237 |
| 29 | Q | 0.30 | 0/1011 | 0.58 | 0/1362 |
| 30 | R | 0.27 | 0/1010 | 0.55 | 0/1355 |
| 31 | S | 0.28 | 0/853 | 0.55 | 0/1154 |
| 32 | T | 0.30 | 0/1130 | 0.52 | 1/1517 (0.1%) |
| 33 | Z | 0.45 | 0/519 | 0.77 | 0/696 |
| 34 | D | 0.30 | 0/1459 | 0.56 | 0/1957 |

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|---------|-------------|-------------------|
| | | RMSZ | # Z >5 | RMSZ | # Z >5 |
| 35 | U | 0.26 | 0/681 | 0.49 | 0/917 |
| All | All | 0.47 | 0/86276 | 0.95 | 378/125123 (0.3%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 22 | k | 0 | 3 |
| 24 | y | 0 | 2 |
| All | All | 0 | 5 |

There are no bond length outliers.

The worst 5 of 378 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|------|------|-----------|-------|-------------|----------|
| 3 | 2 | 1634 | C | N1-C2-O2 | 13.79 | 127.17 | 118.90 |
| 3 | 2 | 1634 | C | C2-N1-C1' | 12.73 | 132.81 | 118.80 |
| 3 | 2 | 1148 | C | N1-C2-O2 | 12.30 | 126.28 | 118.90 |
| 3 | 2 | 695 | U | N1-C2-O2 | 12.00 | 131.20 | 122.80 |
| 3 | 2 | 1279 | C | N1-C2-O2 | 11.69 | 125.91 | 118.90 |

There are no chirality outliers.

All (5) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|---------|
| 22 | k | 118 | VAL | Peptide |
| 22 | k | 710 | PHE | Peptide |
| 22 | k | 711 | HIS | Peptide |
| 24 | y | 270 | LYS | Peptide |
| 24 | y | 275 | ASP | Peptide |

5.2 Too-close contacts

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|----------|-------------|-----|
| 1 | b | 79/82 (96%) | 71 (90%) | 8 (10%) | 0 | 100 | 100 |
| 2 | e | 44/63 (70%) | 40 (91%) | 4 (9%) | 0 | 100 | 100 |
| 4 | A | 204/252 (81%) | 189 (93%) | 15 (7%) | 0 | 100 | 100 |
| 5 | B | 212/255 (83%) | 197 (93%) | 15 (7%) | 0 | 100 | 100 |
| 6 | C | 215/254 (85%) | 208 (97%) | 7 (3%) | 0 | 100 | 100 |
| 7 | E | 258/261 (99%) | 249 (96%) | 9 (4%) | 0 | 100 | 100 |
| 8 | G | 230/236 (98%) | 224 (97%) | 6 (3%) | 0 | 100 | 100 |
| 9 | H | 182/190 (96%) | 168 (92%) | 14 (8%) | 0 | 100 | 100 |
| 10 | I | 184/200 (92%) | 175 (95%) | 9 (5%) | 0 | 100 | 100 |
| 11 | J | 183/197 (93%) | 172 (94%) | 11 (6%) | 0 | 100 | 100 |
| 12 | L | 138/156 (88%) | 136 (99%) | 2 (1%) | 0 | 100 | 100 |
| 13 | N | 148/151 (98%) | 143 (97%) | 5 (3%) | 0 | 100 | 100 |
| 14 | V | 85/87 (98%) | 83 (98%) | 2 (2%) | 0 | 100 | 100 |
| 15 | W | 127/130 (98%) | 117 (92%) | 10 (8%) | 0 | 100 | 100 |
| 16 | X | 142/145 (98%) | 136 (96%) | 6 (4%) | 0 | 100 | 100 |
| 17 | Y | 132/135 (98%) | 130 (98%) | 2 (2%) | 0 | 100 | 100 |
| 18 | c | 61/67 (91%) | 57 (93%) | 4 (7%) | 0 | 100 | 100 |
| 19 | d | 33/56 (59%) | 29 (88%) | 4 (12%) | 0 | 100 | 100 |
| 20 | h | 179/274 (65%) | 174 (97%) | 5 (3%) | 0 | 100 | 100 |
| 21 | i | 137/483 (28%) | 133 (97%) | 4 (3%) | 0 | 100 | 100 |
| 22 | k | 597/788 (76%) | 562 (94%) | 34 (6%) | 1 (0%) | 47 | 80 |
| 23 | l | 267/425 (63%) | 233 (87%) | 33 (12%) | 1 (0%) | 34 | 70 |
| 24 | y | 150/318 (47%) | 128 (85%) | 22 (15%) | 0 | 100 | 100 |
| 25 | F | 204/225 (91%) | 195 (96%) | 9 (4%) | 0 | 100 | 100 |
| 26 | M | 123/143 (86%) | 117 (95%) | 6 (5%) | 0 | 100 | 100 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 27 | O | 125/137 (91%) | 119 (95%) | 6 (5%) | 0 | 100 | 100 |
| 28 | P | 112/142 (79%) | 108 (96%) | 4 (4%) | 0 | 100 | 100 |
| 29 | Q | 125/143 (87%) | 120 (96%) | 5 (4%) | 0 | 100 | 100 |
| 30 | R | 123/136 (90%) | 120 (98%) | 3 (2%) | 0 | 100 | 100 |
| 31 | S | 102/146 (70%) | 92 (90%) | 10 (10%) | 0 | 100 | 100 |
| 32 | T | 141/144 (98%) | 138 (98%) | 3 (2%) | 0 | 100 | 100 |
| 33 | Z | 61/108 (56%) | 60 (98%) | 1 (2%) | 0 | 100 | 100 |
| 34 | D | 179/240 (75%) | 173 (97%) | 6 (3%) | 0 | 100 | 100 |
| 35 | U | 78/121 (64%) | 76 (97%) | 2 (3%) | 0 | 100 | 100 |
| All | All | 5360/6890 (78%) | 5072 (95%) | 286 (5%) | 2 (0%) | 100 | 100 |

All (2) Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 23 | l | 183 | SER |
| 22 | k | 483 | PRO |

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------------|------------|----------|-------------|-----|
| 1 | b | 70/71 (99%) | 69 (99%) | 1 (1%) | 67 | 85 |
| 2 | e | 40/54 (74%) | 39 (98%) | 1 (2%) | 47 | 74 |
| 4 | A | 171/210 (81%) | 170 (99%) | 1 (1%) | 86 | 94 |
| 5 | B | 191/224 (85%) | 190 (100%) | 1 (0%) | 88 | 95 |
| 6 | C | 176/205 (86%) | 173 (98%) | 3 (2%) | 60 | 82 |
| 7 | E | 221/222 (100%) | 218 (99%) | 3 (1%) | 67 | 85 |
| 8 | G | 198/201 (98%) | 197 (100%) | 1 (0%) | 88 | 95 |
| 9 | H | 165/170 (97%) | 165 (100%) | 0 | 100 | 100 |
| 10 | I | 150/161 (93%) | 149 (99%) | 1 (1%) | 84 | 93 |

Continued on next page...

Continued from previous page...

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|-----------------|------------|----------|-------------|-----|
| 11 | J | 158/166 (95%) | 157 (99%) | 1 (1%) | 86 | 94 |
| 12 | L | 125/137 (91%) | 123 (98%) | 2 (2%) | 62 | 82 |
| 13 | N | 127/128 (99%) | 127 (100%) | 0 | 100 | 100 |
| 14 | V | 74/74 (100%) | 74 (100%) | 0 | 100 | 100 |
| 15 | W | 110/111 (99%) | 106 (96%) | 4 (4%) | 35 | 65 |
| 16 | X | 119/120 (99%) | 117 (98%) | 2 (2%) | 60 | 82 |
| 17 | Y | 112/113 (99%) | 112 (100%) | 0 | 100 | 100 |
| 18 | c | 56/60 (93%) | 56 (100%) | 0 | 100 | 100 |
| 19 | d | 33/49 (67%) | 31 (94%) | 2 (6%) | 18 | 50 |
| 20 | h | 158/238 (66%) | 157 (99%) | 1 (1%) | 86 | 94 |
| 21 | i | 125/424 (30%) | 125 (100%) | 0 | 100 | 100 |
| 22 | k | 523/703 (74%) | 516 (99%) | 7 (1%) | 69 | 86 |
| 23 | l | 246/384 (64%) | 243 (99%) | 3 (1%) | 71 | 87 |
| 24 | y | 141/283 (50%) | 137 (97%) | 4 (3%) | 43 | 72 |
| 25 | F | 173/191 (91%) | 171 (99%) | 2 (1%) | 71 | 87 |
| 26 | M | 101/119 (85%) | 99 (98%) | 2 (2%) | 55 | 79 |
| 27 | O | 91/105 (87%) | 91 (100%) | 0 | 100 | 100 |
| 28 | P | 95/118 (80%) | 94 (99%) | 1 (1%) | 73 | 88 |
| 29 | Q | 107/119 (90%) | 107 (100%) | 0 | 100 | 100 |
| 30 | R | 113/124 (91%) | 112 (99%) | 1 (1%) | 78 | 91 |
| 31 | S | 93/129 (72%) | 93 (100%) | 0 | 100 | 100 |
| 32 | T | 115/116 (99%) | 114 (99%) | 1 (1%) | 78 | 91 |
| 33 | Z | 56/89 (63%) | 55 (98%) | 1 (2%) | 59 | 81 |
| 34 | D | 149/195 (76%) | 147 (99%) | 2 (1%) | 69 | 86 |
| 35 | U | 79/114 (69%) | 79 (100%) | 0 | 100 | 100 |
| All | All | 4661/5927 (79%) | 4613 (99%) | 48 (1%) | 77 | 89 |

5 of 48 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 22 | k | 288 | ASN |
| 24 | y | 166 | ARG |
| 22 | k | 497 | ARG |

Continued on next page...

Continued from previous page...

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 23 | l | 99 | ASN |
| 24 | y | 270 | LYS |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 16 | X | 48 | HIS |
| 25 | F | 104 | ASN |
| 20 | h | 131 | ASN |
| 25 | F | 103 | ASN |
| 31 | S | 87 | ASN |

5.3.3 RNA ⓘ

| Mol | Chain | Analysed | Backbone Outliers | Pucker Outliers |
|-----|-------|-----------------|-------------------|-----------------|
| 3 | 2 | 1772/1801 (98%) | 617 (34%) | 17 (0%) |

5 of 617 RNA backbone outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | 2 | 4 | C |
| 3 | 2 | 25 | C |
| 3 | 2 | 26 | A |
| 3 | 2 | 34 | G |
| 3 | 2 | 42 | G |

5 of 17 RNA pucker outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|------|------|
| 3 | 2 | 1570 | A |
| 3 | 2 | 1652 | C |
| 3 | 2 | 1150 | G |
| 3 | 2 | 1156 | C |
| 3 | 2 | 1229 | G |

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

| Mol | Chain | Number of breaks |
|-----|-------|------------------|
| 34 | D | 2 |
| 3 | 2 | 2 |
| 22 | k | 1 |
| 19 | d | 1 |

The worst 5 of 6 chain breaks are listed below:

| Model | Chain | Residue-1 | Atom-1 | Residue-2 | Atom-2 | Distance (Å) |
|-------|-------|-----------|--------|-----------|--------|--------------|
| 1 | D | 52:ALA | C | 53:THR | N | 3.81 |
| 1 | 2 | 1377:U | O3' | 1378:U | P | 3.34 |
| 1 | 2 | 1219:A | O3' | 1220:C | P | 3.23 |
| 1 | k | 80:ASN | C | 81:GLY | N | 3.23 |
| 1 | d | 31:ILE | C | 32:ARG | N | 3.21 |


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-4792. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 192




Y Index: 192




Z Index: 192


6.2.2 Raw map



X Index: 192



Y Index: 192




Z Index: 192


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 188




Y Index: 191




Z Index: 194


6.3.2 Raw map



X Index: 191



Y Index: 191




Z Index: 194


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.075. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1165 nm³; this corresponds to an approximate mass of 1053 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.288 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.288 \AA^{-1}

8.2 Resolution estimates [i](#)


| Resolution estimate (Å) | Estimation criterion (FSC cut-off) | | |
|---------------------------|------------------------------------|------|----------|
| | 0.143 | 0.5 | Half-bit |
| Reported by author | 3.47 | - | - |
| Author-provided FSC curve | 3.45 | 4.38 | 3.57 |
| Unmasked-calculated* | 4.71 | 9.35 | 5.69 |

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.71 differs from the reported value 3.47 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-4792 and PDB model 6RBD. Per-residue inclusion information can be found in [section 3](#) on [page 11](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.075 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.075).









































































9.4 Atom inclusion [i](#)



At the recommended contour level, 88% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.075) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion | Q-score |
|-------|--|--|
| All |  0.8580 |  0.3010 |
| 2 |  0.8910 |  0.3250 |
| A |  0.9620 |  0.3880 |
| B |  0.8830 |  0.3430 |
| C |  0.9370 |  0.4320 |
| D |  0.3210 |  0.0700 |
| E |  0.9580 |  0.4940 |
| F |  0.8760 |  0.1080 |
| G |  0.9110 |  0.4150 |
| H |  0.8770 |  0.2990 |
| I |  0.9230 |  0.4640 |
| J |  0.9330 |  0.4440 |
| L |  0.9540 |  0.5010 |
| M |  0.5690 |  0.0680 |
| N |  0.9210 |  0.4260 |
| O |  0.8820 |  0.3060 |
| P |  0.7980 |  0.0860 |
| Q |  0.8900 |  0.1090 |
| R |  0.5430 |  0.1460 |
| S |  0.7700 |  0.0940 |
| T |  0.9040 |  0.0940 |
| U |  0.2470 |  0.0500 |
| V |  0.9580 |  0.4180 |
| W |  0.9570 |  0.4660 |
| X |  0.9070 |  0.4370 |
| Y |  0.9360 |  0.4610 |
| Z |  0.8510 |  0.1010 |
| b |  0.9220 |  0.4210 |
| c |  0.7710 |  0.1320 |
| d |  0.5150 |  0.0700 |
| e |  0.7990 |  0.3590 |
| h |  0.8680 |  0.2590 |
| i |  0.6860 |  0.0500 |
| k |  0.8770 |  0.2590 |
| l |  0.7080 |  0.1000 |
| y |  0.6090 |  0.1220 |

