



## wwPDB EM Validation Summary Report ⓘ

Nov 4, 2024 – 01:06 am GMT

PDB ID : 8RH2  
EMDB ID : EMD-19166  
Title : Trimeric HSV-2G gB ectodomain in postfusion conformation with three bound HDIT102 Fab molecules.  
Authors : Kalbermatter, D.; Seyfizadeh, N.; Imhof, T.; Ries, M.; Mueller, C.; Jenner, L.; Blumenschein, E.; Yendrzheyevskiy, A.; Moog, K.; Eckert, D.; Engel, R.; Diebolder, P.; Chami, M.; Krauss, J.; Schaller, T.; Arndt, M.  
Deposited on : 2023-12-14  
Resolution : 3.12 Å(reported)  
Based on initial model : 2gum

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.12 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	703	
1	B	703	
1	C	703	
2	D	452	
2	F	452	
2	H	452	
3	E	212	
3	G	212	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	L	212	 A horizontal bar chart showing the quality of chain L. The bar is divided into three segments: a green segment representing 39%, a yellow segment representing 11%, and a grey segment representing 50%. The percentages are labeled below the bar.

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 19152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein B.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	583	Total	C	N	O	S	0	0
			4710	2974	830	883	23		
1	B	583	Total	C	N	O	S	0	0
			4710	2974	830	883	23		
1	C	583	Total	C	N	O	S	0	0
			4710	2974	830	883	23		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	553	ALA	VAL	engineered mutation	UNP A0A0D4CHI5
B	553	ALA	VAL	engineered mutation	UNP A0A0D4CHI5
C	553	ALA	VAL	engineered mutation	UNP A0A0D4CHI5

- Molecule 2 is a protein called HDIT102 Fab heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	H	119	Total	C	N	O	S	0	0
			901	556	166	172	7		
2	D	119	Total	C	N	O	S	0	0
			901	556	166	172	7		
2	F	119	Total	C	N	O	S	0	0
			901	556	166	172	7		


- Molecule 3 is a protein called HDIT102 Fab light chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	L	105	Total	C	N	O	S	0	0
			773	482	134	155	2		
3	E	105	Total	C	N	O	S	0	0
			773	482	134	155	2		
3	G	105	Total	C	N	O	S	0	0
			773	482	134	155	2		


### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Envelope glycoprotein B




#### • Molecule 1: Envelope glycoprotein B






• Molecule 1: Envelope glycoprotein B

Chain C: 70% 13% 17%



• Molecule 2: HDIT102 Fab heavy chain

Chain H: 21% 6% 74%





TYR  
SER  
LEU  
LYS  
LEU  
THR  
VAL  
ASP  
LYS  
SER  
ARG  
TRP  
GLN  
GLN  
GLY  
ASN  
VAL  
PHE  
SER  
SER  
CYS  
SER  
VAL  
VAL  
MET  
HIS  
HIS  
GLU  
ALA  
LEU  
HIS  
ASN  
ASN  
HIS  
TYR  
THR  
GLN  
LYS  
SER  
LEU  
SER  
LEU  
SER  
PRO  
GLY  
GLY  
LYS

● Molecule 3: HDIT102 Fab light chain

Chain L: 39%11%50%

GLN  
A2  
G3  
L4  
V12  
T17  
A18  
C22  
I27  
W34  
Q37  
I47  
Y48  
Y49  
P54  
I57  
P58  
E59  
R60  
S75  
R76  
A79  
G80  
D81  
E82  
V89  
V95  
V105  
L106  
GLY  
GLN  
PRO  
LYS  
LYS  
ALA  
ALA  
PRO  
SER  
VAL  
THR  
LEU  
PHE  
PRO  
PRO  
SER  
SER  
GLU

GLU  
LEU  
GLN  
ALA  
ASN  
LYS  
ALA  
THR  
THR  
LEU  
VAL  
CYS  
CYS  
LEU  
ILE  
SER  
ASP  
PHE  
TYR  
PRO  
GLY  
ALA  
VAL  
VAL  
THR  
LYS  
TRP  
ALA  
TRP  
LYS  
ASP  
ASP  
SER  
SER  
PRO  
VAL  
LYS  
GLN  
SER  
SER  
ASN  
LYS  
TYR  
ALA  
ALA  
PRO  
SER  
VAL  
THR  
LEU  
CYS  
GLU

GLN  
TRP  
LYS  
SER  
HIS  
ARG  
SER  
THR  
SER  
CYS  
GLN  
VAL  
THR  
HIS  
GLU  
GLY  
SER  
THR  
VAL  
GLU  
THR  
ALA  
PRO  
THR  
THR  
GLU  
CYS  
SER

● Molecule 3: HDIT102 Fab light chain

Chain E: 42%8%50%

GLN  
A2  
G3  
L4  
V12  
A18  
C22  
W34  
Y35  
I47  
R53  
P54  
I57  
P58  
E59  
R60  
D81  
C87  
Q88  
Q93  
L106  
GLY  
GLN  
PRO  
LYS  
ALA  
ALA  
PRO  
SER  
VAL  
THR  
LEU  
PHE  
PRO  
PRO  
SER  
SER  
GLU  
GLU  
LEU  
GLN  
ALA  
ASN  
LYS  
VAL  
CYS

LEU  
ILE  
SER  
ASP  
PHE  
TYR  
PRO  
GLY  
ALA  
VAL  
THR  
VAL  
ALA  
ALA  
TRP  
LYS  
ALA  
ALA  
SER  
SER  
PRO  
VAL  
LYS  
LYS  
ALA  
GLY  
GLU  
THR  
THR  
THR  
SER  
LYS  
GLN  
SER  
ASN  
LYS  
TYR  
ALA  
ALA  
SER  
SER  
TYR  
LEU  
SER  
SER  
LEU  
THR  
GLN  
TRP  
SER  
HIS  
ARG  
SER  
TYR  
THR  
CYS  
VAL  
CYS

VAL  
THR  
HIS  
GLU  
GLY  
SER  
THR  
VAL  
GLU  
LYS  
THR  
VAL  
VAL  
PRO  
THR  
GLU  
CYS  
SER

● Molecule 3: HDIT102 Fab light chain

Chain G: 41%8%50%

GLN  
A2  
G3  
L4  
V12  
T17  
A18  
W34  
K38  
I47  
Y48  
Y49  
D50  
P54  
I57  
P58  
E59  
S75  
R76  
A83  
V89  
V95  
L106  
GLY  
GLN  
PRO  
LYS  
LYS  
ALA  
ALA  
PRO  
SER  
VAL  
THR  
THR  
LEU  
PHE  
PRO  
PRO  
SER  
SER  
TYR  
LEU  
SER  
THR  
PRO  
GLU  
GLU  
GLN  
TRP  
LEU  
GLN  
ASN  
HIS  
SER  
ALA  
LYS  
ASN  
LYS  
THR

LEU  
VAL  
CYS  
LEU  
ILE  
SER  
ASP  
PHE  
TYR  
PRO  
GLY  
ALA  
VAL  
THR  
VAL  
VAL  
ALA  
ALA  
TRP  
TRP  
LYS  
ALA  
ASP  
SER  
SER  
PRO  
VAL  
LYS  
ALA  
GLY  
VAL  
GLU  
THR  
THR  
PRO  
SER  
LYS  
GLN  
SER  
ASN  
ASN  
LYS  
TYR  
ALA  
ALA  
SER  
SER  
SER  
SER  
TYR  
LEU  
SER  
LEU  
THR  
PRO  
GLU  
GLN  
TRP  
SER  
HIS  
SER  
ARG  
SER  
TYR

SER  
CYS  
GLN  
VAL  
THR  
HIS  
GLU  
GLY  
SER  
THR  
VAL  
GLU  
LYS  
THR  
VAL  
ALA  
ALA  
PRO  
TRP  
THR  
THR  
CYS  
SER



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	575340	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.256	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.159	Depositor
Minimum map value	-0.087	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0175	Depositor
Map size ( $\text{\AA}$ )	386.28, 386.28, 386.28	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.2876, 1.2876, 1.2876	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/4827	0.49	0/6556
1	B	0.29	0/4827	0.49	0/6556
1	C	0.30	0/4827	0.49	0/6556
2	D	0.29	0/920	0.51	0/1247
2	F	0.29	0/920	0.53	0/1247
2	H	0.29	0/920	0.54	0/1247
3	E	0.31	0/791	0.51	0/1078
3	G	0.31	0/791	0.51	0/1078
3	L	0.30	0/791	0.49	0/1078
All	All	0.30	0/19614	0.50	0/26643

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4710	0	4544	73	0
1	B	4710	0	4544	74	0
1	C	4710	0	4544	69	0
2	D	901	0	861	10	0
2	F	901	0	861	23	0
2	H	901	0	861	16	0
3	E	773	0	743	8	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	G	773	0	743	12	0
3	L	773	0	743	15	0
All	All	19152	0	18444	252	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 252 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:107:VAL:HG12	1:C:617:LEU:HB2	1.68	0.75
1:A:367:MET:HG2	1:A:380:SER:HB2	1.68	0.74
1:C:608:GLU:HB3	1:C:621:ARG:HH12	1.52	0.74
2:F:96:THR:HA	2:F:123:VAL:O	1.88	0.74
3:L:34:TRP:HB2	3:L:47:ILE:HB	1.70	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	579/703 (82%)	572 (99%)	7 (1%)	0	100	100
1	B	579/703 (82%)	573 (99%)	6 (1%)	0	100	100
1	C	579/703 (82%)	572 (99%)	7 (1%)	0	100	100
2	D	117/452 (26%)	113 (97%)	4 (3%)	0	100	100
2	F	117/452 (26%)	113 (97%)	4 (3%)	0	100	100
2	H	117/452 (26%)	114 (97%)	3 (3%)	0	100	100
3	E	103/212 (49%)	98 (95%)	5 (5%)	0	100	100
3	G	103/212 (49%)	97 (94%)	6 (6%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	L	103/212 (49%)	99 (96%)	4 (4%)	0	100	100
All	All	2397/4101 (58%)	2351 (98%)	46 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	507/593 (86%)	507 (100%)	0	100	100
1	B	507/593 (86%)	507 (100%)	0	100	100
1	C	507/593 (86%)	506 (100%)	1 (0%)	92	96
2	D	93/395 (24%)	93 (100%)	0	100	100
2	F	93/395 (24%)	93 (100%)	0	100	100
2	H	93/395 (24%)	92 (99%)	1 (1%)	70	83
3	E	84/176 (48%)	84 (100%)	0	100	100
3	G	84/176 (48%)	83 (99%)	1 (1%)	67	82
3	L	84/176 (48%)	83 (99%)	1 (1%)	67	82
All	All	2052/3492 (59%)	2048 (100%)	4 (0%)	91	96

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	169	HIS
2	H	77	ARG
3	L	76	ARG
3	G	76	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	335	ASN
1	C	393	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19166. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


### 6.2.2 Raw map



X Index: 150



Y Index: 150



Z Index: 150


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 147




Y Index: 145




Z Index: 159


### 6.3.2 Raw map



X Index: 147



Y Index: 145




Z Index: 159

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y



Z


### 6.4.2 Raw map



X



Y



Z


The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



X



Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.0175. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_19166\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 137 nm<sup>3</sup>; this corresponds to an approximate mass of 123 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.321 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.321  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.12	-	-
Author-provided FSC curve	3.10	3.37	3.12
Unmasked-calculated*	3.51	4.04	3.55


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.51 differs from the reported value 3.12 by more than 10 %



## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-19166 and PDB model 8RH2. Per-residue inclusion information can be found in section [3](#) on page [5](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0175 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0175).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0175) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8670	<div><div></div></div> 0.5620
A	<div><div></div></div> 0.8610	<div><div></div></div> 0.5570
B	<div><div></div></div> 0.8580	<div><div></div></div> 0.5600
C	<div><div></div></div> 0.8600	<div><div></div></div> 0.5620
D	<div><div></div></div> 0.9100	<div><div></div></div> 0.5780
E	<div><div></div></div> 0.8650	<div><div></div></div> 0.5610
F	<div><div></div></div> 0.8950	<div><div></div></div> 0.5710
G	<div><div></div></div> 0.8710	<div><div></div></div> 0.5660
H	<div><div></div></div> 0.9020	<div><div></div></div> 0.5640
L	<div><div></div></div> 0.8630	<div><div></div></div> 0.5670

1.0

0.0

<0.0