



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 22, 2025 – 05:40 PM EDT

PDB ID : 4RIT
Title : The yellow crystal structure of pyridoxal-dependent decarboxylase from
sphaerobacter thermophilus dsm 20745
Authors : Wu, R.; Clancy, S.; Joachimiak, A.; Midwest Center for Structural Genomics
(MCSG)
Deposited on : 2014-10-07
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	
1	B	486	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	506	-	-	X	-
4	TRS	B	508	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7997 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridoxal-dependent decarboxylase.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	B	469	Total	C	N	O	P	S	Se	0	13	0
			3695	2335	665	674	1	4	16			
1	A	461	Total	C	N	O	P	S	Se	0	14	0
			3651	2309	655	668	1	3	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	SER	-	expression tag	UNP D1C7D8
B	-1	ASN	-	expression tag	UNP D1C7D8
B	0	ALA	-	expression tag	UNP D1C7D8
A	-2	SER	-	expression tag	UNP D1C7D8
A	-1	ASN	-	expression tag	UNP D1C7D8
A	0	ALA	-	expression tag	UNP D1C7D8

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Cl	0	0
			2	2		
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			8	4	1	3		
4	B	1	Total	C	N	O	0	0
			8	4	1	3		
4	A	1	Total	C	N	O	0	0
			8	4	1	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	291	Total	O	0	0
			291	291		
5	A	267	Total	O	0	0
			267	267		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyridoxal-dependent decarboxylase



- Molecule 1: Pyridoxal-dependent decarboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.48Å 118.57Å 126.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.59 – 1.80 39.59 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (39.59-1.80) 99.3 (39.59-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.94 (at 1.79Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1496)	Depositor
R, R_{free}	0.150 , 0.173 0.158 , 0.176	Depositor DCC
R_{free} test set	5352 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7997	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, TRS, GOL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	2/3685 (0.1%)	0.72	3/4984 (0.1%)
1	B	0.76	6/3730 (0.2%)	0.75	5/5045 (0.1%)
All	All	0.73	8/7415 (0.1%)	0.74	8/10029 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	330	PRO	N-CD	5.54	1.55	1.47
1	B	93	PRO	N-CD	5.46	1.55	1.47
1	B	47	PRO	N-CD	5.40	1.55	1.47
1	A	93	PRO	N-CD	5.34	1.55	1.47
1	B	407[A]	CYS	CB-SG	-5.29	1.73	1.81

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	392	ARG	C-N-CD	5.55	140.05	128.40
1	B	92	ASP	C-N-CD	5.41	139.77	128.40
1	B	373	ARG	NE-CZ-NH1	-5.38	117.61	120.30
1	A	92	ASP	C-N-CD	5.34	139.62	128.40
1	B	46	MSE	C-N-CD	5.33	139.60	128.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3651	0	3642	32	0
1	B	3695	0	3679	29	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
3	A	36	0	48	6	0
3	B	30	0	40	4	0
4	A	8	0	12	1	0
4	B	16	0	24	1	0
5	A	267	0	0	4	0
5	B	291	0	0	3	0
All	All	7997	0	7445	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 56 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:457:PRO:HA	3:A:504:GOL:H31	1.57	0.84
1:B:409:ARG:HH22	3:B:506:GOL:H2	1.43	0.80
1:B:46:MSE:HE3	1:B:47:PRO:HD2	1.64	0.78
1:B:288[A]:ARG:NH1	5:B:829:HOH:O	2.21	0.73
1:A:131:ARG:HH22	3:A:503:GOL:H32	1.52	0.73

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/486 (96%)	455 (97%)	13 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	477/486 (98%)	463 (97%)	14 (3%)	0	100	100
All	All	945/972 (97%)	918 (97%)	27 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/362 (102%)	364 (99%)	5 (1%)	62	56
1	B	371/362 (102%)	369 (100%)	2 (0%)	86	86
All	All	740/724 (102%)	733 (99%)	7 (1%)	79	72

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	376[A]	ASP
1	A	376[B]	ASP
1	A	477	ARG
1	A	380	ARG
1	A	373	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	B	304	1	23,24,25	2.06	5 (21%)	25,32,34	1.71	5 (20%)
1	LLP	A	304	1	23,24,25	1.97	6 (26%)	25,32,34	1.71	6 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	B	304	1	-	4/16/17/19	0/1/1/1
1	LLP	A	304	1	-	4/16/17/19	0/1/1/1

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	304	LLP	C4'-NZ	6.44	1.48	1.27
1	A	304	LLP	C4'-NZ	6.42	1.48	1.27
1	B	304	LLP	C4-C4'	3.57	1.54	1.46
1	A	304	LLP	C2'-C2	2.99	1.55	1.50
1	A	304	LLP	C4-C4'	2.96	1.52	1.46

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	304	LLP	OP4-C5'-C5	4.09	117.02	109.36
1	B	304	LLP	OP4-C5'-C5	3.99	116.83	109.36
1	B	304	LLP	C5-C6-N1	-3.70	117.82	123.83
1	A	304	LLP	C5-C6-N1	-3.20	118.62	123.83
1	A	304	LLP	C4-C4'-NZ	-2.91	110.64	124.04

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	304	LLP	C4-C4'-NZ-CE
1	A	304	LLP	C4-C4'-NZ-CE
1	A	304	LLP	CG-CD-CE-NZ
1	B	304	LLP	CG-CD-CE-NZ
1	B	304	LLP	C3-C4-C4'-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 3 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TRS	A	508	-	7,7,7	0.55	0	9,9,9	1.55	1 (11%)
3	GOL	B	505	-	5,5,5	0.47	0	5,5,5	0.89	0
3	GOL	A	503	-	5,5,5	0.35	0	5,5,5	0.34	0
3	GOL	B	504	-	5,5,5	0.40	0	5,5,5	0.31	0
4	TRS	B	508	-	7,7,7	0.98	1 (14%)	9,9,9	2.92	4 (44%)
3	GOL	B	507	-	5,5,5	0.17	0	5,5,5	0.72	0
4	TRS	B	509	-	7,7,7	0.42	0	9,9,9	1.09	0
3	GOL	A	504	-	5,5,5	0.79	0	5,5,5	0.63	0
3	GOL	A	506	-	5,5,5	0.30	0	5,5,5	0.46	0
3	GOL	A	502	-	5,5,5	0.15	0	5,5,5	1.00	0
3	GOL	A	507	-	5,5,5	0.26	0	5,5,5	0.31	0
3	GOL	B	503	-	5,5,5	0.38	0	5,5,5	0.26	0
3	GOL	B	506	-	5,5,5	0.26	0	5,5,5	0.32	0
3	GOL	A	505	-	5,5,5	0.32	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TRS	A	508	-	-	3/9/9/9	-
3	GOL	B	505	-	-	2/4/4/4	-
3	GOL	A	503	-	-	4/4/4/4	-
3	GOL	B	504	-	-	1/4/4/4	-
4	TRS	B	508	-	-	5/9/9/9	-
3	GOL	B	507	-	-	2/4/4/4	-
4	TRS	B	509	-	-	6/9/9/9	-
3	GOL	A	504	-	-	4/4/4/4	-
3	GOL	A	506	-	-	3/4/4/4	-
3	GOL	A	502	-	-	2/4/4/4	-
3	GOL	A	507	-	-	2/4/4/4	-
3	GOL	B	503	-	-	4/4/4/4	-
3	GOL	B	506	-	-	4/4/4/4	-
3	GOL	A	505	-	-	3/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	508	TRS	C2-C	-2.14	1.47	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	508	TRS	O2-C2-C	-6.92	91.60	110.88
4	B	508	TRS	C2-C-N	-3.45	99.37	108.17
4	A	508	TRS	O3-C3-C	-3.02	102.45	110.88
4	B	508	TRS	O1-C1-C	2.88	118.90	110.88
4	B	508	TRS	C1-C-N	2.32	114.09	108.17

There are no chirality outliers.

5 of 45 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	503	GOL	O1-C1-C2-O2
3	B	503	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	503	GOL	C1-C2-C3-O3
3	B	507	GOL	O1-C1-C2-O2
3	B	507	GOL	O1-C1-C2-C3

There are no ring outliers.

7 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	508	TRS	1	0
3	A	503	GOL	2	0
4	B	508	TRS	1	0
3	A	504	GOL	1	0
3	A	506	GOL	1	0
3	B	506	GOL	4	0
3	A	505	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	448/486 (92%)	-0.25	15 (3%) 49 47	10, 26, 50, 71	11 (2%)
1	B	456/486 (93%)	-0.33	15 (3%) 49 47	10, 23, 49, 76	9 (1%)
All	All	904/972 (93%)	-0.29	30 (3%) 49 47	10, 24, 49, 76	20 (2%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	52	LEU	6.7
1	B	51	ASP	5.8
1	B	50	VAL	4.7
1	B	339	GLY	4.1
1	A	55	PRO	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	B	304	24/25	0.96	0.08	14,21,36,39	2
1	LLP	A	304	24/25	0.96	0.08	16,22,33,36	3

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	507	6/6	0.56	0.25	46,47,74,81	0
3	GOL	B	506	6/6	0.72	0.20	64,72,77,78	0
3	GOL	B	507	6/6	0.77	0.20	42,55,70,72	0
3	GOL	A	503	6/6	0.78	0.15	60,66,73,74	0
4	TRS	B	509	8/8	0.78	0.17	58,62,68,68	0
4	TRS	A	508	8/8	0.80	0.20	49,58,64,65	0
3	GOL	A	504	6/6	0.81	0.20	54,58,60,66	0
3	GOL	B	505	6/6	0.82	0.17	60,67,72,74	0
4	TRS	B	508	8/8	0.83	0.16	51,57,61,65	0
3	GOL	A	506	6/6	0.84	0.19	34,69,74,81	0
3	GOL	B	503	6/6	0.85	0.15	56,61,63,65	0
3	GOL	B	504	6/6	0.86	0.12	33,35,38,53	0
3	GOL	A	505	6/6	0.87	0.17	63,69,71,73	0
3	GOL	A	502	6/6	0.88	0.13	43,48,49,49	0
2	CL	A	501	1/1	0.94	0.14	45,45,45,45	0
2	CL	B	501	1/1	0.97	0.08	31,31,31,31	0
2	CL	B	502	1/1	0.99	0.10	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.