



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 13, 2024 – 02:03 AM EDT

PDB ID : 6RNT  
Title : CRYSTAL STRUCTURE OF RIBONUCLEASE T1 COMPLEXED WITH  
ADENOSINE 2'-MONOPHOSPHATE AT 1.8-ANGSTROMS RESOLUTION  
Authors : Ding, J.; Koellner, G.; Grunert, H.-P.; Saenger, W.  
Deposited on : 1991-08-20  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	104	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 915 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RIBONUCLEASE T1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	104	Total	C	N	O	S	0	4	0
			787	485	127	171	4			

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is ADENOSINE-2'-MONOPHOSPHATE (three-letter code: 2AM) (formula:  $C_{10}H_{14}N_5O_7P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	104	Total 104	O 104	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: RIBONUCLEASE T1

Chain A: 



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.93Å 46.57Å 41.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.80	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-1.80)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROFFT	Depositor
R, $R_{free}$	0.149 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	915	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	11.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, 2AM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	6.55	13/831 (1.6%)	6.31	91/1132 (8.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	49[A]	ASP	CG-OD1	107.66	3.73	1.25
1	A	49[B]	ASP	CG-OD1	107.66	3.73	1.25
1	A	49[A]	ASP	CG-OD2	71.31	2.89	1.25
1	A	49[B]	ASP	CG-OD2	71.31	2.89	1.25
1	A	11	TYR	CE1-CZ	10.29	1.51	1.38
1	A	49[A]	ASP	CB-CG	8.63	1.69	1.51
1	A	49[B]	ASP	CB-CG	8.63	1.69	1.51
1	A	11	TYR	CG-CD2	6.15	1.47	1.39
1	A	88	GLY	N-CA	6.11	1.55	1.46
1	A	24	TYR	CG-CD2	5.46	1.46	1.39
1	A	17	SER	CA-CB	5.27	1.60	1.52
1	A	70	GLY	N-CA	5.14	1.53	1.46
1	A	59	TRP	CD2-CE2	-5.06	1.35	1.41

All (91) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49[A]	ASP	CB-CG-OD1	-102.39	26.15	118.30
1	A	49[B]	ASP	CB-CG-OD1	-102.39	26.15	118.30
1	A	49[A]	ASP	CB-CG-OD2	-70.41	54.93	118.30
1	A	49[B]	ASP	CB-CG-OD2	-70.41	54.93	118.30
1	A	49[A]	ASP	OD1-CG-OD2	-45.83	36.23	123.30
1	A	49[B]	ASP	OD1-CG-OD2	-45.83	36.23	123.30
1	A	50	PHE	CB-CG-CD1	16.96	132.67	120.80
1	A	50	PHE	CB-CG-CD2	-16.33	109.37	120.80
1	A	11	TYR	CB-CG-CD2	-15.00	112.00	121.00
1	A	42	TYR	CB-CG-CD1	-13.28	113.03	121.00
1	A	42	TYR	CG-CD1-CE1	-12.55	111.26	121.30
1	A	78[A]	VAL	CA-CB-CG2	11.94	128.81	110.90
1	A	78[B]	VAL	CA-CB-CG2	11.94	128.81	110.90
1	A	11	TYR	CB-CG-CD1	11.20	127.72	121.00
1	A	46	GLU	OE1-CD-OE2	11.02	136.52	123.30
1	A	77	ARG	NE-CZ-NH2	10.71	125.66	120.30
1	A	49[A]	ASP	CA-CB-CG	-10.50	90.31	113.40
1	A	49[B]	ASP	CA-CB-CG	-10.50	90.31	113.40
1	A	11	TYR	O-C-N	10.42	139.37	122.70
1	A	42	TYR	CD1-CG-CD2	9.59	128.45	117.90
1	A	3	ASP	CB-CG-OD1	9.57	126.91	118.30
1	A	4	TYR	CB-CG-CD1	-9.53	115.28	121.00
1	A	11	TYR	CD1-CE1-CZ	-9.25	111.47	119.80
1	A	96	SER	N-CA-CB	9.15	124.22	110.50
1	A	11	TYR	CG-CD2-CE2	-9.03	114.07	121.30
1	A	32	THR	N-CA-CB	-8.53	94.09	110.30
1	A	24	TYR	CB-CG-CD2	-8.49	115.91	121.00
1	A	100	PHE	CB-CG-CD2	-8.48	114.87	120.80
1	A	29	ASP	CB-CG-OD2	-8.47	110.68	118.30
1	A	99	ASN	O-C-N	8.38	136.12	122.70
1	A	79	VAL	O-C-N	8.27	135.93	122.70
1	A	45	TYR	CB-CG-CD2	-7.98	116.21	121.00
1	A	4	TYR	CB-CG-CD2	7.91	125.74	121.00
1	A	1	ALA	CB-CA-C	7.19	120.88	110.10
1	A	83	ASN	CA-CB-CG	-7.08	97.82	113.40
1	A	11	TYR	CG-CD1-CE1	7.06	126.95	121.30
1	A	24	TYR	CG-CD2-CE2	-7.00	115.70	121.30
1	A	52	VAL	N-CA-CB	-6.92	96.27	111.50
1	A	89	VAL	O-C-N	6.83	133.63	122.70
1	A	52	VAL	CG1-CB-CG2	6.76	121.71	110.90
1	A	63	SER	O-C-N	6.57	133.21	122.70
1	A	75	ALA	CB-CA-C	-6.51	100.34	110.10
1	A	5	THR	O-C-N	6.49	133.08	122.70

Continued on next page...



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	TYR	CB-CG-CD1	-6.48	117.11	121.00
1	A	10	CYS	O-C-N	6.46	133.04	122.70
1	A	40	HIS	CB-CA-C	6.42	123.24	110.40
1	A	76	ASP	O-C-N	6.41	132.96	122.70
1	A	11	TYR	CZ-CE2-CD2	6.39	125.56	119.80
1	A	42	TYR	CG-CD2-CE2	-6.37	116.21	121.30
1	A	10	CYS	N-CA-CB	-6.35	99.17	110.60
1	A	46	GLU	CB-CA-C	-6.29	97.83	110.40
1	A	38	TYR	CB-CG-CD2	-6.27	117.24	121.00
1	A	95	ALA	N-CA-CB	6.26	118.86	110.10
1	A	75	ALA	N-CA-CB	-6.25	101.35	110.10
1	A	78[A]	VAL	CA-CB-CG1	-6.23	101.55	110.90
1	A	78[B]	VAL	CA-CB-CG1	-6.23	101.55	110.90
1	A	40	HIS	CE1-NE2-CD2	6.21	122.11	106.60
1	A	29	ASP	O-C-N	6.09	133.55	123.20
1	A	80	PHE	CB-CG-CD2	-6.09	116.54	120.80
1	A	33	VAL	O-C-N	6.05	133.48	123.20
1	A	18	THR	CA-CB-CG2	6.04	120.85	112.40
1	A	84	ASN	OD1-CG-ND2	-6.01	108.08	121.90
1	A	37	SER	CA-CB-OG	-5.99	95.03	111.20
1	A	11	TYR	CE1-CZ-OH	-5.90	104.17	120.10
1	A	24	TYR	CB-CG-CD1	5.89	124.54	121.00
1	A	31	GLU	OE1-CD-OE2	5.87	130.35	123.30
1	A	70	GLY	C-N-CA	-5.87	109.98	122.30
1	A	70	GLY	O-C-N	-5.82	113.30	123.20
1	A	17	SER	CA-CB-OG	-5.79	95.57	111.20
1	A	84	ASN	O-C-N	-5.74	113.52	122.70
1	A	33	VAL	C-N-CA	-5.72	110.28	122.30
1	A	31	GLU	O-C-N	5.70	131.82	122.70
1	A	77	ARG	CG-CD-NE	5.67	123.71	111.80
1	A	58	GLU	N-CA-CB	5.62	120.72	110.60
1	A	41	LYS	N-CA-CB	-5.60	100.52	110.60
1	A	100	PHE	CB-CG-CD1	5.59	124.71	120.80
1	A	24	TYR	CA-CB-CG	-5.50	102.94	113.40
1	A	100	PHE	CB-CA-C	-5.44	99.51	110.40
1	A	83	ASN	CB-CG-OD1	-5.39	110.82	121.60
1	A	32	THR	CA-CB-CG2	5.35	119.89	112.40
1	A	40	HIS	CG-CD2-NE2	-5.30	99.12	109.20
1	A	8	SER	CA-CB-OG	5.24	125.34	111.20
1	A	11	TYR	OH-CZ-CE2	5.23	134.21	120.10
1	A	82	GLU	OE1-CD-OE2	-5.22	117.04	123.30
1	A	5	THR	CA-C-O	-5.20	109.17	120.10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	ASP	CB-CG-OD2	5.17	122.95	118.30
1	A	57	TYR	CD1-CE1-CZ	-5.15	115.16	119.80
1	A	38	TYR	CG-CD2-CE2	-5.12	117.21	121.30
1	A	59	TRP	NE1-CE2-CZ2	-5.10	124.79	130.40
1	A	87	ALA	O-C-N	5.09	131.85	123.20
1	A	48	PHE	CB-CG-CD1	5.05	124.33	120.80

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	32	THR	CB

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	787	0	687	3	0
2	A	1	0	0	0	0
3	A	23	0	12	0	0
4	A	104	0	0	0	0
All	All	915	0	699	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:22:ALA:O	1:A:26[B]:LEU:HD12	2.00	0.61
1:A:54:SER:HB2	1:A:55:PRO:HA	1.85	0.57

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	106/104 (102%)	104 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/85 (105%)	85 (96%)	4 (4%)	23	11

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	THR
1	A	41	LYS
1	A	52	VAL
1	A	84	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
3	2AM	A	105	-	21,25,25	1.84	6 (28%)	28,38,38	2.23	12 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	2AM	A	105	-	-	3/7/27/27	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	105	2AM	P-O2'	3.60	1.65	1.59
3	A	105	2AM	O4'-C1'	3.20	1.45	1.40
3	A	105	2AM	C2-N3	3.11	1.36	1.32

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	105	2AM	C2-N1	2.87	1.39	1.33
3	A	105	2AM	P-O2P	-2.75	1.44	1.54
3	A	105	2AM	C6-N6	-2.20	1.26	1.34

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	105	2AM	C4'-O4'-C1'	-4.70	105.62	109.92
3	A	105	2AM	O4'-C1'-C2'	-3.85	100.05	106.61
3	A	105	2AM	C5-C6-N1	-3.69	111.44	120.23
3	A	105	2AM	C5-C6-N6	3.19	125.16	120.31
3	A	105	2AM	O2P-P-O1P	3.13	123.02	110.83
3	A	105	2AM	C3'-C2'-C1'	2.74	108.05	102.81
3	A	105	2AM	O4'-C4'-C3'	2.63	110.37	105.15
3	A	105	2AM	C2'-C3'-C4'	-2.61	96.37	101.99
3	A	105	2AM	C6-C5-C4	2.52	122.81	117.90
3	A	105	2AM	C2'-C1'-N9	2.49	118.09	112.56
3	A	105	2AM	N6-C6-N1	2.36	123.39	118.33
3	A	105	2AM	O3P-P-O1P	2.28	119.72	110.83

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	105	2AM	C2'-O2'-P-O3P
3	A	105	2AM	C2'-O2'-P-O1P
3	A	105	2AM	O4'-C4'-C5'-O5'

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.