



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 24, 2024 – 12:17 am GMT

PDB ID : 8RNF
Title : HLA-E*01:03 in complex with SARS-CoV-2 Omicron Nsp13 peptide, VIPLSAPTL
Authors : Sun, R.; Achour, A.; Sala, B.M.; Sandalova, T.
Deposited on : 2024-01-09
Resolution : 1.87 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>0%</div> <div>93%</div> <div>7%</div> </div>
1	C	274	<div> <div>6%</div> <div>94%</div> <div>5%</div> </div>
1	F	274	<div> <div>3%</div> <div>91%</div> <div>7%</div> </div>
2	B	100	<div> <div>3%</div> <div>93%</div> <div>7%</div> </div>
2	D	100	<div> <div>3%</div> <div>97%</div> <div>0%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	100	<div><div></div><div>3%</div><div>91%</div><div>9%</div></div>
3	E	9	<div><div></div><div>11%</div><div>100%</div><div></div></div>
3	H	9	<div><div></div><div></div><div>100%</div><div></div></div>
3	P	9	<div><div></div><div></div><div>100%</div><div></div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 10417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class I histocompatibility antigen, E alpha chain variant.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	274	Total	C	N	O	S	0	0	0
			2231	1394	399	431	7			
1	C	272	Total	C	N	O	S	0	0	0
			2218	1387	397	427	7			
1	F	269	Total	C	N	O	S	0	0	0
			2196	1375	392	422	7			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	D	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	G	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	initiating methionine	UNP P61769
D	1	MET	-	initiating methionine	UNP P61769
G	1	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called Non-structural protein 7.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	9	Total	C	N	O	0	0	0
			64	43	9	12			
3	H	9	Total	C	N	O	0	0	0
			64	43	9	12			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	0	0	0
			64	43	9	12			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	2	ILE	MET	variant	UNP P0DTD1
H	2	ILE	MET	variant	UNP P0DTD1
P	2	ILE	MET	variant	UNP P0DTD1

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	276	Total	O	0	0
			276	276		
4	B	139	Total	O	0	0
			139	139		
4	C	178	Total	O	0	0
			178	178		
4	D	91	Total	O	0	0
			91	91		
4	E	6	Total	O	0	0
			6	6		
4	F	253	Total	O	0	0
			253	253		
4	G	113	Total	O	0	0
			113	113		
4	H	8	Total	O	0	0
			8	8		
4	P	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class I histocompatibility antigen, E alpha chain variant





- Molecule 1: HLA class I histocompatibility antigen, E alpha chain variant



- Molecule 1: HLA class I histocompatibility antigen, E alpha chain variant




- Molecule 2: Beta-2-microglobulin




- Molecule 2: Beta-2-microglobulin



- Molecule 2: Beta-2-microglobulin



- Molecule 3: Non-structural protein 7



- Molecule 3: Non-structural protein 7



There are no outlier residues recorded for this chain.

- Molecule 3: Non-structural protein 7



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.23Å 66.53Å 158.45Å 90.00° 95.26° 90.00°	Depositor
Resolution (Å)	47.95 – 1.87 47.95 – 1.87	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.95-1.87) 99.7 (47.95-1.87)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 1.87Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.178 , 0.220 0.188 , 0.227	Depositor DCC
R_{free} test set	5263 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.050	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	10417	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2297	0.79	1/3123 (0.0%)
1	C	0.35	0/2283	0.72	0/3103
1	F	0.38	0/2260	0.80	1/3072 (0.0%)
2	B	0.39	0/860	0.78	1/1162 (0.1%)
2	D	0.36	0/860	0.72	0/1162
2	G	0.38	0/860	0.80	0/1162
3	E	0.48	0/65	0.63	0/89
3	H	0.48	0/65	0.75	0/89
3	P	0.48	0/65	0.74	0/89
All	All	0.38	0/9615	0.77	3/13051 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	100	MET	CG-SD-CE	5.79	109.46	100.20
1	A	44	ARG	CB-CA-C	-5.25	99.91	110.40
1	F	114	GLU	CG-CD-OE2	-5.12	108.06	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	65	ARG	Sidechain
1	C	44	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2231	0	2068	12	0
1	C	2218	0	2060	6	0
1	F	2196	0	2044	8	0
2	B	837	0	803	3	0
2	D	837	0	803	1	0
2	G	837	0	803	6	0
3	E	64	0	75	0	0
3	H	64	0	75	0	0
3	P	64	0	75	0	0
4	A	276	0	0	6	0
4	B	139	0	0	1	0
4	C	178	0	0	0	0
4	D	91	0	0	0	0
4	E	6	0	0	0	0
4	F	253	0	0	3	0
4	G	113	0	0	3	0
4	H	8	0	0	0	0
4	P	5	0	0	0	0
All	All	10417	0	8806	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 33 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:7:LYS:HD2	4:G:288:HOH:O	1.86	0.74
1:A:19:GLU:HG3	1:A:20:PRO:HD2	1.81	0.62
1:C:21:ARG:CD	1:C:39:ASP:HB2	2.36	0.56
1:A:136:VAL:HG21	4:A:465:HOH:O	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:THR:HG23	4:A:490:HOH:O	2.10	0.51

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/274 (99%)	267 (98%)	5 (2%)	0	100	100
1	C	268/274 (98%)	259 (97%)	9 (3%)	0	100	100
1	F	265/274 (97%)	261 (98%)	4 (2%)	0	100	100
2	B	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
2	D	98/100 (98%)	97 (99%)	1 (1%)	0	100	100
2	G	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
3	E	7/9 (78%)	7 (100%)	0	0	100	100
3	H	7/9 (78%)	7 (100%)	0	0	100	100
3	P	7/9 (78%)	7 (100%)	0	0	100	100
All	All	1120/1149 (98%)	1097 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/235 (100%)	232 (99%)	3 (1%)	65	55
1	C	234/235 (100%)	231 (99%)	3 (1%)	65	55
1	F	232/235 (99%)	225 (97%)	7 (3%)	36	19
2	B	95/95 (100%)	93 (98%)	2 (2%)	48	34
2	D	95/95 (100%)	93 (98%)	2 (2%)	48	34
2	G	95/95 (100%)	94 (99%)	1 (1%)	70	62
3	E	8/8 (100%)	8 (100%)	0	100	100
3	H	8/8 (100%)	8 (100%)	0	100	100
3	P	8/8 (100%)	8 (100%)	0	100	100
All	All	1010/1014 (100%)	992 (98%)	18 (2%)	54	40

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	162	ASP
2	G	71	PHE
1	F	272	LEU
2	D	71	PHE
1	F	82	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
2	G	84	ASN
2	G	32	HIS
1	F	96	GLN
2	D	3	GLN
1	F	156	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/274 (100%)	-0.08	3 (1%) 77 82	13, 21, 39, 54	0
1	C	272/274 (99%)	0.37	17 (6%) 27 28	15, 29, 55, 84	0
1	F	269/274 (98%)	-0.04	8 (2%) 52 56	12, 21, 43, 64	0
2	B	100/100 (100%)	-0.15	3 (3%) 52 56	14, 21, 36, 82	0
2	D	100/100 (100%)	0.18	3 (3%) 52 56	18, 27, 46, 62	0
2	G	100/100 (100%)	-0.03	3 (3%) 52 56	14, 21, 42, 67	0
3	E	9/9 (100%)	0.58	1 (11%) 12 13	27, 28, 38, 50	0
3	H	9/9 (100%)	-0.28	0 100 100	18, 19, 21, 22	0
3	P	9/9 (100%)	-0.34	0 100 100	16, 18, 21, 23	0
All	All	1142/1149 (99%)	0.06	38 (3%) 49 52	12, 23, 46, 84	0

The worst 5 of 38 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	223	GLY	4.7
2	G	1	MET	4.2
2	B	1	MET	4.1
1	F	40	ALA	3.5
1	F	41	ALA	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.