



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 09:12 AM EDT

PDB ID : 3RSR
Title : Crystal Structure of 5-NITP Inhibition of Yeast Ribonucleotide Reductase
Authors : Wan, Q.; Mohammed, F.; Jha, S.; Motea, E.; Berdis, A.; Dealwis, C.G.
Deposited on : 2011-05-02
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	888	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase large chain 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	610	Total	C	N	O	S	0	0	0
			4794	3066	800	900	28			

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 1-{2-DEOXY-5-O-[(R)-HYDROXY{[(R)-HYDROXY(PHOSPHONOOXY)P HOSPHORYL]OXY}PHOSPHORYL]-BETA-D-ERYTHRO-PENTOFURANOSYL}-5-NITRO -1H-INDOLE (three-letter code: N5P) (formula: C₁₃H₁₇N₂O₁₄P₃).



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	243	Total 243	O 243	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- [illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	108.35Å 118.21Å 63.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.97 – 2.30 19.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.97-2.30) 99.8 (19.97-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 2.19Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7_650)	Depositor
R, R_{free}	0.188 , 0.233 0.185 , 0.230	Depositor DCC
R_{free} test set	1861 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	36.7	Xtriage
Anisotropy	0.627	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5070	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, N5P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/4903	0.58	0/6653

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4794	0	4643	102	0
2	A	1	0	0	0	0
3	A	32	0	13	1	0
4	A	243	0	0	7	0
All	All	5070	0	4656	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:214:GLN:HG2	1:A:244:THR:HB	1.51	0.93
1:A:95:SER:H	1:A:132:ASN:HD21	1.16	0.92
1:A:582:PRO:O	1:A:596:PHE:HZ	1.59	0.86
1:A:582:PRO:HB2	1:A:596:PHE:CZ	2.11	0.85
1:A:595:CYS:SG	1:A:596:PHE:HA	2.24	0.78
1:A:596:PHE:N	1:A:596:PHE:CD1	2.50	0.77
1:A:595:CYS:HB3	1:A:596:PHE:HD1	1.51	0.75
1:A:669:ARG:O	1:A:698:GLN:HB2	1.89	0.72
1:A:596:PHE:HE2	1:A:662:HIS:HB2	1.54	0.72
1:A:398:ASN:HD22	1:A:398:ASN:H	1.37	0.72
1:A:487:GLN:HE22	1:A:590:LEU:HB2	1.57	0.70
1:A:423:CYS:HB3	1:A:425:LEU:HD21	1.73	0.70
1:A:614:TYR:CZ	1:A:618:GLN:HG3	2.27	0.69
1:A:90:THR:HG23	1:A:166:ARG:HE	1.57	0.69
1:A:408:CYS:SG	1:A:584:PRO:HG2	2.35	0.68
1:A:596:PHE:CE2	1:A:662:HIS:HB2	2.29	0.67
1:A:483:ALA:HB1	1:A:581:ALA:HB3	1.76	0.67
1:A:213:PRO:HD2	1:A:466:TYR:HB2	1.77	0.66
1:A:95:SER:H	1:A:132:ASN:ND2	1.91	0.65
1:A:373:LEU:HG	1:A:676:LEU:HD13	1.77	0.65
1:A:665:ASN:ND2	1:A:694:TYR:HB2	2.12	0.64
1:A:200:HIS:HE1	1:A:457:ASN:HD22	1.46	0.64
1:A:314:ILE:HD12	1:A:384:VAL:HG13	1.80	0.63
1:A:582:PRO:C	1:A:596:PHE:HZ	2.01	0.63
1:A:507:LEU:O	1:A:511:GLN:HG3	1.99	0.62
3:A:841:N5P:O1G	3:A:841:N5P:O1A	2.18	0.61
1:A:582:PRO:C	1:A:596:PHE:CZ	2.75	0.60
1:A:625:LEU:HD22	1:A:628:VAL:HG23	1.82	0.60
1:A:487:GLN:HA	1:A:595:CYS:HB2	1.83	0.60
1:A:566:ARG:O	1:A:570:MET:HG3	2.02	0.60
1:A:290:TYR:OH	1:A:405:SER:HB3	2.02	0.59
1:A:200:HIS:HD2	4:A:889:HOH:O	1.85	0.59
1:A:582:PRO:HB2	1:A:596:PHE:CE2	2.38	0.58
1:A:214:GLN:HG2	1:A:244:THR:CB	2.32	0.56
1:A:140:ASP:OD1	1:A:168:GLN:HG2	2.05	0.56
1:A:130:LYS:HB2	4:A:895:HOH:O	2.04	0.56
1:A:535:GLU:HG3	4:A:872:HOH:O	2.05	0.56
1:A:537:PHE:CZ	1:A:573:GLY:HA2	2.40	0.55
1:A:669:ARG:O	1:A:671:PRO:HD3	2.06	0.55
1:A:126:GLU:OE1	1:A:181:ARG:NH1	2.40	0.55
1:A:583:MET:O	1:A:596:PHE:HE1	1.90	0.55
1:A:115:MET:HE3	1:A:159:ILE:HG12	1.88	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:MET:CE	1:A:159:ILE:HG12	2.37	0.54
1:A:292:GLU:CD	1:A:387:LYS:HE2	2.28	0.54
1:A:407:LEU:HD23	1:A:407:LEU:O	2.08	0.54
1:A:672:THR:HG23	1:A:675:LYS:CB	2.38	0.54
1:A:662:HIS:HD2	4:A:865:HOH:O	1.92	0.53
1:A:289:LEU:HD11	1:A:308:LEU:HD23	1.91	0.52
1:A:582:PRO:O	1:A:596:PHE:CZ	2.50	0.52
1:A:95:SER:N	1:A:132:ASN:HD21	1.97	0.52
1:A:666:LEU:HD22	1:A:692:MET:HG3	1.92	0.51
1:A:595:CYS:CB	1:A:596:PHE:HA	2.41	0.51
1:A:94:PHE:HB3	1:A:132:ASN:HD22	1.76	0.50
1:A:101:LEU:HD23	1:A:115:MET:HE3	1.93	0.50
1:A:288:ALA:HB2	1:A:407:LEU:HA	1.94	0.50
1:A:582:PRO:CB	1:A:596:PHE:CZ	2.92	0.50
1:A:461:VAL:O	1:A:465:ASN:HB2	2.12	0.49
1:A:139:ARG:HH12	1:A:191:ASN:HD22	1.59	0.49
1:A:270:ASN:HB2	4:A:898:HOH:O	2.13	0.49
1:A:381:THR:HB	1:A:382:PRO:HA	1.95	0.49
1:A:617:THR:HG22	1:A:618:GLN:HG2	1.94	0.49
1:A:645:LYS:HG3	1:A:682:TYR:CE2	2.47	0.48
1:A:416:ALA:HB1	1:A:417:PRO:HD2	1.96	0.48
1:A:583:MET:O	1:A:596:PHE:CE1	2.68	0.47
1:A:256:ARG:HA	1:A:333:PRO:HD2	1.98	0.46
1:A:682:TYR:O	1:A:686:LYS:HG2	2.14	0.46
1:A:280:ASN:OD1	1:A:308:LEU:HD13	2.16	0.46
1:A:128:LYS:HE2	1:A:128:LYS:HB2	1.64	0.46
1:A:307:ASP:OD2	1:A:307:ASP:N	2.49	0.46
1:A:491:ASP:OD1	1:A:593:ASN:HB2	2.16	0.45
1:A:602:LEU:HD13	1:A:636:TYR:CD1	2.51	0.45
1:A:156:LEU:HG	1:A:165:GLU:O	2.16	0.45
1:A:665:ASN:HD22	1:A:694:TYR:HB2	1.81	0.45
1:A:273:ILE:HD12	1:A:304:ILE:HD11	1.98	0.45
1:A:638:THR:O	1:A:641:GLU:N	2.46	0.45
1:A:668:LEU:HD13	1:A:671:PRO:HA	1.98	0.45
1:A:304:ILE:HA	1:A:304:ILE:HD13	1.57	0.43
1:A:336:ALA:HB1	1:A:354:TYR:CD2	2.52	0.43
1:A:614:TYR:CE2	1:A:618:GLN:HG3	2.52	0.43
1:A:398:ASN:HD22	1:A:398:ASN:N	2.09	0.43
1:A:452:LYS:O	1:A:456:ARG:HG3	2.18	0.43
1:A:541:PRO:HD3	4:A:880:HOH:O	2.17	0.43
1:A:487:GLN:NE2	1:A:590:LEU:HB2	2.30	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:596:PHE:HE2	1:A:662:HIS:HD1	1.66	0.43
1:A:407:LEU:HD23	1:A:407:LEU:C	2.38	0.42
1:A:478:ARG:NH1	4:A:1080:HOH:O	2.53	0.42
1:A:583:MET:CB	1:A:584:PRO:CD	2.96	0.42
1:A:220:LEU:HB2	1:A:421:ALA:HB3	2.02	0.42
1:A:408:CYS:HB3	1:A:584:PRO:CG	2.50	0.42
1:A:425:LEU:HD22	1:A:483:ALA:HB3	2.01	0.42
1:A:151:LEU:HD13	1:A:589:ILE:HD13	2.02	0.42
1:A:344:GLY:O	1:A:348:GLU:HG3	2.19	0.42
1:A:619:ASN:OD1	1:A:619:ASN:O	2.37	0.42
1:A:622:ILE:HD11	1:A:636:TYR:HB2	2.01	0.42
1:A:292:GLU:OE1	1:A:387:LYS:HE2	2.19	0.41
1:A:697:THR:HG23	1:A:698:GLN:N	2.34	0.41
1:A:166:ARG:HD2	1:A:169:HIS:CE1	2.56	0.41
1:A:223:MET:HG2	1:A:255:ILE:HD11	2.03	0.41
1:A:628:VAL:HA	1:A:629:PRO:HD3	1.86	0.41
1:A:692:MET:SD	1:A:695:LEU:HB2	2.61	0.41
1:A:210:THR:HB	1:A:211:PRO:CD	2.51	0.41
1:A:139:ARG:O	1:A:142:GLN:HG2	2.21	0.40
1:A:416:ALA:HB1	1:A:417:PRO:CD	2.50	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	596/888 (67%)	584 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	509/761 (67%)	488 (96%)	21 (4%)	26	39

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	156	LEU
1	A	287	PHE
1	A	304	ILE
1	A	341	ASP
1	A	387	LYS
1	A	398	ASN
1	A	425	LEU
1	A	436	THR
1	A	491	ASP
1	A	495	LEU
1	A	565	LEU
1	A	595	CYS
1	A	596	PHE
1	A	638	THR
1	A	664	LEU
1	A	666	LEU
1	A	672	THR
1	A	673	MET
1	A	676	LEU
1	A	695	LEU
1	A	697	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	106	ASN
1	A	132	ASN
1	A	160	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	191	ASN
1	A	200	HIS
1	A	398	ASN
1	A	487	GLN
1	A	554	GLN
1	A	618	GLN
1	A	665	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	N5P	A	841	2	32,34,34	2.42	6 (18%)	43,53,53	1.87	10 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	N5P	A	841	2	-	0/20/38/38	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	841	N5P	OR2-NRO	9.75	1.39	1.22
3	A	841	N5P	PB-O3B	7.39	1.67	1.59
3	A	841	N5P	CZ3-NRO	-2.53	1.39	1.45
3	A	841	N5P	CD2-CE2	2.42	1.46	1.41
3	A	841	N5P	PA-O3A	2.20	1.61	1.59
3	A	841	N5P	PB-O3A	2.20	1.61	1.59

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	841	N5P	O2G-PG-O3B	-4.58	89.28	104.64
3	A	841	N5P	O2G-PG-O3G	4.52	124.75	107.80
3	A	841	N5P	CG-CD2-CE2	4.43	109.99	106.20
3	A	841	N5P	CH2-CZ3-NRO	-3.63	116.17	119.34
3	A	841	N5P	O3G-PG-O3B	3.59	116.68	104.64
3	A	841	N5P	O3G-PG-O1G	-3.59	96.85	110.83
3	A	841	N5P	O3B-PB-O1B	2.87	119.33	110.70
3	A	841	N5P	O2G-PG-O1G	2.54	120.73	110.83
3	A	841	N5P	CZ2-CE2-NE1	2.23	134.00	132.11
3	A	841	N5P	CE3-CZ3-NRO	2.05	120.50	118.74

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	841	N5P	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	610/888 (68%)	-0.16	20 (3%)	49 51	26, 40, 73, 91	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	595	CYS	4.8
1	A	583	MET	4.4
1	A	613	GLN	4.3
1	A	587	SER	4.2
1	A	694	TYR	4.0
1	A	584	PRO	3.8
1	A	594	GLU	3.5
1	A	612	TRP	3.5
1	A	592	TYR	3.4
1	A	437	SER	2.9
1	A	616	ILE	2.6
1	A	438	LYS	2.6
1	A	596	PHE	2.5
1	A	590	LEU	2.5
1	A	611	ILE	2.4
1	A	370	TYR	2.4
1	A	589	ILE	2.3
1	A	620	GLY	2.3
1	A	673	MET	2.2
1	A	593	ASN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	A	2001	1/1	0.77	0.18	54,54,54,54	0
3	N5P	A	841	32/32	0.87	0.11	29,35,46,74	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around N5P A 841:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.