



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 29, 2024 – 07:30 AM EDT

PDB ID : 3RY3
Title : Putative solute-binding protein from Yersinia pestis.
Authors : Osipiuk, J.; Zhou, M.; Kwon, K.; Anderson, W.F.; Joachimiak, A.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2011-05-10
Resolution : 2.43 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.43 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2124 (2.46-2.42)
Clashscore	180529	2259 (2.46-2.42)
Ramachandran outliers	177936	2244 (2.46-2.42)
Sidechain outliers	177891	2244 (2.46-2.42)
RSRZ outliers	164620	2124 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	528	
1	B	528	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7888 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative solute-binding protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	Se	0	4	0
			3831	2440	656	724	1	10			
1	B	490	Total	C	N	O	S	Se	0	5	0
			3849	2453	660	725	1	10			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q7CJS8
A	-1	ASN	-	expression tag	UNP Q7CJS8
A	0	ALA	-	expression tag	UNP Q7CJS8
B	-2	SER	-	expression tag	UNP Q7CJS8
B	-1	ASN	-	expression tag	UNP Q7CJS8
B	0	ALA	-	expression tag	UNP Q7CJS8

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	97	Total	O	0	0
			97	97		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Putative solute-binding protein



• Molecule 1: Putative solute-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	157.04Å 171.32Å 128.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.10 – 2.43 43.10 – 2.43	Depositor EDS
% Data completeness (in resolution range)	97.9 (43.10-2.43) 98.3 (43.10-2.43)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.31 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.5.0109, PHENIX	Depositor
R, R_{free}	0.173 , 0.208 0.174 , 0.205	Depositor DCC
R_{free} test set	3276 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.540	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7888	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	1/3933 (0.0%)	0.82	2/5343 (0.0%)
1	B	0.93	1/3958 (0.0%)	0.86	6/5378 (0.1%)
All	All	0.94	2/7891 (0.0%)	0.84	8/10721 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	396	GLU	CG-CD	5.66	1.60	1.51
1	A	70	TRP	CB-CG	-5.20	1.40	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	285	ARG	NE-CZ-NH2	-9.32	115.64	120.30
1	A	285	ARG	NE-CZ-NH2	-7.08	116.76	120.30
1	B	285	ARG	NE-CZ-NH1	6.97	123.78	120.30
1	B	134	ASP	CB-CG-OD1	6.51	124.16	118.30
1	B	327	ASP	CB-CG-OD1	6.00	123.70	118.30
1	A	139	ARG	NE-CZ-NH2	-5.96	117.32	120.30
1	B	499	ASP	CB-CG-OD1	5.63	123.36	118.30
1	B	190	MSE	CG-SE-CE	-5.01	87.87	98.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3831	0	3720	23	0
1	B	3849	0	3748	32	0
2	B	8	0	6	0	0
3	A	103	0	0	1	0
3	B	97	0	0	1	0
All	All	7888	0	7474	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:169:TYR:OH	1:B:174:ILE:O	2.00	0.79
1:A:46:MSE:HE1	1:A:111:THR:HG23	1.65	0.77
1:A:375:GLN:HG3	1:B:375:GLN:HG3	1.66	0.75
1:A:330:ILE:HG22	1:A:334:LYS:HE2	1.79	0.63
1:A:133:ILE:HD11	1:A:139:ARG:HB2	1.82	0.61
1:A:32:GLN:HG2	1:A:209:ILE:HB	1.83	0.60
1:B:134:ASP:HB2	1:B:135:PRO:CD	2.33	0.58
1:B:32:GLN:HG2	1:B:209:ILE:HB	1.85	0.58
1:A:38:GLU:OE1	1:A:187:GLY:N	2.35	0.55
1:A:425:SER:HB2	1:A:439:LYS:HG2	1.89	0.54
1:B:117:ALA:O	1:B:120:GLY:N	2.42	0.53
1:B:258:VAL:HG13	1:B:258:VAL:O	2.09	0.53
1:B:45:PRO:C	1:B:47:LEU:H	2.11	0.52
1:B:49:TRP:CD2	1:B:155:SER:HB3	2.44	0.52
1:B:301:ILE:HD12	1:B:372:ASP:HB3	1.93	0.50
1:B:86:THR:HG23	3:B:576:HOH:O	2.12	0.49
1:B:272:TYR:CE1	1:B:400:ARG:HG2	2.48	0.49
1:A:106:LYS:HE3	3:A:627:HOH:O	2.11	0.49
1:B:506:GLU:HB3	1:B:512:SER:HB2	1.94	0.49
1:B:257:ILE:HD11	1:B:377:LEU:HD13	1.94	0.48
1:B:302:MSE:HE1	1:B:369:THR:CG2	2.43	0.48
1:B:417:MSE:HE3	1:B:420:TYR:CD2	2.48	0.48
1:A:65:ASN:OD1	1:A:71:ASP:OD2	2.32	0.48
1:B:228:GLY:O	1:B:494:ALA:HA	2.15	0.47
1:B:364:THR:HG21	1:B:395:TRP:CE2	2.51	0.46
1:B:446:HIS:HE1	1:B:469:ASP:OD2	1.99	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:ASN:ND2	1:A:71:ASP:OD2	2.47	0.46
1:B:382:LYS:HB2	1:B:383:PRO:HD3	1.98	0.45
1:B:363:TYR:O	1:B:392:SER:HA	2.16	0.45
1:B:67:ASP:O	1:B:68:PHE:HB2	2.16	0.45
1:A:434:ASN:N	1:A:435:PRO:CD	2.80	0.45
1:A:363:TYR:O	1:A:392:SER:HA	2.17	0.44
1:A:285:ARG:HD2	1:A:480:ALA:O	2.17	0.44
1:A:425:SER:CB	1:A:439:LYS:HG2	2.48	0.44
1:A:172:LYS:HB3	1:A:173:PRO:HD2	2.00	0.43
1:A:30:THR:O	1:A:30:THR:HG22	2.19	0.43
1:B:39:PRO:HA	1:B:51:HIS:HE1	1.83	0.43
1:B:443:VAL:HG13	1:B:466:VAL:HG13	2.01	0.42
1:A:134:ASP:HB2	1:A:135:PRO:CD	2.49	0.42
1:A:434:ASN:N	1:A:435:PRO:HD3	2.34	0.42
1:B:330:ILE:HG22	1:B:334:LYS:HE2	2.01	0.42
1:A:30:THR:HG23	1:A:31:LEU:N	2.34	0.42
1:A:128:LEU:O	1:A:129:SER:HB3	2.19	0.42
1:B:72:SER:HB2	1:B:75:LEU:O	2.20	0.41
1:B:66:GLU:OE2	1:B:502:LYS:HE3	2.21	0.41
1:A:67:ASP:O	1:A:68:PHE:HB2	2.20	0.41
1:A:172:LYS:HB3	1:A:173:PRO:CD	2.50	0.41
1:B:167:LYS:HB3	1:B:168:THR:HG23	2.03	0.41
1:B:302:MSE:HE1	1:B:369:THR:HG21	2.03	0.41
1:B:517:ILE:HA	1:B:520:TRP:CE2	2.56	0.41
1:A:63:LYS:NZ	1:A:518:ASP:OD2	2.54	0.40
1:B:296:LEU:HD11	1:B:300[B]:GLN:NE2	2.36	0.40
1:B:86:THR:HA	1:B:140:ILE:O	2.21	0.40
1:B:66:GLU:OE2	1:B:502:LYS:CE	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	489/528 (93%)	470 (96%)	19 (4%)	0	100	100
1	B	491/528 (93%)	465 (95%)	25 (5%)	1 (0%)	44	53
All	All	980/1056 (93%)	935 (95%)	44 (4%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	46	MSE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	405/424 (96%)	395 (98%)	10 (2%)	42	55
1	B	408/424 (96%)	396 (97%)	12 (3%)	37	49
All	All	813/848 (96%)	791 (97%)	22 (3%)	40	52

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	THR
1	A	73	LEU
1	A	77	GLN
1	A	86	THR
1	A	106	LYS
1	A	240	SER
1	A	245	LYS
1	A	300	GLN
1	A	379	SER
1	A	389	ASP
1	B	40	THR
1	B	50	SER
1	B	63	LYS
1	B	72	SER
1	B	114	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	171	GLN
1	B	202	LYS
1	B	217	SER
1	B	343	GLN
1	B	352	LYS
1	B	419	LEU
1	B	435	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	137	ASN
1	A	243	ASN
1	A	319	ASN
1	A	339	GLN
1	B	51	HIS
1	B	137	ASN
1	B	319	ASN
1	B	446	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	B	701	-	3,3,3	0.98	0	3,3,3	0.41	0
2	ACT	B	702	-	3,3,3	0.86	0	3,3,3	1.13	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	479/528 (90%)	-0.42	4 (0%) 82 83	20, 41, 64, 83	4 (0%)
1	B	480/528 (90%)	-0.38	14 (2%) 54 55	21, 41, 71, 90	5 (1%)
All	All	959/1056 (90%)	-0.40	18 (1%) 66 67	20, 41, 68, 90	9 (0%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	40	THR	5.1
1	B	120	GLY	4.6
1	A	38	GLU	4.0
1	A	42	GLY	3.3
1	B	41	GLU	3.1
1	B	121	LYS	2.9
1	B	170	ALA	2.7
1	B	43	PHE	2.6
1	B	125	GLY	2.6
1	B	167	LYS	2.5
1	B	42	GLY	2.4
1	B	47	LEU	2.4
1	B	206	ASP	2.3
1	B	215	GLU	2.2
1	A	119	GLY	2.2
1	B	37	ASP	2.2
1	A	120	GLY	2.1
1	B	30	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	B	701	4/4	0.96	0.09	34,35,35,35	0
2	ACT	B	702	4/4	0.96	0.10	45,45,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.