



wwPDB EM Validation Summary Report ⓘ

Dec 14, 2024 – 10:01 am GMT

PDB ID : 8S0E
EMDB ID : EMD-19623
Title : H. sapiens OCCM bound to double stranded DNA
Authors : Greiwe, J.F.; Weissmann, F.; Diffley, J.F.X.; Costa, A.
Deposited on : 2024-02-13
Resolution : 3.80 Å(reported)
Based on initial models : 7JPS, 7JPR, .

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	39	<div> <div>13%</div> <div>87%</div> </div>
2	Y	39	<div> <div>18%</div> <div>82%</div> </div>
3	5	734	<div> <div>27%</div> <div>35%</div> <div>62%</div> </div>
4	8	546	<div> <div>23%</div> <div>37%</div> <div>10%</div> <div>53%</div> </div>
5	2	904	<div> <div>49%</div> <div>14%</div> <div>37%</div> </div>
6	4	863	<div> <div>46%</div> <div>13%</div> <div>41%</div> </div>
7	6	821	<div> <div>63%</div> <div>14%</div> <div>23%</div> </div>
8	7	719	<div> <div>16%</div> <div>40%</div> <div>10%</div> <div>50%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	B	577	
10	C	711	
11	G	560	
12	3	810	
13	A	861	
14	D	436	
15	E	435	

2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 44676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (39-mer).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	X	39	Total	C	N	O	P	0	0
			799	379	146	235	39		

- Molecule 2 is a DNA chain called DNA (39-mer).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	Y	39	Total	C	N	O	P	0	0
			800	379	149	233	39		

- Molecule 3 is a protein called DNA replication licensing factor MCM5.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	5	279	Total	C	N	O	S	0	0
			2169	1364	383	408	14		

- Molecule 4 is a protein called DNA replication factor Cdt1.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	8	259	Total	C	N	O	S	0	0
			2097	1306	390	386	15		

- Molecule 5 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	2	572	Total	C	N	O	S	0	0
			4513	2838	809	843	23		

- Molecule 6 is a protein called DNA replication licensing factor MCM4.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	4	512	Total	C	N	O	S	0	0
			4110	2598	728	760	24		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
4	650	MET	LEU	variant	UNP P33991

- Molecule 7 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	6	629	Total	C	N	O	S	0	0
			5091	3223	888	955	25		

- Molecule 8 is a protein called DNA replication licensing factor MCM7.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	7	357	Total	C	N	O	S	0	0
			2841	1774	515	534	18		

- Molecule 9 is a protein called Origin recognition complex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	B	273	Total	C	N	O	S	0	0
			2247	1446	379	416	6		

- Molecule 10 is a protein called Origin recognition complex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	C	574	Total	C	N	O	S	0	0
			4689	3027	781	854	27		

- Molecule 11 is a protein called Cell division control protein 6 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	G	337	Total	C	N	O	S	0	0
			2677	1701	473	483	20		

- Molecule 12 is a protein called DNA replication licensing factor MCM3.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	3	421	Total	C	N	O	S	0	0
			3281	2046	588	627	20		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
3	-1	GLY	-	expression tag	UNP P25205
3	0	GLU	-	expression tag	UNP P25205

- Molecule 13 is a protein called Origin recognition complex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	A	364	Total	C	N	O	S	0	0
			2889	1817	509	540	23		


- Molecule 14 is a protein called Origin recognition complex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	D	406	Total	C	N	O	S	0	0
			3309	2113	575	601	20		

- Molecule 15 is a protein called Origin recognition complex subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	E	364	Total	C	N	O	S	0	0
			3003	1967	497	529	10		

- Molecule 16 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (three-letter code: AGS) (formula: $C_{10}H_{16}N_5O_{12}P_3S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
16	2	1	Total	C	N	O	P	S
			31	10	5	12	3	1

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						AltConf
16	6	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	
16	A	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	
16	D	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	
16	E	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	

- Molecule 17 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		AltConf
17	2	1	Total	Mg	0
			1	1	
17	6	1	Total	Mg	0
			1	1	
17	A	1	Total	Mg	0
			1	1	
17	D	1	Total	Mg	0
			1	1	
17	E	1	Total	Mg	0
			1	1	


- Molecule 18 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
18	4	1	Total	Zn	0
			1	1	




Chain 2:






- Molecule 7: DNA replication licensing factor MCM6




- Molecule 8: DNA replication licensing factor MCM7






• Molecule 10: Origin recognition complex subunit 3

Chain C: 63% 18% 19%





• Molecule 11: Cell division control protein 6 homolog

Chain G: 18% 52% 8% 40%





● Molecule 15: Origin recognition complex subunit 5



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	34116	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	49.28	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	130000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.830	Depositor
Minimum map value	-0.311	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.027	Depositor
Recommended contour level	0.145	Depositor
Map size (Å)	432.00003, 432.00003, 432.00003	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.08, 1.08, 1.08	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, AGS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.56	0/895	0.91	0/1379
2	Y	0.55	0/897	0.88	0/1382
3	5	0.25	0/2198	0.48	0/2954
4	8	0.24	0/2130	0.51	0/2866
5	2	0.24	0/4594	0.49	0/6207
6	4	0.24	0/4178	0.49	0/5636
7	6	0.25	0/5172	0.49	0/6973
8	7	0.24	0/2884	0.51	0/3893
9	B	0.24	0/2294	0.45	0/3101
10	C	0.25	0/4788	0.45	0/6476
11	G	0.24	0/2705	0.47	0/3631
12	3	0.24	0/3324	0.48	0/4470
13	A	0.24	0/2938	0.47	0/3975
14	D	0.24	0/3373	0.43	0/4548
15	E	0.25	0/3076	0.45	0/4172
All	All	0.26	0/45446	0.50	0/61663

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	799	0	439	20	0
2	Y	800	0	438	17	0
3	5	2169	0	2216	13	0
4	8	2097	0	2135	39	0
5	2	4513	0	4537	79	0
6	4	4110	0	4160	74	0
7	6	5091	0	5115	76	0
8	7	2841	0	2852	46	0
9	B	2247	0	2236	29	0
10	C	4689	0	4724	79	0
11	G	2677	0	2830	30	0
12	3	3281	0	3329	34	0
13	A	2889	0	2905	32	0
14	D	3309	0	3343	23	0
15	E	3003	0	3033	28	0
16	2	31	0	12	2	0
16	6	31	0	12	5	0
16	A	31	0	12	2	0
16	D	31	0	12	3	0
16	E	31	0	12	2	0
17	2	1	0	0	0	0
17	6	1	0	0	0	0
17	A	1	0	0	0	0
17	D	1	0	0	0	0
17	E	1	0	0	0	0
18	4	1	0	0	0	0
All	All	44676	0	44352	570	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 570 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:2:222:MET:HG2	5:2:227:ARG:HB2	1.62	0.82
7:6:204:GLN:HB2	7:6:233:VAL:HG13	1.69	0.74
12:3:329:LEU:HD12	12:3:333:SER:HB3	1.67	0.74
10:C:130:THR:HG22	10:C:132:TYR:H	1.52	0.73
6:4:212:CYS:HB3	6:4:223:TYR:HE1	1.55	0.72

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	5	273/734 (37%)	267 (98%)	6 (2%)	0	100	100
4	8	253/546 (46%)	248 (98%)	5 (2%)	0	100	100
5	2	564/904 (62%)	558 (99%)	6 (1%)	0	100	100
6	4	498/863 (58%)	483 (97%)	15 (3%)	0	100	100
7	6	609/821 (74%)	597 (98%)	12 (2%)	0	100	100
8	7	349/719 (48%)	340 (97%)	9 (3%)	0	100	100
9	B	265/577 (46%)	260 (98%)	5 (2%)	0	100	100
10	C	556/711 (78%)	546 (98%)	10 (2%)	0	100	100
11	G	323/560 (58%)	318 (98%)	5 (2%)	0	100	100
12	3	415/810 (51%)	408 (98%)	7 (2%)	0	100	100
13	A	360/861 (42%)	348 (97%)	12 (3%)	0	100	100
14	D	402/436 (92%)	391 (97%)	11 (3%)	0	100	100
15	E	354/435 (81%)	346 (98%)	8 (2%)	0	100	100
All	All	5221/8977 (58%)	5110 (98%)	111 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	5	235/625 (38%)	235 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	8	230/456 (50%)	230 (100%)	0	100	100
5	2	496/781 (64%)	495 (100%)	1 (0%)	92	94
6	4	456/753 (61%)	456 (100%)	0	100	100
7	6	571/724 (79%)	571 (100%)	0	100	100
8	7	304/619 (49%)	304 (100%)	0	100	100
9	B	255/529 (48%)	254 (100%)	1 (0%)	89	91
10	C	535/659 (81%)	534 (100%)	1 (0%)	92	94
11	G	302/499 (60%)	301 (100%)	1 (0%)	91	92
12	3	355/708 (50%)	355 (100%)	0	100	100
13	A	321/766 (42%)	321 (100%)	0	100	100
14	D	375/403 (93%)	373 (100%)	2 (0%)	86	90
15	E	335/399 (84%)	335 (100%)	0	100	100
All	All	4770/7921 (60%)	4764 (100%)	6 (0%)	92	95

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
11	G	269	LYS
14	D	260	ARG
14	D	386	LYS
9	B	349	LYS
5	2	469	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
10	C	317	HIS
15	E	383	GLN
6	4	483	GLN
6	4	531	GLN
10	C	81	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
16	AGS	2	1001	17	26,33,33	0.71	1 (3%)	26,52,52	0.97	2 (7%)
16	AGS	A	901	17	26,33,33	0.71	1 (3%)	26,52,52	1.01	2 (7%)
16	AGS	D	901	17	26,33,33	0.71	1 (3%)	26,52,52	1.09	2 (7%)
16	AGS	6	901	17	26,33,33	0.71	1 (3%)	26,52,52	0.99	2 (7%)
16	AGS	E	501	17	26,33,33	0.71	1 (3%)	26,52,52	1.04	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
16	AGS	2	1001	17	-	3/17/38/38	0/3/3/3
16	AGS	A	901	17	-	3/17/38/38	0/3/3/3
16	AGS	D	901	17	-	3/17/38/38	0/3/3/3
16	AGS	6	901	17	-	4/17/38/38	0/3/3/3
16	AGS	E	501	17	-	2/17/38/38	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
16	A	901	AGS	PG-S1G	2.14	1.95	1.90
16	2	1001	AGS	PG-S1G	2.13	1.95	1.90
16	D	901	AGS	PG-S1G	2.11	1.95	1.90
16	6	901	AGS	PG-S1G	2.11	1.95	1.90
16	E	501	AGS	PG-S1G	2.11	1.95	1.90

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
16	D	901	AGS	PA-O3A-PB	-3.83	119.67	132.83
16	E	501	AGS	PA-O3A-PB	-3.58	120.53	132.83
16	A	901	AGS	PA-O3A-PB	-3.56	120.61	132.83
16	6	901	AGS	PA-O3A-PB	-3.37	121.27	132.83
16	2	1001	AGS	PA-O3A-PB	-2.96	122.67	132.83

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
16	2	1001	AGS	C5'-O5'-PA-O3A
16	6	901	AGS	C5'-O5'-PA-O3A
16	6	901	AGS	O4'-C4'-C5'-O5'
16	D	901	AGS	C5'-O5'-PA-O1A
16	D	901	AGS	C5'-O5'-PA-O2A


There are no ring outliers.



5 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
16	2	1001	AGS	2	0
16	A	901	AGS	2	0
16	D	901	AGS	3	0
16	6	901	AGS	5	0
16	E	501	AGS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19623. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 151




Y Index: 205




Z Index: 194


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.145. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_19623_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 568 nm³; this corresponds to an approximate mass of 513 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.263 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.80	-	-
Author-provided FSC curve	3.80	4.38	3.87
Unmasked-calculated*	7.84	10.38	8.11

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.84 differs from the reported value 3.8 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-19623 and PDB model 8S0E. Per-residue inclusion information can be found in section [3](#) on page [8](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.145 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.145).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.145) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7060	<div></div> 0.2400
2	<div></div> 0.8050	<div></div> 0.2610
3	<div></div> 0.2480	<div></div> 0.0230
4	<div></div> 0.7800	<div></div> 0.2410
5	<div></div> 0.2630	<div></div> 0.0300
6	<div></div> 0.8370	<div></div> 0.3220
7	<div></div> 0.5830	<div></div> 0.0980
8	<div></div> 0.4080	<div></div> 0.1290
A	<div></div> 0.8140	<div></div> 0.2900
B	<div></div> 0.7800	<div></div> 0.3160
C	<div></div> 0.8480	<div></div> 0.3070
D	<div></div> 0.8630	<div></div> 0.3540
E	<div></div> 0.8770	<div></div> 0.3840
G	<div></div> 0.5300	<div></div> 0.2000
X	<div></div> 0.8770	<div></div> 0.1980
Y	<div></div> 0.8650	<div></div> 0.2250

