



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 15, 2024 – 01:40 AM EST

PDB ID : 1S3V
Title : Structure Determination of Tetrahydroquinazoline Antifolates in Complex with Human and *Pneumocystis carinii* Dihydrofolate Reductase: Correlations of Enzyme Selectivity and Stereochemistry
Authors : Cody, V.; Luft, J.R.; Pangborn, W.; Gangjee, A.; Queener, S.F.
Deposited on : 2004-01-14
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	TQD	A	187	X	-	-	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1651 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrofolate reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	186	Total	C	N	O	S	0	0	0
			1502	963	253	279	7			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is (2R,6S)-6-{[methyl(3,4,5-trimethoxyphenyl)amino]methyl}-1,2,5,6,7,8-hexahydroquinazoline-2,4-diamine (three-letter code: TQD) (formula: C₁₉H₂₉N₅O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			27	19	5	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	112	Total O 112 112	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydrofolate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	106.97Å 106.97Å 43.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-1.80) 88.6 (50.00-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 1.80Å)	Xtriage
Refinement program	PROFFT	Depositor
R, R_{free}	0.173 , 0.192 0.172 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	21.3	Xtriage
Anisotropy	0.089	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 62.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1651	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TQD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.01	0/1537	2.08	50/2073 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (50) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ASN	CA-CB-CG	15.89	148.36	113.40
1	A	91	ARG	CD-NE-CZ	14.31	143.63	123.60
1	A	21	ASP	CB-CG-OD1	11.94	129.04	118.30
1	A	126	ASN	N-CA-CB	-10.39	91.89	110.60
1	A	65	ARG	NE-CZ-NH1	-10.28	115.16	120.30
1	A	33	TYR	CB-CG-CD2	-10.15	114.91	121.00
1	A	70	ARG	NE-CZ-NH1	10.10	125.35	120.30
1	A	91	ARG	NE-CZ-NH2	9.95	125.28	120.30
1	A	146	THR	CA-CB-CG2	-8.90	99.95	112.40
1	A	77	ARG	CG-CD-NE	8.77	130.22	111.80
1	A	126	ASN	CA-CB-CG	8.70	132.54	113.40
1	A	64	ASN	CA-CB-CG	7.76	130.48	113.40
1	A	180	GLU	CG-CD-OE1	7.75	133.79	118.30
1	A	162	TYR	CB-CG-CD1	-7.65	116.41	121.00
1	A	146	THR	O-C-N	7.64	134.92	122.70
1	A	162	TYR	CB-CG-CD2	7.58	125.55	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	77	ARG	NE-CZ-NH1	7.24	123.92	120.30
1	A	33	TYR	CB-CG-CD1	6.67	125.00	121.00
1	A	150	GLU	OE1-CD-OE2	6.62	131.25	123.30
1	A	33	TYR	CG-CD1-CE1	-6.48	116.11	121.30
1	A	81	GLU	OE1-CD-OE2	6.38	130.96	123.30
1	A	121	TYR	O-C-N	6.20	132.62	122.70
1	A	179	PHE	CB-CG-CD2	-6.16	116.49	120.80
1	A	126	ASN	CB-CA-C	5.93	122.26	110.40
1	A	182	TYR	CB-CG-CD2	5.91	124.54	121.00
1	A	152	ASP	CB-CG-OD2	-5.86	113.03	118.30
1	A	180	GLU	CG-CD-OE2	-5.76	106.78	118.30
1	A	146	THR	CA-C-O	-5.65	108.24	120.10
1	A	36	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	43	VAL	CA-CB-CG1	5.54	119.22	110.90
1	A	36	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	29	ASN	O-C-N	5.50	131.50	122.70
1	A	91	ARG	CG-CD-NE	5.49	123.34	111.80
1	A	108	LYS	CB-CA-C	-5.47	99.47	110.40
1	A	21	ASP	CB-CG-OD2	-5.46	113.39	118.30
1	A	37	MET	CG-SD-CE	5.46	108.93	100.20
1	A	75	LEU	CB-CA-C	5.33	120.32	110.20
1	A	143	GLU	CG-CD-OE1	5.32	128.95	118.30
1	A	172	GLU	OE1-CD-OE2	5.32	129.68	123.30
1	A	91	ARG	CB-CG-CD	5.26	125.28	111.60
1	A	133	LEU	CA-CB-CG	5.22	127.31	115.30
1	A	177	TYR	CA-CB-CG	5.21	123.30	113.40
1	A	78	GLU	CG-CD-OE2	-5.21	107.88	118.30
1	A	12	GLN	CG-CD-OE1	-5.16	111.28	121.60
1	A	59	SER	CA-CB-OG	-5.12	97.39	111.20
1	A	103	PRO	N-CA-CB	-5.05	97.04	102.60
1	A	91	ARG	NH1-CZ-NH2	-5.04	113.85	119.40
1	A	118	SER	O-C-N	5.02	130.74	122.70
1	A	32	ARG	NE-CZ-NH1	-5.01	117.80	120.30
1	A	123	GLU	CB-CG-CD	5.00	127.70	114.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	77	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1502	0	1511	48	1
2	A	10	0	0	1	0
3	A	27	0	28	1	0
4	A	112	0	0	4	1
All	All	1651	0	1539	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ASN:H	1:A:87:HIS:HD2	1.12	0.94
1:A:44:GLU:OE1	1:A:44:GLU:HA	1.69	0.90
1:A:72:ASN:H	1:A:87:HIS:CD2	1.91	0.87
1:A:111:MET:HE3	1:A:113:TRP:HE1	1.45	0.79
1:A:118:SER:O	1:A:122:LYS:HD3	1.84	0.78
1:A:130:HIS:HE1	1:A:183:GLU:CD	1.89	0.76
1:A:35:GLN:NE2	1:A:70:ARG:HH12	1.85	0.75
1:A:76:SER:OG	2:A:189:SO4:O1	2.04	0.73
1:A:75:LEU:HD11	1:A:92:SER:C	2.08	0.73
1:A:122:LYS:HD2	1:A:149:PRO:HG3	1.70	0.72
1:A:146:THR:HG21	4:A:219:HOH:O	1.95	0.65
1:A:125:MET:HG2	1:A:156:TYR:OH	1.98	0.64
1:A:72:ASN:N	1:A:87:HIS:HD2	1.92	0.62
1:A:35:GLN:HE21	1:A:35:GLN:HA	1.66	0.60
1:A:111:MET:HE1	4:A:207:HOH:O	2.05	0.57
1:A:63:LYS:HD2	1:A:64:ASN:OD1	2.05	0.57
1:A:184:LYS:NZ	1:A:186:ASP:OD2	2.36	0.57
1:A:2:GLY:HA3	1:A:110:ASP:O	2.06	0.55
1:A:91:ARG:O	1:A:91:ARG:HG2	2.06	0.55
1:A:130:HIS:CE1	1:A:183:GLU:CD	2.78	0.55
1:A:172:GLU:OE1	1:A:177:TYR:OH	2.20	0.54
1:A:130:HIS:HB2	1:A:185:ASN:CG	2.29	0.53
1:A:91:ARG:O	1:A:91:ARG:CG	2.56	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:130:HIS:HB2	1:A:185:ASN:ND2	2.23	0.52
1:A:100:THR:HA	1:A:105:LEU:O	2.10	0.51
1:A:130:HIS:HE1	1:A:183:GLU:OE1	1.93	0.51
1:A:168:ASP:O	1:A:170:GLN:NE2	2.41	0.50
1:A:1:VAL:HG12	1:A:2:GLY:N	2.27	0.49
1:A:101:GLU:O	1:A:106:ALA:HB2	2.12	0.49
1:A:35:GLN:HE21	1:A:70:ARG:HH12	1.60	0.49
1:A:130:HIS:CE1	1:A:183:GLU:HG3	2.48	0.49
1:A:111:MET:CE	1:A:113:TRP:HE1	2.20	0.49
1:A:104:GLU:O	1:A:108:LYS:HD2	2.12	0.48
1:A:102:GLN:HA	1:A:103:PRO:HD2	1.66	0.48
1:A:55:LYS:HE3	4:A:282:HOH:O	2.14	0.47
1:A:118:SER:HB3	1:A:147:PHE:O	2.13	0.47
1:A:80:LYS:HE2	4:A:235:HOH:O	2.15	0.46
1:A:75:LEU:HD11	1:A:92:SER:O	2.16	0.46
1:A:183:GLU:CG	1:A:184:LYS:N	2.79	0.46
1:A:93:LEU:HD22	1:A:97:LEU:HG	1.97	0.45
1:A:35:GLN:HE22	1:A:70:ARG:HH12	1.62	0.45
1:A:26:PRO:O	1:A:173:LYS:HE2	2.18	0.44
1:A:24:TRP:HB2	1:A:25:PRO:HD2	2.01	0.43
1:A:1:VAL:HG12	1:A:2:GLY:H	1.84	0.43
1:A:95:ASP:OD2	1:A:98:LYS:HE2	2.19	0.43
1:A:127:HIS:CD2	1:A:128:PRO:O	2.73	0.42
1:A:81:GLU:O	1:A:81:GLU:HG2	2.19	0.42
1:A:114:ILE:HD13	1:A:124:ALA:CB	2.50	0.41
3:A:187:TQD:H9'1	3:A:187:TQD:H2'1	1.78	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:ASP:OD1	4:A:238:HOH:O[5_554]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/186 (99%)	175 (95%)	8 (4%)	1 (0%)	25	14

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	103	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/168 (100%)	154 (92%)	14 (8%)	9	3

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	35	GLN
1	A	44	GLU
1	A	63	LYS
1	A	80	LYS
1	A	81	GLU
1	A	91	ARG
1	A	93	LEU
1	A	118	SER
1	A	122	LYS
1	A	157	LYS
1	A	161	GLU
1	A	167	SER
1	A	171	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN
1	A	87	HIS
1	A	102	GLN
1	A	127	HIS
1	A	130	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TQD	A	187	-	27,29,29	1.48	7 (25%)	28,41,41	2.49	5 (17%)
2	SO4	A	189	-	4,4,4	0.67	0	6,6,6	0.13	0
2	SO4	A	188	-	4,4,4	0.97	0	6,6,6	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TQD	A	187	-	2/2/6/7	0/14/39/39	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	187	TQD	C5D-C4A	3.48	1.56	1.51
3	A	187	TQD	C2D-N1'	-3.02	1.34	1.44
3	A	187	TQD	C8A-C4A	2.23	1.40	1.37
3	A	187	TQD	C3'-C4'	-2.19	1.36	1.41
3	A	187	TQD	C7'-C8	2.15	1.57	1.53
3	A	187	TQD	C4D-N3'	2.14	1.34	1.29
3	A	187	TQD	O3'-C3'	2.04	1.40	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	187	TQD	O4'-C4'-C5'	-7.93	108.76	120.12
3	A	187	TQD	O4'-C4'-C3'	6.70	129.74	120.12
3	A	187	TQD	O3'-C3'-C4'	4.34	122.57	115.14
3	A	187	TQD	C3B-O3'-C3'	3.40	122.50	117.51
3	A	187	TQD	O3'-C3'-C2'	-2.86	119.15	124.08

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	187	TQD	C6D
3	A	187	TQD	C2D

There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	187	TQD	1	0
2	A	189	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	186/186 (100%)	0.03	13 (6%)	24 21	10, 22, 56, 77	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	105	LEU	4.6
1	A	103	PRO	4.0
1	A	43	VAL	3.9
1	A	1	VAL	3.8
1	A	106	ALA	3.2
1	A	186	ASP	2.8
1	A	102	GLN	2.8
1	A	42	SER	2.6
1	A	104	GLU	2.6
1	A	45	GLY	2.3
1	A	2	GLY	2.3
1	A	153	LEU	2.2
1	A	155	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	A	189	5/5	0.94	0.07	52,53,55,55	0
3	TQD	A	187	27/27	0.97	0.05	8,17,24,26	0
2	SO4	A	188	5/5	0.99	0.10	19,21,23,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.