



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2025 – 12:33 PM EDT

PDB ID : 5S76 / pdb_00005s76
Title : XChem group deposition – Crystal Structure of human ACVR1 in complex with FM010916a
Authors : Williams, E.P.; Adamson, R.J.; Smil, D.; Krojer, T.; Burgess-Brown, N.; von Delft, F.; Bountra, C.; Bullock, A.N.
Deposited on : 2020-12-11
Resolution : 1.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2202 (1.34-1.30)
Clashscore	180529	2378 (1.34-1.30)
Ramachandran outliers	177936	2325 (1.34-1.30)
Sidechain outliers	177891	2325 (1.34-1.30)
RSRZ outliers	164620	2199 (1.34-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div>8%</div> <div>86%</div> <div>7% • 6%</div> </div>
1	B	301	<div> <div>5%</div> <div>88%</div> <div>9% •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	02A	A	509	X	-	-	-
6	02A	A	510	X	-	-	-
6	02A	B	505	X	-	-	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5793 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Activin receptor type-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	283	Total	C	N	O	S	0	14	0
			2356	1507	404	430	15			
1	B	292	Total	C	N	O	S	0	8	0
			2385	1519	415	435	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	SER	-	expression tag	UNP Q04771
A	200	MET	-	expression tag	UNP Q04771
A	207	ASP	GLN	engineered mutation	UNP Q04771
B	199	SER	-	expression tag	UNP Q04771
B	200	MET	-	expression tag	UNP Q04771
B	207	ASP	GLN	engineered mutation	UNP Q04771

- Molecule 2 is 4-methyl-3-[4-(1-methylpiperidin-4-yl)phenyl]-5-(3,4,5-trimethoxyphenyl)pyridine (CCD ID: LU8) (formula: C₂₇H₃₂N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	1
			64	54	4	6		
2	A	1	Total	C	N	O	0	0
			32	27	2	3		
2	A	1	Total	C	N	O	0	0
			32	27	2	3		
2	B	1	Total	C	N	O	0	0
			32	27	2	3		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: $C_4H_6O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 6 is (2S)-azetidine-2-carboxylic acid (CCD ID: 02A) (formula: $C_4H_7NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			7	4	1	2		
6	A	1	Total	C	N	O	0	0
			7	4	1	2		
6	B	1	Total	C	N	O	0	0
			7	4	1	2		

- Molecule 7 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	382	Total	O	0	1
			383	383		
8	B	405	Total	O	0	0
			405	405		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Activin receptor type-1



• Molecule 1: Activin receptor type-1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	126.93Å 84.82Å 87.93Å 90.00° 130.84° 90.00°	Depositor
Resolution (Å)	63.65 – 1.31 63.65 – 1.31	Depositor EDS
% Data completeness (in resolution range)	73.7 (63.65-1.31) 73.6 (63.65-1.31)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.31Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.155 , 0.173 0.165 , 0.181	Depositor DCC
R_{free} test set	6229 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.011 for -h+k-l,-l,-k 0.000 for -h-k-l,l,k 0.015 for -h-2*l,-k,l	Xtriage
F_o , F_c correlation	0.97	EDS
Total number of atoms	5793	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LU8, TLA, EDO, SO4, DMS, 02A

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/2417	0.93	3/3273 (0.1%)
1	B	0.80	0/2447	0.99	10/3319 (0.3%)
All	All	0.78	0/4864	0.96	13/6592 (0.2%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	454	ARG	NE-CZ-NH1	6.94	123.77	120.30
1	B	474	MET	CG-SD-CE	6.67	110.87	100.20
1	B	401	ARG	NE-CZ-NH2	-6.55	117.02	120.30
1	B	417	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	B	485	ARG	NE-CZ-NH2	-6.12	117.24	120.30
1	B	454	ARG	NE-CZ-NH2	-5.89	117.35	120.30
1	A	474	MET	CG-SD-CE	5.86	109.57	100.20
1	B	247	ARG	NE-CZ-NH2	-5.76	117.42	120.30
1	B	335	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	B	294	TYR	CB-CG-CD2	5.46	124.27	121.00
1	A	247	ARG	NE-CZ-NH2	-5.37	117.62	120.30
1	B	485	ARG	NE-CZ-NH1	5.29	122.95	120.30
1	A	375	ARG	NE-CZ-NH2	-5.24	117.68	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2356	0	2342	31	0
1	B	2385	0	2361	24	0
2	A	128	0	0	0	0
2	B	32	0	0	0	0
3	A	28	0	42	1	0
3	B	12	0	18	1	0
4	A	4	0	6	0	0
4	B	4	0	6	0	0
5	A	10	0	4	1	0
6	A	14	0	0	0	0
6	B	7	0	0	0	0
7	A	15	0	0	0	0
7	B	10	0	0	0	0
8	A	383	0	0	11	0
8	B	405	0	0	14	0
All	All	5793	0	4779	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:ARG:NE	8:A:602:HOH:O	1.95	0.99
1:B:260:GLU:OE2	8:B:601:HOH:O	1.88	0.89
1:B:457:ILE:HG13	1:B:475:LYS:HZ1	1.41	0.85
1:B:480:GLN:CG	8:B:651:HOH:O	2.29	0.80
1:B:307:ARG:NH1	8:B:602:HOH:O	2.15	0.79
1:A:297:LEU:H	1:A:297:LEU:HD13	1.48	0.78
1:B:260:GLU:CD	8:B:601:HOH:O	2.23	0.76
1:B:480:GLN:HB2	8:B:651:HOH:O	1.86	0.74
1:B:260:GLU:CG	8:B:601:HOH:O	2.36	0.72
1:B:457:ILE:HG13	1:B:475:LYS:NZ	2.04	0.71
1:A:293[B]:ASP:HB2	8:A:614:HOH:O	1.90	0.71
1:B:480:GLN:CB	8:B:651:HOH:O	2.37	0.71
1:A:247:ARG:NH2	8:A:604:HOH:O	2.18	0.68
1:A:302:THR:HG21	1:A:499:ASP:OD2	1.94	0.68
1:A:260:GLU:OE2	8:A:603:HOH:O	2.11	0.67
1:A:454:ARG:NH2	1:A:480[B]:GLN:OE1	2.29	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:PHE:CE2	1:A:400[B]:LYS:HE3	2.31	0.64
1:A:396:PHE:HE2	1:A:400[B]:LYS:HE3	1.63	0.63
1:A:407:PHE:HE2	1:A:411[B]:LEU:HD11	1.63	0.63
1:A:454:ARG:HH22	1:A:480[B]:GLN:CD	2.04	0.61
1:A:407:PHE:CE2	1:A:411[B]:LEU:HD11	2.35	0.61
1:A:442[A]:GLU:HG2	8:A:672:HOH:O	2.03	0.59
1:A:202:ARG:HD3	3:A:514:EDO:O1	2.04	0.58
1:A:407:PHE:CZ	1:A:411[B]:LEU:HD21	2.39	0.58
1:B:480:GLN:CD	8:B:651:HOH:O	2.42	0.57
1:B:260:GLU:OE2	1:B:345:LYS:NZ	2.31	0.57
1:A:247:ARG:NH1	8:A:604:HOH:O	2.33	0.56
1:B:475:LYS:HE2	8:B:904:HOH:O	2.05	0.56
1:A:297:LEU:HD13	1:A:297:LEU:N	2.21	0.55
1:B:252:TYR:CD1	1:B:265:PHE:HB2	2.43	0.54
1:A:475:LYS:HD3	8:A:797:HOH:O	2.07	0.54
1:A:297:LEU:H	1:A:297:LEU:CD1	2.20	0.53
1:A:247:ARG:CZ	8:A:604:HOH:O	2.57	0.51
1:A:297:LEU:N	1:A:297:LEU:CD1	2.74	0.50
1:A:396:PHE:HE2	1:A:400[B]:LYS:CE	2.25	0.50
1:A:247:ARG:CZ	8:A:602:HOH:O	2.49	0.50
1:B:208:ILE:CD1	1:B:282:ILE:HD13	2.42	0.49
1:B:208:ILE:HD11	1:B:282:ILE:HD13	1.93	0.49
1:B:370:VAL:HG22	8:B:811:HOH:O	2.12	0.49
1:A:412:TRP:CZ2	1:A:416:ARG:HD2	2.52	0.45
1:B:363:GLN:CB	8:B:964:HOH:O	2.65	0.44
1:B:480:GLN:H	3:B:507:EDO:H11	1.83	0.44
1:B:393:VAL:HG12	8:B:894:HOH:O	2.18	0.43
1:B:497:LYS:HG3	8:B:699:HOH:O	2.18	0.43
1:A:296:GLN:HB2	1:A:297:LEU:HD12	2.01	0.43
1:B:276:SER:HB3	8:B:636:HOH:O	2.18	0.43
1:B:240:ARG:HD3	1:B:241[B]:ASP:OD1	2.19	0.42
1:A:293[B]:ASP:CB	8:A:614:HOH:O	2.59	0.42
1:B:412:TRP:CZ2	1:B:416:ARG:HD2	2.54	0.42
1:A:247:ARG:HD3	8:A:909:HOH:O	2.20	0.42
1:A:273:ARG:HB3	5:A:506:TLA:H3	2.02	0.41
1:A:361:HIS:HB2	1:A:396:PHE:CD1	2.55	0.41
1:A:357:LEU:HG	1:A:375:ARG:HD3	2.02	0.41
1:A:492:LYS:O	1:A:496:THR:HG23	2.21	0.40
1:B:252:TYR:CE1	1:B:265:PHE:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/301 (97%)	288 (99%)	3 (1%)	0	100	100
1	B	296/301 (98%)	292 (99%)	4 (1%)	0	100	100
All	All	587/602 (98%)	580 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/270 (96%)	257 (100%)	1 (0%)	89	74
1	B	263/270 (97%)	261 (99%)	2 (1%)	79	53
All	All	521/540 (96%)	518 (99%)	3 (1%)	84	62

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	297	LEU
1	B	240	ARG
1	B	297	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	284	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	SO4	B	502	-	4,4,4	0.30	0	6,6,6	0.32	0
3	EDO	A	503	-	3,3,3	0.33	0	2,2,2	0.48	0
2	LU8	A	501[A]	-	35,35,35	1.00	1 (2%)	47,49,49	1.55	7 (14%)
7	SO4	A	513	-	4,4,4	0.20	0	6,6,6	0.20	0
3	EDO	A	515	-	3,3,3	0.57	0	2,2,2	0.36	0
3	EDO	A	516	-	3,3,3	0.15	0	2,2,2	0.20	0
3	EDO	A	502	-	3,3,3	0.71	0	2,2,2	0.63	0
2	LU8	B	501	-	35,35,35	1.26	5 (14%)	47,49,49	1.68	10 (21%)
2	LU8	A	508	-	35,35,35	1.43	5 (14%)	47,49,49	1.92	12 (25%)
3	EDO	B	507	-	3,3,3	0.26	0	2,2,2	0.10	0
7	SO4	A	511	-	4,4,4	0.26	0	6,6,6	0.32	0
2	LU8	A	501[B]	-	35,35,35	0.75	1 (2%)	47,49,49	1.15	4 (8%)
7	SO4	A	512	-	4,4,4	0.52	0	6,6,6	0.58	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	DMS	A	505	-	3,3,3	0.20	0	3,3,3	0.19	0
6	02A	A	509	-	6,7,7	3.18	2 (33%)	5,9,9	1.21	0
7	SO4	B	506	-	4,4,4	0.25	0	6,6,6	0.17	0
3	EDO	B	508	-	3,3,3	0.09	0	2,2,2	0.14	0
3	EDO	B	503	-	3,3,3	1.06	0	2,2,2	0.41	0
6	02A	B	505	-	6,7,7	3.66	4 (66%)	5,9,9	2.29	4 (80%)
3	EDO	A	514	-	3,3,3	0.14	0	2,2,2	0.30	0
3	EDO	A	517	-	3,3,3	0.34	0	2,2,2	0.32	0
4	DMS	B	504	-	3,3,3	0.27	0	3,3,3	0.12	0
3	EDO	A	504	-	3,3,3	0.24	0	2,2,2	0.84	0
5	TLA	A	506	-	9,9,9	1.00	0	12,12,12	0.97	0
2	LU8	A	507	-	35,35,35	1.14	3 (8%)	47,49,49	1.69	7 (14%)
6	02A	A	510	-	6,7,7	4.77	2 (33%)	5,9,9	1.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	503	-	-	1/1/1/1	-
2	LU8	A	501[A]	-	-	0/18/28/28	0/4/4/4
3	EDO	A	515	-	-	1/1/1/1	-
3	EDO	A	516	-	-	0/1/1/1	-
3	EDO	A	502	-	-	0/1/1/1	-
2	LU8	B	501	-	-	0/18/28/28	0/4/4/4
2	LU8	A	508	-	-	5/18/28/28	0/4/4/4
3	EDO	B	507	-	-	1/1/1/1	-
2	LU8	A	501[B]	-	-	2/18/28/28	0/4/4/4
6	02A	A	509	-	1/1/2/3	0/2/10/10	0/1/1/1
3	EDO	B	508	-	-	1/1/1/1	-
3	EDO	B	503	-	-	0/1/1/1	-
6	02A	B	505	-	1/1/2/3	0/2/10/10	0/1/1/1
3	EDO	A	514	-	-	1/1/1/1	-
3	EDO	A	517	-	-	1/1/1/1	-
5	TLA	A	506	-	-	10/12/12/12	-
3	EDO	A	504	-	-	0/1/1/1	-
2	LU8	A	507	-	-	0/18/28/28	0/4/4/4
6	02A	A	510	-	1/1/2/3	2/2/10/10	0/1/1/1

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	510	02A	C27-CA	-11.32	1.40	1.55
6	B	505	02A	C27-CA	-7.44	1.45	1.55
6	A	509	02A	C27-CA	-6.28	1.47	1.55
2	A	508	LU8	C26-C27	-3.66	1.32	1.38
6	B	505	02A	OXT-C	-3.63	1.19	1.30
6	A	509	02A	C27-C26	-3.31	1.49	1.54
2	A	507	LU8	C26-C05	3.19	1.45	1.39
2	A	501[A]	LU8	C12-C13	3.12	1.43	1.38
2	A	508	LU8	C14-C15	-3.05	1.46	1.52
2	B	501	LU8	O02-C03	2.84	1.41	1.37
2	B	501	LU8	C06-C24	2.63	1.44	1.40
2	A	508	LU8	C23-C11	2.59	1.44	1.39
2	B	501	LU8	C07-C06	-2.46	1.35	1.39
2	B	501	LU8	C12-C13	2.46	1.42	1.38
2	A	507	LU8	C10-C11	-2.45	1.45	1.49
2	A	507	LU8	O02-C03	2.35	1.41	1.37
2	A	508	LU8	C06-C24	-2.33	1.36	1.40
6	B	505	02A	CA-C	-2.33	1.46	1.52
6	B	505	02A	C27-C26	-2.20	1.50	1.54
2	B	501	LU8	C27-C30	-2.09	1.36	1.41
2	A	508	LU8	C26-C05	2.08	1.43	1.39
2	A	501[B]	LU8	C27-C30	-2.04	1.36	1.41
6	A	510	02A	C27-C26	-2.03	1.51	1.54

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	508	LU8	C20-N18-C17	5.27	117.93	109.54
2	A	507	LU8	C07-C06-C05	-5.19	112.06	119.56
2	A	501[A]	LU8	C17-C16-C15	5.05	116.92	111.00
2	A	508	LU8	C21-C20-N18	4.77	117.35	111.20
2	A	507	LU8	C26-C05-C06	-4.66	112.80	120.61
2	A	507	LU8	C17-C16-C15	-4.62	105.59	111.00
2	A	501[A]	LU8	C20-N18-C17	4.56	116.80	109.54
2	B	501	LU8	C23-C22-C14	-4.11	117.08	121.18
2	A	508	LU8	C23-C22-C14	-4.08	117.11	121.18
2	B	501	LU8	C22-C14-C13	4.05	123.33	118.30
2	A	508	LU8	C16-C17-N18	3.45	115.65	111.20
2	B	501	LU8	C20-N18-C17	3.38	114.92	109.54
2	A	508	LU8	C12-C13-C14	3.35	124.52	121.18
2	A	501[A]	LU8	C09-C10-C11	-3.35	114.72	119.56
2	A	508	LU8	C06-C07-N08	-3.18	119.80	124.27
2	B	501	LU8	C05-C06-C24	-3.14	117.29	122.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	LU8	C06-C07-N08	-3.04	119.99	124.27
2	A	501[A]	LU8	C11-C10-C24	2.99	127.40	122.47
2	A	507	LU8	C04-C05-C06	2.87	125.42	120.61
6	B	505	02A	C26-C27-CA	2.85	90.90	87.50
2	A	501[A]	LU8	C16-C17-N18	2.78	114.78	111.20
6	B	505	02A	OXT-C-CA	2.77	122.89	113.51
2	B	501	LU8	C19-N18-C20	-2.67	105.55	110.63
2	B	501	LU8	C06-C24-C10	-2.66	115.16	118.39
2	B	501	LU8	C13-C12-C11	-2.59	117.80	121.12
2	A	508	LU8	C04-C03-C30	2.50	123.00	120.22
2	A	508	LU8	C09-N08-C07	2.49	120.90	117.51
2	A	507	LU8	O02-C03-C04	-2.47	119.82	124.08
2	A	508	LU8	C06-C24-C10	2.39	121.30	118.39
2	A	501[A]	LU8	C29-O28-C27	2.38	121.01	117.51
2	B	501	LU8	C23-C11-C12	2.37	121.92	117.68
2	A	508	LU8	C27-C30-C03	-2.36	117.27	119.56
2	A	501[B]	LU8	C19-N18-C17	2.27	114.96	110.63
2	A	501[B]	LU8	C09-C10-C11	-2.26	116.29	119.56
6	B	505	02A	O-C-CA	-2.25	114.99	122.26
2	A	508	LU8	C29-O28-C27	2.25	120.82	117.51
2	B	501	LU8	C20-C21-C15	2.25	113.63	111.00
2	A	507	LU8	C05-C06-C24	2.24	126.16	122.47
2	A	501[B]	LU8	C16-C15-C14	-2.23	107.40	112.67
2	A	507	LU8	C04-C05-C26	2.16	121.51	118.35
2	A	501[A]	LU8	C06-C07-N08	-2.16	121.23	124.27
6	B	505	02A	C27-CA-C	-2.11	110.19	116.15
2	A	501[B]	LU8	C22-C14-C15	-2.07	115.83	121.09
2	A	508	LU8	C16-C15-C21	-2.04	105.25	109.68

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	509	02A	CA
6	A	510	02A	CA
6	B	505	02A	CA

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	506	TLA	O1-C1-C2-O2
5	A	506	TLA	O11-C1-C2-O2
5	A	506	TLA	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	506	TLA	C1-C2-C3-C4
5	A	506	TLA	O2-C2-C3-O3
5	A	506	TLA	O2-C2-C3-C4
5	A	506	TLA	O3-C3-C4-O4
5	A	506	TLA	O3-C3-C4-O41
5	A	506	TLA	O1-C1-C2-C3
5	A	506	TLA	O11-C1-C2-C3
6	A	510	02A	O-C-CA-C27
3	A	503	EDO	O1-C1-C2-O2
3	A	515	EDO	O1-C1-C2-O2
2	A	508	LU8	C04-C05-C06-C07
6	A	510	02A	OXT-C-CA-C27
3	B	507	EDO	O1-C1-C2-O2
3	B	508	EDO	O1-C1-C2-O2
3	A	514	EDO	O1-C1-C2-O2
2	A	508	LU8	C26-C05-C06-C07
2	A	501[B]	LU8	C22-C14-C15-C21
2	A	501[B]	LU8	C13-C14-C15-C21
2	A	508	LU8	C22-C14-C15-C21
2	A	508	LU8	C04-C05-C06-C24
2	A	508	LU8	C13-C14-C15-C21
3	A	517	EDO	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	507	EDO	1	0
3	A	514	EDO	1	0
5	A	506	TLA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient


equivalents in the CSD to analyse the geometry.




Ligand LU8 A 508




Bond lengths



Bond angles




Torsions




Rings


Ligand LU8 A 501 (B)




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/301 (94%)	0.23	23 (8%) 19 21	5, 14, 32, 49	14 (4%)
1	B	292/301 (97%)	0.23	16 (5%) 32 37	5, 15, 29, 41	8 (2%)
All	All	575/602 (95%)	0.23	39 (6%) 25 27	5, 14, 31, 49	22 (3%)

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	297	LEU	6.1
1	A	325	GLY	5.3
1	B	205	ALA	5.3
1	B	297	LEU	4.7
1	A	324	PHE	4.2
1	B	324	PHE	4.1
1	B	499	ASP	4.0
1	B	323	ILE	3.9
1	B	255	VAL	3.7
1	A	393	VAL	3.4
1	B	372	ASN	3.3
1	B	219	TYR	3.3
1	A	275	SER	3.1
1	B	229	GLY	3.1
1	A	395	CYS	3.0
1	A	394	ASP	3.0
1	A	227	TRP	2.9
1	B	498	ILE	2.8
1	B	328	GLY	2.8
1	B	211	LEU	2.7
1	A	239	SER	2.6
1	A	203	THR	2.6
1	A	208	ILE	2.6
1	A	260	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	364	SER	2.4
1	A	396	PHE	2.4
1	A	210	LEU	2.4
1	A	499	ASP	2.4
1	A	211	LEU	2.3
1	A	202	ARG	2.2
1	A	376	VAL	2.2
1	B	374	PRO	2.1
1	A	213	CYS	2.1
1	B	393	VAL	2.1
1	A	219	TYR	2.0
1	A	323	ILE	2.0
1	B	227	TRP	2.0
1	A	274	HIS	2.0
1	A	277	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	503	4/4	0.65	0.21	35,40,40,45	0
5	TLA	A	506	10/10	0.65	0.17	72,75,79,82	0
3	EDO	A	514	4/4	0.67	0.26	51,54,59,60	0
3	EDO	A	517	4/4	0.68	0.18	42,44,46,47	0
3	EDO	B	508	4/4	0.69	0.22	47,48,49,49	0
4	DMS	A	505	4/4	0.77	0.21	40,46,47,52	0
4	DMS	B	504	4/4	0.79	0.20	67,71,72,72	0
6	02A	A	510	7/7	0.81	0.39	1,2,3,3	7

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	02A	B	505	7/7	0.81	0.35	4,8,9,9	7
3	EDO	A	516	4/4	0.82	0.15	42,43,44,48	0
3	EDO	B	507	4/4	0.82	0.18	31,39,41,41	0
7	SO4	B	506	5/5	0.86	0.13	42,43,46,53	5
3	EDO	B	503	4/4	0.88	0.12	20,20,20,25	0
6	02A	A	509	7/7	0.89	0.33	3,4,5,5	7
2	LU8	A	507	32/32	0.90	0.11	16,19,35,42	0
3	EDO	A	515	4/4	0.92	0.11	25,28,31,40	0
3	EDO	A	502	4/4	0.93	0.10	17,17,18,22	0
2	LU8	A	501[B]	32/32	0.94	0.10	9,10,18,21	32
2	LU8	A	501[A]	32/32	0.94	0.10	10,12,30,36	32
2	LU8	A	508	32/32	0.96	0.08	13,15,36,44	0
2	LU8	B	501	32/32	0.96	0.08	11,13,37,41	0
7	SO4	B	502	5/5	0.97	0.06	20,21,23,24	0
3	EDO	A	504	4/4	0.97	0.07	12,18,19,22	0
7	SO4	A	511	5/5	0.98	0.05	15,16,22,22	0
7	SO4	A	512	5/5	0.98	0.07	13,17,26,27	0
7	SO4	A	513	5/5	0.99	0.05	11,13,15,15	5




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LU8 A 507:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LU8 A 501 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around LU8 A 501 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around LU8 A 508:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LU8 B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.