



## wwPDB EM Validation Summary Report ⓘ

May 5, 2025 – 08:16 PM EDT

PDB ID : 7SAE / pdb\_00007sae  
EMDB ID : EMD-24950  
Title : 44SR70P Class1 ribosomal particle  
Authors : Ortega, J.; Seffouh, A.  
Deposited on : 2021-09-22  
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev118  
MolProbity : 4-5-2 with Phenix2.0rc1  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.















Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	2928	
2	C	277	
3	D	209	
4	E	207	
5	J	145	
6	K	122	
7	L	146	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
8	N	120	
9	P	115	
10	Q	119	
11	R	102	
12	S	113	
13	T	95	
14	U	103	
15	V	94	
16	Z	59	
17	b	59	
18	Y	66	
19	d	44	

## 2 Entry composition

There are 19 unique types of molecules in this entry. The entry contains 68571 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (2434-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	2434	Total	C	N	O	P	0	0
			52303	23336	9696	16837	2434		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1558	C	G	conflict	GB 1864548803

- Molecule 2 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	C	267	Total	C	N	O	S	0	0
			2052	1277	402	367	6		

- Molecule 3 is a protein called 50S ribosomal protein L3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	D	179	Total	C	N	O	S	0	0
			1354	854	241	256	3		

- Molecule 4 is a protein called 50S ribosomal protein L4.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	E	205	Total	C	N	O	S	0	0
			1561	980	289	290	2		

- Molecule 5 is a protein called 50S ribosomal protein L13.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	J	142	Total	C	N	O	S	0	0
			1123	710	206	202	5		

- Molecule 6 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	K	122	Total	C	N	O	S	0	0
			920	571	173	172	4		

- Molecule 7 is a protein called 50S ribosomal protein L15.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	L	130	Total	C	N	O	S	0	0
			973	608	183	180	2		

- Molecule 8 is a protein called 50S ribosomal protein L17.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	N	119	Total	C	N	O	S	0	0
			953	583	186	180	4		

- Molecule 9 is a protein called 50S ribosomal protein L19.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	P	114	Total	C	N	O	S	0	0
			936	595	184	157			

- Molecule 10 is a protein called 50S ribosomal protein L20.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	Q	117	Total	C	N	O	S	0	0
			940	591	189	156	4		

- Molecule 11 is a protein called 50S ribosomal protein L21.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	R	101	Total	C	N	O	S	0	0
			780	495	139	146			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	74	HIS	PHE	conflict	UNP P26908
R	83	ALA	HIS	conflict	UNP P26908

- Molecule 12 is a protein called 50S ribosomal protein L22.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	S	109	Total	C	N	O	S	0	0
			842	525	164	150	3		

- Molecule 13 is a protein called 50S ribosomal protein L23.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	T	92	Total	C	N	O	S	0	0
			741	463	136	138	4		

- Molecule 14 is a protein called 50S ribosomal protein L24.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	U	100	Total	C	N	O	S	0	0
			754	473	141	137	3		

- Molecule 15 is a protein called 50S ribosomal protein L27.

Mol	Chain	Residues	Atoms				AltConf	Trace
15	V	72	Total	C	N	O	0	0
			561	349	109	103		

- Molecule 16 is a protein called 50S ribosomal protein L30.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	Z	58	Total	C	N	O	S	0	0
			455	281	89	84	1		

- Molecule 17 is a protein called 50S ribosomal protein L32.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	b	54	Total	C	N	O	S	0	0
			426	262	86	71	7		

- Molecule 18 is a protein called 50S ribosomal protein L29.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	Y	65	Total	C	N	O	S	0	0
			530	328	102	98	2		


- Molecule 19 is a protein called 50S ribosomal protein L34.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	d	44	Total	C	N	O	S	0	0
			367	222	89	54	2		


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: RNA (2434-MER)













• Molecule 2: 50S ribosomal protein L2




• Molecule 3: 50S ribosomal protein L3





• Molecule 4: 50S ribosomal protein L4








- Molecule 5: 50S ribosomal protein L13





- Molecule 6: 50S ribosomal protein L14




- Molecule 7: 50S ribosomal protein L15




- Molecule 8: 50S ribosomal protein L17




- Molecule 9: 50S ribosomal protein L19




- Molecule 10: 50S ribosomal protein L20







- Molecule 11: 50S ribosomal protein L21





- Molecule 12: 50S ribosomal protein L22



- Molecule 13: 50S ribosomal protein L23





- Molecule 14: 50S ribosomal protein L24





- Molecule 15: 50S ribosomal protein L27




- Molecule 16: 50S ribosomal protein L30





- Molecule 17: 50S ribosomal protein L32



• Molecule 18: 50S ribosomal protein L29



• Molecule 19: 50S ribosomal protein L34



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	168942	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	74	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.069	Depositor
Minimum map value	-0.014	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0131	Depositor
Map size ( $\text{\AA}$ )	362.52002, 362.52002, 362.52002	wwPDB
Map dimensions	424, 424, 424	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	0.855, 0.855, 0.855	Depositor

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/58595	0.48	7/91408 (0.0%)
2	C	0.29	0/2087	0.62	1/2798 (0.0%)
3	D	0.40	0/1367	0.62	2/1830 (0.1%)
4	E	0.41	0/1580	0.63	0/2132
5	J	0.40	0/1146	0.59	0/1542
6	K	0.29	0/927	0.60	0/1245
7	L	0.33	0/982	0.58	0/1308
8	N	0.49	0/960	0.64	0/1284
9	P	0.34	0/949	0.59	0/1269
10	Q	0.54	0/952	0.72	0/1266
11	R	0.42	0/790	0.81	1/1061 (0.1%)
12	S	0.42	0/851	0.61	0/1146
13	T	0.41	0/747	0.63	0/995
14	U	0.37	0/764	0.67	0/1022
15	V	0.28	0/569	0.56	0/757
16	Z	0.37	0/457	0.56	0/613
17	b	0.43	0/433	0.53	0/574
18	Y	0.35	0/531	0.60	0/707
19	d	0.46	0/370	0.57	0/483
All	All	0.41	0/75057	0.51	11/113440 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1
10	Q	0	1
11	R	0	1
All	All	0	3

There are no bond length outliers.

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	R	75	ARG	N-CA-C	7.14	119.27	110.91
1	A	880	C	O3'-P-O5'	7.03	114.55	104.00
3	D	74	THR	CA-C-N	6.33	130.31	120.68
3	D	74	THR	C-N-CA	6.33	130.31	120.68
2	C	133	ILE	N-CA-C	-5.84	107.57	113.47

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	154	LEU	Peptide
10	Q	87	GLY	Peptide
11	R	50	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	52303	0	26314	186	0
2	C	2052	0	2138	18	0
3	D	1354	0	1425	13	0
4	E	1561	0	1647	10	0
5	J	1123	0	1162	7	0
6	K	920	0	977	7	0
7	L	973	0	1032	6	0
8	N	953	0	983	9	0
9	P	936	0	1008	7	0
10	Q	940	0	1005	7	0
11	R	780	0	822	2	0
12	S	842	0	899	9	0
13	T	741	0	793	0	0
14	U	754	0	809	7	0
15	V	561	0	567	8	0
16	Z	455	0	491	4	0
17	b	426	0	445	1	0
18	Y	530	0	568	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
19	d	367	0	410	4	0
All	All	68571	0	43495	274	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 274 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2778:A:H62	1:A:2782:A:H61	1.05	1.00
1:A:2771:G:H1	1:A:2791:U:H3	0.94	0.91
1:A:2778:A:H62	1:A:2782:A:N6	1.70	0.90
1:A:912:C:H42	1:A:956:A:H62	1.35	0.74
1:A:2785:U:C4	1:A:2788:G:C6	2.75	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	263/277 (95%)	243 (92%)	20 (8%)	0	100	100
3	D	175/209 (84%)	166 (95%)	9 (5%)	0	100	100
4	E	203/207 (98%)	184 (91%)	19 (9%)	0	100	100
5	J	140/145 (97%)	135 (96%)	5 (4%)	0	100	100
6	K	120/122 (98%)	102 (85%)	18 (15%)	0	100	100
7	L	126/146 (86%)	116 (92%)	10 (8%)	0	100	100
8	N	117/120 (98%)	113 (97%)	4 (3%)	0	100	100
9	P	112/115 (97%)	103 (92%)	9 (8%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
10	Q	115/119 (97%)	104 (90%)	9 (8%)	2 (2%)	7	33
11	R	99/102 (97%)	87 (88%)	12 (12%)	0	100	100
12	S	107/113 (95%)	101 (94%)	6 (6%)	0	100	100
13	T	90/95 (95%)	80 (89%)	10 (11%)	0	100	100
14	U	98/103 (95%)	83 (85%)	15 (15%)	0	100	100
15	V	70/94 (74%)	67 (96%)	3 (4%)	0	100	100
16	Z	56/59 (95%)	54 (96%)	2 (4%)	0	100	100
17	b	52/59 (88%)	49 (94%)	3 (6%)	0	100	100
18	Y	63/66 (96%)	59 (94%)	4 (6%)	0	100	100
19	d	42/44 (96%)	40 (95%)	2 (5%)	0	100	100
All	All	2048/2195 (93%)	1886 (92%)	160 (8%)	2 (0%)	50	81

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
10	Q	88	ILE
10	Q	73	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	C	218/225 (97%)	218 (100%)	0	100	100
3	D	144/170 (85%)	144 (100%)	0	100	100
4	E	169/170 (99%)	169 (100%)	0	100	100
5	J	120/123 (98%)	120 (100%)	0	100	100
6	K	101/101 (100%)	100 (99%)	1 (1%)	73	88
7	L	101/110 (92%)	100 (99%)	1 (1%)	73	88
8	N	99/100 (99%)	98 (99%)	1 (1%)	73	88
9	P	99/100 (99%)	99 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	Q	96/98 (98%)	95 (99%)	1 (1%)	73	88
11	R	82/83 (99%)	82 (100%)	0	100	100
12	S	90/93 (97%)	90 (100%)	0	100	100
13	T	83/85 (98%)	83 (100%)	0	100	100
14	U	84/87 (97%)	84 (100%)	0	100	100
15	V	56/74 (76%)	56 (100%)	0	100	100
16	Z	52/53 (98%)	51 (98%)	1 (2%)	52	79
17	b	48/53 (91%)	48 (100%)	0	100	100
18	Y	56/57 (98%)	56 (100%)	0	100	100
19	d	39/39 (100%)	39 (100%)	0	100	100
All	All	1737/1821 (95%)	1732 (100%)	5 (0%)	90	96

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	K	58	VAL
7	L	105	LEU
8	N	117	ILE
10	Q	74	LEU
16	Z	54	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
10	Q	37	GLN
11	R	74	HIS
15	V	37	GLN
10	Q	107	ASN
11	R	81	ASN

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	2428/2928 (82%)	641 (26%)	38 (1%)

5 of 641 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	9	U
1	A	15	G
1	A	28	A
1	A	31	C
1	A	34	U

5 of 38 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	1779	G
1	A	2716	U
1	A	1784	A
1	A	2351	A
1	A	2812	A

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-24950. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 212



Y Index: 212




Z Index: 212


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 206



Y Index: 227




Z Index: 217


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0131. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.




## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 582 nm<sup>3</sup>; this corresponds to an approximate mass of 526 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.333 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.333  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	3.02	3.26	3.06
Unmasked-calculated*	-	-	-

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-24950 and PDB model 7SAE. Per-residue inclusion information can be found in [section 3](#) on [page 8](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0131 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0131).









































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0131) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8780	 0.5430
A	 0.8990	 0.5380
C	 0.6980	 0.5450
D	 0.8580	 0.5760
E	 0.8270	 0.5490
J	 0.8780	 0.5610
K	 0.6440	 0.5350
L	 0.7150	 0.5240
N	 0.9260	 0.5900
P	 0.6980	 0.5460
Q	 0.9170	 0.5780
R	 0.8490	 0.5710
S	 0.9160	 0.5840
T	 0.8780	 0.5620
U	 0.7840	 0.5590
V	 0.6370	 0.5210
Y	 0.8090	 0.5390
Z	 0.8470	 0.5620
b	 0.9220	 0.5830
d	 0.9570	 0.6020

