



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 8, 2024 – 05:21 AM EDT

PDB ID : 5SJS
Title : CRYSTAL STRUCTURE OF HUMAN PHOSPHODIESTERASE 10 IN COMPLEX WITH c1(c(ccc(n1)C2CC2)Nc3cncnc3)C(Nc4c(ccnc4)C(=O)N(C)C)=O, micromolar IC50=0.087803
Authors : Joseph, C.; Benz, J.; Flohr, A.; Groebke-Zbinden, K.; Rudolph, M.G.
Deposited on : 2022-02-01
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>73%</div> <div>17%</div> <div>• 9%</div> </div>
1	B	343	<div> <div>71%</div> <div>19%</div> <div>• 8%</div> </div>
1	C	343	<div> <div>74%</div> <div>15%</div> <div>• 9%</div> </div>
1	D	343	<div> <div>%</div> <div>68%</div> <div>22%</div> <div>• 9%</div> </div>

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2541	1624	432	461	24			
1	B	315	Total	C	N	O	S	0	0	0
			2551	1630	434	463	24			
1	C	313	Total	C	N	O	S	0	0	0
			2541	1624	432	461	24			
1	D	313	Total	C	N	O	S	0	0	0
			2537	1622	433	458	24			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	447	GLY	-	expression tag	UNP Q9Y233
A	448	SER	-	expression tag	UNP Q9Y233
B	447	GLY	-	expression tag	UNP Q9Y233
B	448	SER	-	expression tag	UNP Q9Y233
C	447	GLY	-	expression tag	UNP Q9Y233
C	448	SER	-	expression tag	UNP Q9Y233
D	447	GLY	-	expression tag	UNP Q9Y233
D	448	SER	-	expression tag	UNP Q9Y233

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		
2	D	1	Total	Zn	0	0
			1	1		


- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0
4	C	1	Total Mg 1 1	0	0
4	D	1	Total Mg 1 1	0	0

- Molecule 5 is 6-cyclopropyl-N-[4-(dimethylcarbamoyl)pyridin-3-yl]-3-[(pyrimidin-5-yl)amino]pyridine-2-carboxamide (three-letter code: K5R) (formula: C₂₁H₂₁N₇O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 30 21 7 2	0	0
5	B	1	Total C N O 30 21 7 2	0	0
5	C	1	Total C N O 30 21 7 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			30	21	7	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	45	Total	O	0	0
			45	45		
7	B	51	Total	O	0	0
			51	51		
7	C	56	Total	O	0	0
			56	56		
7	D	16	Total	O	0	0
			16	16		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



- Molecule 1: cAMP and cAMP-inhibited cGMP 3',5'-cyclic phosphodiesterase 10A



1692	C560	GLY
D701	H563	SER
E702	D564	ILE
K703		CYS
K704	H637	THR
K705		SER
Q708	L575	GLU
Q709	Q576	GLU
Y730	K577	GLU
A734	F578	
I735	D579	L459
P736	H580	F462
C737	F581	E471
P746	L582	F477
L751	Y586	F490
C755	S587	V491
N758	T588	Y492
Q761	Q593	M493
V765	F596	V494
I766	S601	H495
R767	S615	R496
G768	S616	S497
GLU	S617	C498
GLU	V622	G499
THR		T500
THR	I631	E504
TRP	Y638	R510
ILE		F511
SER	M643	V515
SER	I666	K516
PRO		K517
SER	T671	Y524
VAL	A672	V533
GLN	C673	A534
LYS	D674	H535
ALA	L675	C536
ALA	C676	M537
ALA	S677	Y538
SER	L679	A539
SER	K680	I540
GLU	L681	L541
ASP	P683	Q542
	A689	T549
		E552
		R553
		L556

4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	135.23Å 135.23Å 234.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.48 – 2.70 43.48 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.1 (43.48-2.70) 96.1 (43.48-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.173 , 0.254 0.182 , 0.255	Depositor DCC
R_{free} test set	2152 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 32.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10473	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MG, K5R, CME, ZN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/2592	0.93	0/3507
1	B	0.76	0/2602	0.97	1/3521 (0.0%)
1	C	0.75	0/2592	0.96	2/3507 (0.1%)
1	D	0.80	0/2588	0.91	0/3502
All	All	0.77	0/10374	0.94	3/14037 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	690	ASN	CB-CA-C	-5.77	98.86	110.40
1	B	664	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	C	553	ARG	NE-CZ-NH1	5.17	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2541	0	2513	36	0
1	B	2551	0	2515	35	0
1	C	2541	0	2511	33	0
1	D	2537	0	2511	45	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	30	0	0	4	0
5	B	30	0	0	5	0
5	C	30	0	0	4	0
5	D	30	0	0	1	0
6	B	6	0	8	1	0
7	A	45	0	0	2	0
7	B	51	0	0	2	1
7	C	56	0	0	2	2
7	D	16	0	0	3	1
All	All	10473	0	10058	157	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:804:K5R:C19	5:A:804:K5R:C27	2.53	0.84
1:C:493:MET:O	1:C:497:SER:HB2	1.84	0.77
1:B:683:PRO:O	1:B:687:LEU:HD12	1.85	0.76
1:D:490:PHE:O	1:D:494:VAL:HG23	1.88	0.73
5:C:803:K5R:C19	5:C:803:K5R:C27	2.69	0.70

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:941:HOH:O	7:C:941:HOH:O[2_655]	1.78	0.42
7:B:907:HOH:O	7:D:910:HOH:O[8_544]	1.84	0.36
7:C:905:HOH:O	7:C:905:HOH:O[2_655]	2.19	0.01

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/343 (90%)	283 (91%)	25 (8%)	2 (1%)	22	45
1	B	312/343 (91%)	286 (92%)	23 (7%)	3 (1%)	13	33
1	C	310/343 (90%)	285 (92%)	25 (8%)	0	100	100
1	D	310/343 (90%)	291 (94%)	17 (6%)	2 (1%)	22	45
All	All	1242/1372 (90%)	1145 (92%)	90 (7%)	7 (1%)	22	45

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	612	SER
1	B	652	GLY
1	D	615	SER
1	A	483	GLU
1	D	579	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/305 (92%)	264 (94%)	17 (6%)	16	38
1	B	281/305 (92%)	262 (93%)	19 (7%)	13	32
1	C	281/305 (92%)	259 (92%)	22 (8%)	10	26
1	D	280/305 (92%)	265 (95%)	15 (5%)	18	42
All	All	1123/1220 (92%)	1050 (94%)	73 (6%)	14	34

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	760	SER
1	D	709	GLN
1	D	500	THR
1	D	616	SER
1	B	551	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	476	HIS
1	D	621	GLN
1	D	743	GLN
1	D	650	GLN
1	B	604	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CME	B	509	1	8,9,10	0.47	0	6,9,11	0.86	0
1	CME	A	509	1	8,9,10	0.58	0	6,9,11	0.72	0
1	CME	D	509	1	8,9,10	0.41	0	6,9,11	0.82	0
1	CME	C	509	1	8,9,10	0.42	0	6,9,11	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	B	509	1	-	1/5/8/10	-
1	CME	A	509	1	-	2/5/8/10	-
1	CME	D	509	1	-	2/5/8/10	-
1	CME	C	509	1	-	3/5/8/10	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	509	CME	SD-CE-CZ-OH
1	C	509	CME	CE-SD-SG-CB
1	A	509	CME	SD-CE-CZ-OH
1	C	509	CME	SD-CE-CZ-OH
1	A	509	CME	CZ-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 9 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	K5R	C	803	-	33,33,33	1.67	7 (21%)	42,46,46	2.18	15 (35%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	K5R	D	803	-	33,33,33	1.65	11 (33%)	42,46,46	2.16	9 (21%)
6	GOL	B	803	-	5,5,5	0.25	0	5,5,5	0.62	0
5	K5R	A	804	-	33,33,33	2.14	8 (24%)	42,46,46	2.52	18 (42%)
5	K5R	B	804	-	33,33,33	2.38	9 (27%)	42,46,46	2.56	15 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	K5R	C	803	-	-	2/24/26/26	0/4/4/4
5	K5R	D	803	-	-	0/24/26/26	0/4/4/4
6	GOL	B	803	-	-	2/4/4/4	-
5	K5R	A	804	-	-	2/24/26/26	0/4/4/4
5	K5R	B	804	-	-	3/24/26/26	0/4/4/4

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	804	K5R	C5-N14	8.01	1.46	1.34
5	A	804	K5R	C5-N14	6.93	1.44	1.34
5	A	804	K5R	C4-C8	5.35	1.49	1.41
5	C	803	K5R	C5-N14	4.41	1.40	1.34
5	B	804	K5R	C4-C8	4.16	1.47	1.41

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	804	K5R	C4-C5-N14	7.98	126.63	118.32
5	B	804	K5R	C27-N18-C23	7.16	124.26	115.80
5	D	803	K5R	C27-N18-C23	6.37	123.32	115.80
5	C	803	K5R	C27-N18-C23	6.02	122.91	115.80
5	B	804	K5R	C8-C4-C5	5.77	128.76	120.72

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	803	K5R	C13-C10-C9-N2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	803	GOL	O1-C1-C2-C3
5	B	804	K5R	C13-C10-C9-C22
5	B	804	K5R	C12-C10-C9-C22
5	C	803	K5R	C12-C10-C9-C22


There are no ring outliers.

5 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	803	K5R	4	0
5	D	803	K5R	1	0
6	B	803	GOL	1	0
5	A	804	K5R	4	0
5	B	804	K5R	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand K5R C 803




Bond lengths



Bond angles




Torsions




Rings


Ligand K5R D 803




Bond lengths



Bond angles




Torsions




Rings


Ligand K5R A 804




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	312/343 (90%)	-0.62	1 (0%) 90 89	37, 52, 84, 119	0
1	B	314/343 (91%)	-0.54	1 (0%) 90 89	35, 55, 91, 112	0
1	C	312/343 (90%)	-0.65	1 (0%) 90 89	37, 52, 83, 111	0
1	D	312/343 (90%)	-0.07	3 (0%) 79 79	63, 88, 109, 126	0
All	All	1250/1372 (91%)	-0.47	6 (0%) 87 86	35, 60, 102, 126	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	457	GLN	2.8
1	D	459	LEU	2.7
1	C	459	LEU	2.5
1	A	459	LEU	2.4
1	D	737	CYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CME	B	509	10/11	0.86	0.15	58,71,100,105	0
1	CME	C	509	10/11	0.87	0.12	52,67,108,116	0
1	CME	D	509	10/11	0.88	0.12	77,100,124,124	0
1	CME	A	509	10/11	0.92	0.10	59,66,91,97	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	D	802	1/1	0.42	0.21	72,72,72,72	0
3	CL	A	802	1/1	0.65	0.11	101,101,101,101	0
5	K5R	D	803	30/30	0.82	0.16	63,94,120,122	0
4	MG	A	803	1/1	0.84	0.09	34,34,34,34	0
5	K5R	B	804	30/30	0.87	0.13	49,69,83,90	0
5	K5R	C	803	30/30	0.89	0.13	53,68,91,105	0
6	GOL	B	803	6/6	0.89	0.18	74,88,91,92	0
4	MG	C	802	1/1	0.91	0.08	38,38,38,38	0
5	K5R	A	804	30/30	0.92	0.11	52,65,81,88	0
2	ZN	D	801	1/1	0.99	0.03	73,73,73,73	0
2	ZN	B	801	1/1	0.99	0.02	45,45,45,45	0
2	ZN	A	801	1/1	1.00	0.01	49,49,49,49	0
4	MG	B	802	1/1	1.00	0.02	36,36,36,36	0
2	ZN	C	801	1/1	1.00	0.01	48,48,48,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around K5R D 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around K5R B 804:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K5R C 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around K5R A 804:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.