



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 09:44 PM EDT

PDB ID : 6TE1
Title : Structure of the KDM1A/CoREST complex with the inhibitor 2-[3-{4-chloro-3-[(4-chlorophenyl)ethynyl]phenyl}-1-(3-morpholin-4-ylpropyl)-1,4,6,7-tetrahydro-5H-pyrazolo[4,3-c]pyridin-5-yl]-2-oxoethanol
Authors : Pasqualato, S.; Cecatiello, V.
Deposited on : 2019-11-11
Resolution : 3.11 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 3.11 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1292 (3.14-3.10)
Clashscore	141614	1389 (3.14-3.10)
Ramachandran outliers	138981	1337 (3.14-3.10)
Sidechain outliers	138945	1337 (3.14-3.10)
RSRZ outliers	127900	1260 (3.14-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	852	
2	B	482	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	905	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Lysine-specific histone demethylase 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	666	Total	C	N	O	S	0	0	0
			5217	3324	906	967	20			

- Molecule 2 is a protein called REST corepressor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	133	Total	C	N	O	S	0	0	0
			1076	676	194	203	3			

- Molecule 3 is 5-[4-cyclobutyl-1-[2-(4-piperidin-4-yloxyphenoxy)ethyl]imidazol-2-yl]-4-methyl-thieno[3,2-b]pyrrole (three-letter code: N4K) (formula: C₂₇H₃₂N₄O₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			34	27	4	2	1		

Continued on next page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			34	27	4	2	1		

- # FAD


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- GOL
-
- The diagram shows the skeletal structure of 1,2,3-propanetriol (glycerol). It consists of a three-carbon chain. The first carbon (C1) is bonded to a hydroxyl group (HO) labeled O1. The second carbon (C2) is bonded to a hydroxyl group (OH) labeled O2. The third carbon (C3) is bonded to a hydroxyl group (OH) labeled O3. The labels C1, C2, and C3 are in green, while the labels O1, O2, and O3 are in green. The hydroxyl groups are written in red.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	14	Total	O	0	0
			14	14		



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	120.31 Å 179.74 Å 236.77 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.05 – 3.11 46.23 – 3.11	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.05-3.11) 99.8 (46.23-3.11)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.17 (at 3.12 Å)	Xtriage
Refinement program	PHENIX 1.17_3644	Depositor
R, R_{free}	0.168 , 0.192 0.168 , 0.192	Depositor DCC
R_{free} test set	2315 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	87.0	Xtriage
Anisotropy	0.331	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6458	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, N4K, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/5331	0.42	0/7232
2	B	0.25	0/1091	0.38	0/1471
All	All	0.25	0/6422	0.41	0/8703

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5217	0	5252	40	0
2	B	1076	0	1091	9	0
3	A	68	0	0	0	0
4	A	53	0	31	2	0
5	A	30	0	40	1	0
6	A	14	0	0	0	0
All	All	6458	0	6414	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:716:GLU:O	1:A:750:ARG:NH1	2.13	0.82
1:A:332:MET:H	1:A:571:TYR:HD2	1.32	0.75
1:A:495:ASP:OD1	2:B:371:ARG:NH2	2.33	0.62
1:A:384:ARG:HH22	2:B:313:GLY:HA3	1.65	0.60
1:A:526:ARG:NH2	5:A:906:GOL:O1	2.37	0.58
1:A:384:ARG:HB3	2:B:314:MET:HE3	1.86	0.57
1:A:661:LYS:HB3	1:A:704:LEU:HD21	1.86	0.57
1:A:801:GLU:HG2	1:A:809:ALA:H	1.73	0.54
1:A:331:ALA:HA	4:A:903:FAD:N5	2.23	0.53
1:A:306:LEU:HD13	1:A:584:ILE:HG12	1.92	0.52
1:A:463:LYS:O	1:A:467:GLU:HG2	2.10	0.52
1:A:760:SER:HB2	4:A:903:FAD:HM83	1.92	0.51
1:A:594:ARG:HG2	1:A:640:VAL:HB	1.92	0.51
1:A:238:LEU:HB3	1:A:243:ASN:HB3	1.95	0.48
1:A:780:ILE:HG12	1:A:798:PHE:HE1	1.78	0.48
1:A:541:ALA:HB2	1:A:659:LEU:HD23	1.94	0.48
1:A:209:VAL:HG13	1:A:242:TYR:HD1	1.78	0.48
1:A:364:GLU:HA	1:A:681:VAL:HB	1.95	0.48
1:A:319:THR:OG1	1:A:328:ASP:OD1	2.23	0.47
1:A:442:LYS:HE3	2:B:355:THR:HG21	1.97	0.47
1:A:707:VAL:HG12	1:A:712:ALA:HA	1.97	0.46
1:A:669:VAL:HG13	1:A:671:TRP:CE2	2.50	0.46
1:A:458:LEU:HB3	1:A:487:LEU:HD13	1.97	0.46
1:A:601:GLU:HB3	1:A:617:LYS:HD3	1.98	0.45
2:B:432:GLU:O	2:B:435:GLN:HG2	2.16	0.45
1:A:780:ILE:HB	1:A:796:LEU:HB3	1.98	0.44
1:A:289:SER:HB3	1:A:814:ALA:HB1	2.01	0.43
1:A:384:ARG:NH1	2:B:312:LYS:O	2.36	0.43
1:A:458:LEU:HD23	1:A:458:LEU:HA	1.89	0.43
1:A:699:LYS:H	1:A:699:LYS:HG3	1.49	0.43
2:B:421:PHE:O	2:B:425:ARG:HB2	2.19	0.43
1:A:435:VAL:HG13	2:B:349:ILE:HG12	2.01	0.42
1:A:534:ALA:HB2	1:A:688:ARG:HG3	2.01	0.42
1:A:318:ALA:H	1:A:330:GLY:HA3	1.84	0.42
1:A:364:GLU:OE2	1:A:524:ARG:NH2	2.52	0.42
1:A:654:MET:HE2	1:A:654:MET:HB3	1.76	0.42
1:A:487:LEU:HD23	2:B:372:LEU:HD11	2.01	0.41
1:A:447:LYS:HB2	1:A:447:LYS:HE3	1.86	0.41
1:A:282:ILE:HD13	1:A:305:THR:HB	2.03	0.41
1:A:548:SER:O	1:A:552:TRP:HB3	2.21	0.40
1:A:353:LEU:HB3	1:A:565:LEU:HD22	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:695:TRP:HE1	1:A:706:LEU:HD22	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/852 (78%)	648 (98%)	16 (2%)	0	100	100
2	B	131/482 (27%)	123 (94%)	8 (6%)	0	100	100
All	All	795/1334 (60%)	771 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	566/699 (81%)	560 (99%)	6 (1%)	73	88
2	B	117/395 (30%)	116 (99%)	1 (1%)	78	91
All	All	683/1094 (62%)	676 (99%)	7 (1%)	76	89

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	295	ARG
1	A	429	GLU
1	A	458	LEU
1	A	568	ARG
1	A	571	TYR
1	A	591	ARG
2	B	347	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	904	-	5,5,5	0.98	0	5,5,5	1.14	0
5	GOL	A	906	-	5,5,5	0.92	0	5,5,5	1.16	1 (20%)
5	GOL	A	905	-	5,5,5	0.96	0	5,5,5	1.06	0
5	GOL	A	908	-	5,5,5	0.94	0	5,5,5	1.07	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	N4K	A	901	-	35,39,39	2.56	4 (11%)	36,55,55	1.99	5 (13%)
5	GOL	A	907	-	5,5,5	0.86	0	5,5,5	1.11	0
3	N4K	A	902	-	35,39,39	2.70	5 (14%)	36,55,55	1.92	5 (13%)
4	FAD	A	903	-	54,58,58	0.49	0	71,89,89	0.54	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	904	-	-	2/4/4/4	-
5	GOL	A	906	-	-	2/4/4/4	-
5	GOL	A	905	-	-	2/4/4/4	-
5	GOL	A	908	-	-	2/4/4/4	-
3	N4K	A	901	-	-	8/12/32/32	0/6/6/6
5	GOL	A	907	-	-	2/4/4/4	-
3	N4K	A	902	-	-	7/12/32/32	0/6/6/6
4	FAD	A	903	-	-	4/30/50/50	0/6/6/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	902	N4K	C30-S34	-13.54	1.63	1.74
3	A	901	N4K	C30-S34	-12.87	1.64	1.74
3	A	902	N4K	C11-N12	-5.69	1.31	1.38
3	A	901	N4K	C11-N12	-5.17	1.31	1.38
3	A	901	N4K	C32-C31	2.81	1.54	1.42
3	A	902	N4K	C32-C31	2.77	1.54	1.42
3	A	902	N4K	C03-C04	2.64	1.53	1.48
3	A	901	N4K	C03-C04	2.54	1.53	1.48
3	A	902	N4K	C33-S34	-2.12	1.61	1.71

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	902	N4K	C33-S34-C30	8.83	101.19	91.55
3	A	901	N4K	C33-S34-C30	8.35	100.67	91.55
3	A	901	N4K	C19-O20-C21	-4.23	111.53	119.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	N4K	O15-C14-C13	4.20	111.57	107.28
3	A	901	N4K	C07-C06-N05	4.06	126.23	120.23
3	A	902	N4K	C07-C06-N05	4.04	126.19	120.23
3	A	902	N4K	C19-O20-C21	-3.02	113.72	119.18
3	A	902	N4K	O20-C21-C22	-2.98	101.37	108.31
3	A	902	N4K	C11-C06-C07	-2.45	125.88	129.26
4	A	903	FAD	C5A-C6A-N6A	2.43	124.01	120.31
3	A	901	N4K	C11-C06-C07	-2.09	126.37	129.26
5	A	906	GOL	C3-C2-C1	-2.08	104.15	111.80

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	N4K	N12-C13-C14-O15
3	A	901	N4K	C14-C13-N12-C11
5	A	904	GOL	O1-C1-C2-C3
5	A	906	GOL	C1-C2-C3-O3
5	A	907	GOL	O1-C1-C2-C3
5	A	908	GOL	O1-C1-C2-O2
5	A	908	GOL	O1-C1-C2-C3
3	A	901	N4K	C27-C19-O20-C21
3	A	901	N4K	C18-C19-O20-C21
5	A	905	GOL	O1-C1-C2-C3
5	A	904	GOL	O1-C1-C2-O2
5	A	906	GOL	O2-C2-C3-O3
5	A	907	GOL	O1-C1-C2-O2
3	A	902	N4K	N12-C13-C14-O15
4	A	903	FAD	C2'-C3'-C4'-O4'
5	A	905	GOL	O1-C1-C2-O2
4	A	903	FAD	C2'-C3'-C4'-C5'
3	A	901	N4K	C14-C13-N12-C04
4	A	903	FAD	O3'-C3'-C4'-O4'
3	A	902	N4K	C27-C19-O20-C21
3	A	902	N4K	C18-C19-O20-C21
4	A	903	FAD	O3'-C3'-C4'-C5'
3	A	902	N4K	C13-C14-O15-C16
3	A	901	N4K	C22-C21-O20-C19
3	A	902	N4K	C22-C21-O20-C19
3	A	902	N4K	C26-C21-O20-C19
3	A	901	N4K	C26-C21-O20-C19
3	A	901	N4K	C11-C06-C07-C10

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	902	N4K	C11-C06-C07-C10

There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	906	GOL	1	0
4	A	903	FAD	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand N4K A 902



Ligand FAD A 903



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	666/852 (78%)	-0.19	1 (0%) 95 91	45, 83, 132, 167	0
2	B	133/482 (27%)	0.00	0 100 100	70, 114, 150, 169	0
All	All	799/1334 (59%)	-0.16	1 (0%) 95 92	45, 88, 138, 169	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	171	PRO	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	905	6/6	0.70	0.48	126,133,137,141	0
5	GOL	A	908	6/6	0.82	0.31	108,127,146,157	0
5	GOL	A	907	6/6	0.87	0.32	114,120,123,130	0
5	GOL	A	904	6/6	0.91	0.26	92,101,102,107	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	N4K	A	901	34/34	0.91	0.29	94,118,166,224	0
3	N4K	A	902	34/34	0.95	0.28	44,60,90,99	0
5	GOL	A	906	6/6	0.96	0.23	74,83,89,106	0
4	FAD	A	903	53/53	0.97	0.23	34,55,70,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around N4K A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD A 903:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.