



## Full wwPDB EM Validation Report ⓘ

Oct 6, 2024 – 10:22 PM EDT

PDB ID : 7TLB  
EMDB ID : EMD-25986  
Title : Down-state locked rS2d SARS-CoV-2 spike ectodomain in the RBD-down conformation, State 2  
Authors : Henderson, R.; Acharya, P.  
Deposited on : 2022-01-18  
Resolution : 3.06 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1208	<p>9% 78% 20%</p>
1	B	1208	<p>8% 78% 20%</p>
1	C	1208	<p>9% 78% 20%</p>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 22806 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein.


Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	972	Total	C	N	O	S	0	0
			7602	4860	1262	1445	35		
1	B	972	Total	C	N	O	S	0	0
			7602	4860	1262	1445	35		
1	C	972	Total	C	N	O	S	0	0
			7602	4860	1262	1445	35		

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	383	CYS	SER	conflict	UNP P0DTC2
A	682	GLY	ARG	conflict	UNP P0DTC2
A	683	SER	ARG	conflict	UNP P0DTC2
A	685	SER	ARG	conflict	UNP P0DTC2
A	985	CYS	ASP	conflict	UNP P0DTC2
A	986	PRO	LYS	conflict	UNP P0DTC2
A	987	PRO	VAL	conflict	UNP P0DTC2
B	383	CYS	SER	conflict	UNP P0DTC2
B	682	GLY	ARG	conflict	UNP P0DTC2
B	683	SER	ARG	conflict	UNP P0DTC2
B	685	SER	ARG	conflict	UNP P0DTC2
B	985	CYS	ASP	conflict	UNP P0DTC2
B	986	PRO	LYS	conflict	UNP P0DTC2
B	987	PRO	VAL	conflict	UNP P0DTC2
C	383	CYS	SER	conflict	UNP P0DTC2
C	682	GLY	ARG	conflict	UNP P0DTC2
C	683	SER	ARG	conflict	UNP P0DTC2
C	685	SER	ARG	conflict	UNP P0DTC2
C	985	CYS	ASP	conflict	UNP P0DTC2
C	986	PRO	LYS	conflict	UNP P0DTC2
C	987	PRO	VAL	conflict	UNP P0DTC2







## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	218600	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	3.508	Depositor
Minimum map value	-2.271	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.085	Depositor
Recommended contour level	0.432	Depositor
Map size (Å)	317.4, 317.4, 317.4	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.058, 1.058, 1.058	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.65	0/7770	0.94	22/10569 (0.2%)
1	B	0.65	0/7770	0.94	23/10569 (0.2%)
1	C	0.64	0/7770	0.93	19/10569 (0.2%)
All	All	0.65	0/23310	0.94	64/31707 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	5
All	All	0	8

There are no bond length outliers.

All (64) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	904	TYR	CB-CG-CD2	-8.41	115.95	121.00
1	C	904	TYR	CB-CG-CD2	-8.17	116.10	121.00
1	C	408	ARG	NE-CZ-NH1	8.07	124.33	120.30
1	B	273	ARG	NE-CZ-NH1	7.76	124.18	120.30
1	A	1019	ARG	NE-CZ-NH1	7.67	124.13	120.30
1	A	273	ARG	NE-CZ-NH1	7.54	124.07	120.30
1	C	1019	ARG	NE-CZ-NH1	7.46	124.03	120.30
1	A	1014	ARG	NE-CZ-NH1	7.19	123.89	120.30
1	A	328	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	C	1014	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	C	273	ARG	NE-CZ-NH1	7.09	123.84	120.30
1	A	319	ARG	NE-CZ-NH1	7.04	123.82	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	328	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	B	983	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	C	403	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	B	1014	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	C	815	ARG	NE-CZ-NH1	6.31	123.46	120.30
1	C	995	ARG	NE-CZ-NH1	6.31	123.46	120.30
1	A	403	ARG	NE-CZ-NH1	6.31	123.45	120.30
1	B	815	ARG	NE-CZ-NH1	6.23	123.42	120.30
1	B	1019	ARG	NE-CZ-NH1	6.19	123.39	120.30
1	B	995	ARG	NE-CZ-NH1	6.17	123.39	120.30
1	B	346	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	C	983	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	A	983	ARG	NE-CZ-NH1	6.09	123.34	120.30
1	A	815	ARG	NE-CZ-NH1	6.08	123.34	120.30
1	A	451	TYR	CB-CG-CD1	-6.06	117.36	121.00
1	B	466	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	B	355	ARG	NE-CZ-NH1	6.02	123.31	120.30
1	A	995	ARG	NE-CZ-NH1	6.02	123.31	120.30
1	C	34	ARG	NE-CZ-NH1	6.01	123.31	120.30
1	A	466	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	A	403	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	B	328	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	C	567	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	C	346	ARG	NE-CZ-NH1	5.78	123.19	120.30
1	A	346	ARG	NE-CZ-NH1	5.75	123.18	120.30
1	A	454	ARG	NE-CZ-NH1	5.75	123.18	120.30
1	B	403	ARG	NE-CZ-NH1	5.74	123.17	120.30
1	C	214	ARG	NE-CZ-NH1	5.71	123.15	120.30
1	B	451	TYR	CB-CG-CD1	-5.60	117.64	121.00
1	C	355	ARG	NE-CZ-NH1	5.58	123.09	120.30
1	C	237	ARG	NE-CZ-NH1	5.57	123.09	120.30
1	A	357	ARG	NE-CZ-NH1	5.53	123.06	120.30
1	B	319	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	B	34	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	B	102	ARG	NE-CZ-NH1	5.41	123.00	120.30
1	A	237	ARG	NE-CZ-NH1	5.39	123.00	120.30
1	C	319	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	C	815	ARG	NE-CZ-NH2	-5.33	117.64	120.30
1	B	567	ARG	NE-CZ-NH1	5.30	122.95	120.30
1	B	815	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	815	ARG	NE-CZ-NH2	-5.27	117.67	120.30
1	B	391	CYS	CA-CB-SG	5.26	123.47	114.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	454	ARG	NE-CZ-NH1	5.23	122.91	120.30
1	A	328	ARG	NE-CZ-NH2	-5.17	117.72	120.30
1	A	34	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	B	1000	ARG	NE-CZ-NH1	5.14	122.87	120.30
1	A	905	ARG	NE-CZ-NH1	5.11	122.86	120.30
1	C	1000	ARG	NE-CZ-NH1	5.09	122.84	120.30
1	A	355	ARG	NE-CZ-NH1	5.02	122.81	120.30
1	B	357	ARG	NE-CZ-NH1	5.01	122.81	120.30
1	A	214	ARG	NE-CZ-NH1	5.01	122.80	120.30
1	B	214	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	983	ARG	Sidechain
1	B	355	ARG	Sidechain
1	B	983	ARG	Sidechain
1	C	269	TYR	Sidechain
1	C	319	ARG	Sidechain
1	C	355	ARG	Sidechain
1	C	396	TYR	Sidechain
1	C	983	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7602	0	7429	2	0
1	B	7602	0	7429	1	0
1	C	7602	0	7429	2	0
All	All	22806	0	22287	5	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (5) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:726:ILE:HG22	1:A:948:LEU:HD13	1.97	0.46
1:C:350:VAL:HG23	1:C:400:PHE:CD2	2.52	0.45
1:B:431:GLY:HA2	1:B:515:PHE:CD2	2.53	0.43
1:A:431:GLY:HA2	1:A:515:PHE:CD2	2.56	0.41
1:C:410:ILE:N	1:C:410:ILE:HD12	2.36	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	948/1208 (78%)	894 (94%)	53 (6%)	1 (0%)	48	77
1	B	948/1208 (78%)	899 (95%)	48 (5%)	1 (0%)	48	77
1	C	948/1208 (78%)	899 (95%)	49 (5%)	0	100	100
All	All	2844/3624 (78%)	2692 (95%)	150 (5%)	2 (0%)	50	77

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	614	ASP
1	A	463	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	852/1053 (81%)	846 (99%)	6 (1%)	81	89
1	B	852/1053 (81%)	848 (100%)	4 (0%)	86	91
1	C	852/1053 (81%)	846 (99%)	6 (1%)	81	89
All	All	2556/3159 (81%)	2540 (99%)	16 (1%)	82	90

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	366	SER
1	A	493	GLN
1	A	721	SER
1	A	760	CYS
1	A	902	MET
1	A	916	LEU
1	B	366	SER
1	B	574	ASP
1	B	586	ASP
1	B	721	SER
1	C	619	GLU
1	C	721	SER
1	C	760	CYS
1	C	902	MET
1	C	916	LEU
1	C	1082	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-25986. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 144



Y Index: 154




Z Index: 116


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.432. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

## 6.6 Mask visualisation [i](#)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 137 nm<sup>3</sup>; this corresponds to an approximate mass of 124 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.327 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-25986 and PDB model 7TLB. Per-residue inclusion information can be found in [section 3](#) on [page 4](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.432 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.432).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 82% of all backbone atoms, 73% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.432) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7270	<div></div> 0.4530
A	<div></div> 0.7220	<div></div> 0.4490
B	<div></div> 0.7390	<div></div> 0.4600
C	<div></div> 0.7190	<div></div> 0.4500

