



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2024 – 12:33 PM EDT

PDB ID : 2TPI  
Title : ON THE DISORDERED ACTIVATION DOMAIN IN TRYPSINOGEN.  
CHEMICAL LABELLING AND LOW-TEMPERATURE CRYSTALLOG-  
RAPHY  
Authors : Walter, J.; Steigemann, W.; Singh, T.P.; Bartunik, H.; Bode, W.; Huber, R.  
Deposited on : 1981-10-26  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Z	229	<div> <div></div> <div>68%</div> <div>24%</div> <div>• •</div> </div>
2	I	58	<div> <div>74%</div> <div>14%</div> <div>7%</div> <div>• •</div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2214 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRYPSINOGEN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	Z	220	1618	1003	276	321	18	65	2	0

- Molecule 2 is a protein called TRYPSIN INHIBITOR.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	I	57	438	275	80	76	7	17	0	1

- Molecule 3 is ISOLEUCINE (three-letter code: ILE) (formula:  $C_6H_{13}NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	Z	1	8	6	1	1	0	0

- Molecule 4 is VALINE (three-letter code: VAL) (formula:  $C_5H_{11}NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	Z	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 5 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	Z	1	Total	Hg	0	1
			3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	Z	105	Total	O	0	0
			105	105		
6	I	34	Total	O	0	0
			34	34		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: TRYPSINOGEN



#### • Molecule 2: TRYPSIN INHIBITOR



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.67Å 84.16Å 122.89Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.50 – 2.10 36.02 – 2.00	Depositor EDS
% Data completeness (in resolution range)	(Not available) (6.50-2.10) 54.3 (36.02-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	unknown	Depositor
R, $R_{free}$	0.200 , (Not available) 0.206 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtriage
Anisotropy	0.393	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 61.7	EDS
L-test for twinning <sup>1</sup>	$\langle  L  \rangle = 0.38$ , $\langle L^2 \rangle = 0.21$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	2214	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	Z	1.06	5/1665 (0.3%)	1.09	1/2256 (0.0%)
2	I	0.96	0/449	1.16	1/602 (0.2%)
All	All	1.04	5/2114 (0.2%)	1.10	2/2858 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	Z	0	17
2	I	0	5
All	All	0	22

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	Z	51	TRP	NE1-CE2	-8.23	1.26	1.37
1	Z	237	TRP	NE1-CE2	-7.71	1.27	1.37
1	Z	215	TRP	NE1-CE2	-7.43	1.27	1.37
1	Z	141	TRP	NE1-CE2	-6.95	1.28	1.37
1	Z	186	GLU	CD-OE2	5.89	1.32	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	53	ARG	NE-CZ-NH2	-9.24	115.68	120.30
1	Z	165	ASP	CB-CG-OD2	-8.19	110.93	118.30

There are no chirality outliers.

All (22) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	2	PRO	Mainchain
2	I	3	ASP	Sidechain
2	I	31	GLN	Sidechain
2	I	43	ASN	Sidechain
2	I	44	ASN	Sidechain
1	Z	147	SER	Mainchain
1	Z	165	ASP	Sidechain
1	Z	175	GLN	Sidechain
1	Z	186	GLU	Sidechain
1	Z	192	GLN	Sidechain
1	Z	194	ASP	Sidechain
1	Z	198	PRO	Mainchain
1	Z	210	GLN	Sidechain
1	Z	223	ASN	Sidechain
1	Z	233	ASN	Mainchain
1	Z	245	ASN	Sidechain
1	Z	27	VAL	Mainchain
1	Z	30	GLN	Sidechain
1	Z	70	GLU	Mainchain
1	Z	79	ASN	Sidechain
1	Z	80	GLU	Sidechain
1	Z	95	ASN	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Z	1618	0	1569	26	2
2	I	438	0	418	7	3
3	Z	8	0	10	0	0
4	Z	8	0	9	0	0
5	Z	3	0	0	0	0
6	I	34	0	0	0	2
6	Z	105	0	0	0	0
All	All	2214	0	2006	33	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 8.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:Z:143:ASN:O	1:Z:191[C]:CYS:SG	2.02	1.17
1:Z:143:ASN:HB3	1:Z:191[A]:CYS:SG	1.89	1.12
1:Z:143:ASN:HB3	1:Z:191[B]:CYS:SG	2.14	0.88
1:Z:48:ASN:HD22	1:Z:50:GLN:H	1.43	0.64
1:Z:137:LEU:HD11	1:Z:157:CYS:HB3	1.82	0.62
2:I:44:ASN:C	2:I:44:ASN:HD22	2.04	0.59
1:Z:100:ASN:HD21	1:Z:179:ASN:HD22	1.52	0.57
1:Z:190:SER:O	1:Z:220[C]:CYS:SG	2.62	0.57
1:Z:50:GLN:HG2	1:Z:111:ALA:HA	1.89	0.54
1:Z:189:ASP:HB3	1:Z:220[C]:CYS:SG	2.48	0.53
1:Z:238:ILE:O	1:Z:242:ILE:HD13	2.11	0.51
2:I:23:TYR:H	2:I:43:ASN:HD21	1.59	0.50
1:Z:219:GLY:C	1:Z:220[A]:CYS:SG	2.89	0.50
1:Z:199:VAL:HG21	1:Z:228:TYR:CD1	2.49	0.48
1:Z:64:GLN:NE2	1:Z:66:ARG:HH21	2.11	0.48
2:I:20:ARG:HE	2:I:44:ASN:HD21	1.62	0.48
1:Z:87:LYS:HE3	1:Z:245:ASN:HB3	1.96	0.47
1:Z:30:GLN:HG2	1:Z:155:LEU:HD11	2.00	0.44
2:I:50:ASP:O	2:I:54:THR:HG23	2.18	0.43
2:I:23:TYR:H	2:I:43:ASN:ND2	2.16	0.43
1:Z:64:GLN:HE21	1:Z:66:ARG:HE	1.67	0.43
2:I:17:ARG:HH11	2:I:17:ARG:HD2	1.63	0.42
2:I:44:ASN:C	2:I:44:ASN:ND2	2.72	0.42
1:Z:123:LEU:HD23	1:Z:235:VAL:HG13	2.01	0.42
1:Z:56:ALA:HB1	1:Z:90:VAL:HG13	2.02	0.42
1:Z:105:LEU:HD13	1:Z:242:ILE:CD1	2.50	0.42
1:Z:105:LEU:HD22	1:Z:242:ILE:HD11	2.01	0.42
1:Z:87:LYS:HB2	1:Z:107:LYS:HB3	2.02	0.41
1:Z:23:GLY:O	1:Z:26:THR:HB	2.20	0.41
1:Z:29:TYR:CE1	1:Z:200:VAL:HG21	2.56	0.40
1:Z:48:ASN:ND2	1:Z:50:GLN:H	2.16	0.40
1:Z:49:SER:HA	1:Z:114:LEU:HD11	2.03	0.40
1:Z:211:GLY:HA2	1:Z:229:THR:O	2.21	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:I:589:HOH:O	6:I:589:HOH:O[4_575]	0.83	1.37
1:Z:74:ASN:ND2	2:I:58:ALA:N[4_575]	1.44	0.76
1:Z:109:LYS:NZ	2:I:54:THR:O[2_575]	1.57	0.63
2:I:31:GLN:CG	6:I:589:HOH:O[4_575]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Z	222/229 (97%)	212 (96%)	10 (4%)	0	100	100
2	I	55/58 (95%)	51 (93%)	4 (7%)	0	100	100
All	All	277/287 (96%)	263 (95%)	14 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Z	186/190 (98%)	172 (92%)	14 (8%)	13	10
2	I	45/46 (98%)	38 (84%)	7 (16%)	2	1
All	All	231/236 (98%)	210 (91%)	21 (9%)	10	6

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Z	26	THR
1	Z	28	PRO
1	Z	48	ASN
1	Z	75	VAL
1	Z	99	LEU
1	Z	113	SER
1	Z	155	LEU
1	Z	163	LEU
1	Z	170	SER
1	Z	192	GLN
1	Z	220[A]	CYS
1	Z	220[B]	CYS
1	Z	220[C]	CYS
1	Z	242	ILE
2	I	17	ARG
2	I	20	ARG
2	I	29	LEU
2	I	43	ASN
2	I	44	ASN
2	I	49	GLU
2	I	54	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	Z	30	GLN
1	Z	48	ASN
1	Z	64	GLN
1	Z	100	ASN
1	Z	101	ASN
1	Z	192	GLN
2	I	43	ASN
2	I	44	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ILE	Z	1016	4	6,7,8	0.81	0	5,8,10	1.99	2 (40%)
4	VAL	Z	1017	3	5,7,7	0.93	0	7,9,9	1.21	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ILE	Z	1016	4	-	1/7/8/10	-
4	VAL	Z	1017	3	-	4/8/8/8	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	Z	1016	ILE	CB-CA-C	-3.27	107.83	112.83
4	Z	1017	VAL	OXT-C-O	-2.85	117.61	124.09
3	Z	1016	ILE	CG1-CB-CA	2.03	116.28	111.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	Z	1017	VAL	N-CA-CB-CG1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
4	Z	1017	VAL	C-CA-CB-CG1
4	Z	1017	VAL	C-CA-CB-CG2
4	Z	1017	VAL	OXT-C-CA-N
3	Z	1016	ILE	C-CA-CB-CG1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	Z	220/229 (96%)	-0.18	3 (1%) 75 78	11, 19, 26, 32	36 (16%)
2	I	56/58 (96%)	-0.41	0 100 100	12, 18, 24, 30	7 (12%)
All	All	276/287 (96%)	-0.23	3 (1%) 80 84	11, 19, 26, 32	43 (15%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Z	220[A]	CYS	3.3
1	Z	146	SER	2.9
1	Z	114	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	VAL	Z	1017	8/8	0.93	0.12	15,20,20,23	0
3	ILE	Z	1016	8/9	0.95	0.13	17,20,20,20	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	HG	Z	1[A]	1/1	0.95	0.11	85,85,85,85	1
5	HG	Z	1[B]	1/1	0.95	0.11	49,49,49,49	1
5	HG	Z	1[C]	1/1	0.95	0.11	83,83,83,83	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.