



wwPDB EM Validation Summary Report ⓘ

Feb 20, 2025 – 07:47 AM EST

PDB ID : 1TUB
Title : TUBULIN ALPHA-BETA DIMER, ELECTRON DIFFRACTION
Authors : Nogales, E.; Downing, K.H.
Deposited on : 1997-09-23
Resolution : 3.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON CRYSTALLOGRAPHY

The reported resolution of this entry is 3.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	440	
2	B	427	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GTP	A	500	-	-	X	-
4	GDP	B	500	-	-	X	-
5	TXL	B	501	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TUBULIN.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	440	3430	2168	583	657	22	0	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	GLY	ALA	conflict	UNP P02550

- Molecule 2 is a protein called TUBULIN.


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	427	3359	2110	576	647	26	0	0

- Molecule 3 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					AltConf
3	A	1	Total	C	N	O	P	0
			32	10	5	14	3	

- Molecule 4 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
5	B	1	58	43	1	14	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TUBULIN



• Molecule 2: TUBULIN



Y432	Y433	Y434	Y435	Y436	Y437	K372	K373	K374	K375	K376	K377	K378	K379	K380	K381	K382	K383	K384	K385	K386	K387	K388	K389	K390	K391	K392	K393	K394	K395	K396	K397	K398	K399	F399	F400	F401	F402	F403	F404	F405	F406	F407	F408	F409	F410	F411	F412	F413	F414	F415	F416	F417	F418	F419	F420	F421	F422	F423	F424	F425	F426	F427	F428	F429	F430	F431																																																																																																																									
A304	A305	A306	A307	A308	A309	A310	A311	A312	A313	A314	A315	A316	A317	A318	A319	A320	A321	A322	A323	A324	A325	A326	A327	A328	A329	A330	A331	A332	A333	A334	A335	A336	A337	A338	A339	A340	A341	A342	A343	A344	A345	A346	A347	A348	A349	A350	A351	A352	A353	A354	A355	A356	A357	A358	A359	A360	A361	A362	A363	A364	A365	A366	A367	A368	A369	A370	A371																																																																																																																								
P184	P185	P186	P187	P188	P189	P190	P191	P192	P193	P194	P195	P196	P197	P198	P199	P200	P201	P202	P203	P204	P205	P206	P207	P208	P209	P210	P211	P212	P213	P214	P215	P216	P217	P218	P219	P220	P221	P222	P223	P224	P225	P226	P227	P228	P229	P230	P231	P232	P233	P234	P235	P236	P237	P238	P239	P240	P241	P242	P243	P244	P245	P246	P247	P248	P249	P250	P251	P252	P253	P254	P255	P256	P257	P258	P259	P260	P261	P262	P263	P264	P265	P266	P267	P268	P269	P270	P271	P272	P273	P274	P275	P276	P277	P278	P279	P280	P281	P282	P283	P284	P285	P286	P287	P288	P289	P290	P291	P292	P293	P294	P295	P296	P297	P298	P299	P300	P301	P302	P303	P304	P305	P306	P307	P308	P309	P310	P311	P312	P313	P314	P315	P316	P317	P318	P319	P320	P321	P322	P323	P324	P325	P326	P327	P328	P329	P330	P331	P332	P333	P334	P335	P336	P337	P338	P339	P340	P341	P342	P343	P344	P345	P346	P347	P348	P349	P350	P351	P352	P353	P354	P355	P356	P357	P358	P359	P360	P361	P362	P363	P364	P365	P366	P367	P368	P369	P370	P371

4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.00Å 92.00Å 90.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 3.70	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-3.70)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Wilson B-factor (Å ²)	144.0	Xtriage
Anisotropy	0.504	Xtriage
L-test for twinning ¹	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	0.000 for -h,-l,-k 0.000 for -h,l,k 0.006 for h,-k,-l	Xtriage
Total number of atoms	6907	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 61.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2123e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TXL, GDP, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	2.34	110/3508 (3.1%)	2.76	214/4762 (4.5%)
2	B	2.47	111/3434 (3.2%)	3.07	266/4652 (5.7%)
All	All	2.41	221/6942 (3.2%)	2.92	480/9414 (5.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	56
2	B	0	59
All	All	0	115

The worst 5 of 221 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	278	ARG	CA-CB	34.96	2.30	1.53
2	B	105	LYS	C-N	-29.38	0.80	1.33
2	B	73	GLY	C-N	-28.00	0.69	1.34
1	A	38	SER	C-N	-27.48	0.70	1.34
1	A	347	CYS	C-N	-23.43	0.89	1.34

The worst 5 of 480 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	368	LEU	O-C-N	-54.19	36.00	122.70
1	A	363	VAL	C-N-CD	-48.68	13.51	120.60
2	B	273	ALA	C-N-CD	-46.48	18.35	120.60
2	B	105	LYS	O-C-N	-44.51	47.53	123.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	88	ARG	C-N-CD	-44.07	23.65	120.60

There are no chirality outliers.

5 of 115 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	18	ASN	Mainchain
1	A	24	TYR	Mainchain,Peptide
1	A	30	ILE	Mainchain,Peptide
1	A	34	GLY	Peptide
1	A	36	MET	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3430	0	3257	1657	34
2	B	3359	0	3172	1921	15
3	A	32	0	11	30	0
4	B	28	0	12	17	0
5	B	58	0	51	59	0
All	All	6907	0	6503	3480	35

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 260.

The worst 5 of 3480 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:229:HIS:CE1	5:B:501:TXL:H343	1.28	1.68
2:B:346:TRP:CE3	2:B:347:ILE:HG13	1.25	1.66
1:A:212:ILE:HD11	1:A:230:LEU:CD2	1.25	1.65
2:B:151:THR:CB	2:B:192:HIS:CD2	1.75	1.62
1:A:115:ILE:CD1	1:A:152:LEU:HG	1.15	1.62

The worst 5 of 35 symmetry-related close contacts are listed below. The label for Atom-2 includes

the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:HIS:CG	1:A:420:GLU:CD[2_444]	0.73	1.47
1:A:1:MET:CE	2:B:72:PRO:CG[1_655]	1.00	1.20
1:A:283:HIS:CG	1:A:420:GLU:OE1[2_444]	1.01	1.19
1:A:283:HIS:ND1	1:A:420:GLU:CD[2_444]	1.11	1.09
1:A:283:HIS:CD2	1:A:420:GLU:CA[2_444]	1.18	1.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/440 (100%)	323 (74%)	62 (14%)	53 (12%)	0	4
2	B	425/427 (100%)	297 (70%)	49 (12%)	79 (19%)	0	1
All	All	863/867 (100%)	620 (72%)	111 (13%)	132 (15%)	0	3

5 of 132 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	31	GLN
1	A	33	ASP
1	A	35	GLN
1	A	39	ASP
1	A	42	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/369 (100%)	351 (95%)	18 (5%)	21	47
2	B	368/368 (100%)	348 (95%)	20 (5%)	18	46
All	All	737/737 (100%)	699 (95%)	38 (5%)	22	46

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	300	ASN
2	B	426	ASN
2	B	302	MET
2	B	390	ARG
2	B	436	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 34 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	336	GLN
2	B	337	ASN
2	B	406	HIS
1	A	380	ASN
1	A	356	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GDP	B	500	-	25,30,30	1.59	5 (20%)	30,47,47	1.06	3 (10%)
5	TXL	B	501	-	63,63,63	4.48	48 (76%)	100,100,100	3.47	55 (55%)
3	GTP	A	500	-	29,34,34	2.32	9 (31%)	35,54,54	1.77	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GDP	B	500	-	-	4/12/32/32	0/3/3/3
5	TXL	B	501	-	-	0/38/124/124	0/6/6/6
3	GTP	A	500	-	-	1/18/38/38	0/3/3/3

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	501	TXL	O3-C4	-10.05	1.26	1.46
5	B	501	TXL	C41-C36	9.35	1.53	1.39
5	B	501	TXL	C37-C36	-8.74	1.26	1.39
5	B	501	TXL	C8-C3	-8.13	1.38	1.57
5	B	501	TXL	O6-C9	-8.05	1.08	1.21

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	501	TXL	C39-C38-C37	11.06	133.89	120.24
5	B	501	TXL	C15-C1-C2	10.89	123.89	111.93
5	B	501	TXL	C38-C37-C36	-9.75	110.78	120.36
5	B	501	TXL	O7-C10-C9	7.21	121.53	109.51
5	B	501	TXL	C39-C40-C41	-6.27	112.51	120.24

There are no chirality outliers.

All (5) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	B	500	GDP	PA-O3A-PB-O2B
4	B	500	GDP	C5'-O5'-PA-O1A
4	B	500	GDP	PA-O3A-PB-O1B
4	B	500	GDP	PA-O3A-PB-O3B
3	A	500	GTP	PA-O3A-PB-O2B


There are no ring outliers.

3 monomers are involved in 106 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	500	GDP	17	0
5	B	501	TXL	59	0
3	A	500	GTP	30	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	52
1	A	41

The worst 5 of 93 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	387:LEU	C	388:PHE	N	1.63
1	A	283:HIS	C	284:GLU	N	1.20
1	A	380:ASN	C	381:THR	N	1.20
1	A	437:VAL	C	438:ASP	N	1.20
1	B	52:TYR	C	53:TYR	N	1.20