



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 04:49 PM EDT

PDB ID : 6TVV
Title : Crystal structure of 3'-5' RecJ exonuclease from M. Jannaschii
Authors : De March, M.; Medagli, B.; Krastanova, I.; Saha, I.; Pisani, F.; Onesti, S.
Deposited on : 2020-01-10
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	
1	B	432	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6373 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called MjaRecJ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	429	Total	C	N	O	S	0	0	0
			3279	2129	541	595	14			
1	B	428	Total	C	N	O	S	0	0	0
			3059	1972	516	558	13			

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		
2	B	2	Total	Mn	0	0
			2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	11	Total	O	0	0
			11	11		
4	B	2	Total	O	0	0
			2	2		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: MjaRecJ

Chain A: 



• Molecule 1: MjaRecJ

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	92.07Å 92.07Å 275.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	87.33 – 2.80 87.33 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (87.33-2.80) 99.8 (87.33-2.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.78 (at 2.82Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.214 , 0.264 0.214 , 0.263	Depositor DCC
R_{free} test set	1476 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	62.9	Xtriage
Anisotropy	0.364	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6373	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/3338	0.88	2/4518 (0.0%)
1	B	0.72	0/3114	0.84	0/4244
All	All	0.72	0/6452	0.86	2/8762 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	84	ASP	CB-CA-C	6.39	123.18	110.40
1	A	383	ASN	CB-CA-C	6.35	123.10	110.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	ASN	Peptide
1	A	398	GLY	Peptide
1	B	168	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3279	0	3257	46	0
1	B	3059	0	2791	24	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	18	0	24	0	0
4	A	11	0	0	0	0
4	B	2	0	0	0	0
All	All	6373	0	6072	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:VAL:O	1:A:130:ASP:CB	2.32	0.77
1:A:383:ASN:HD21	1:A:386:VAL:HG23	1.49	0.76
1:B:1:MET:O	1:B:1:MET:HG3	1.95	0.66
1:A:383:ASN:HD21	1:A:386:VAL:CG2	2.10	0.64
1:B:260:ASN:HD22	1:B:260:ASN:N	1.95	0.64
1:A:280:GLU:OE2	1:A:316:LYS:HE2	2.00	0.62
1:A:117:GLU:O	1:A:118:ASN:ND2	2.33	0.61
1:A:115:ILE:O	1:A:116:ASN:CG	2.39	0.61
1:B:75:VAL:O	1:B:75:VAL:HG23	2.01	0.60
1:B:361:HIS:O	1:B:361:HIS:CG	2.54	0.60
1:A:129:VAL:O	1:A:130:ASP:HB2	2.03	0.59
1:A:426:GLY:O	1:A:428:GLN:N	2.34	0.59
1:B:1:MET:O	1:B:1:MET:CG	2.52	0.58
1:A:75:VAL:HG23	1:A:75:VAL:O	2.04	0.58
1:B:383:ASN:ND2	1:B:427:GLU:O	2.38	0.56
1:A:129:VAL:O	1:A:130:ASP:HB3	2.07	0.55
1:A:17:LYS:HE3	1:A:118:ASN:HA	1.88	0.54
1:A:252:ILE:HD12	1:A:258:ILE:HG12	1.89	0.54
1:B:259:GLU:HG2	1:B:260:ASN:HD22	1.74	0.53
1:A:84:ASP:HB2	1:A:137:ALA:CB	2.38	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:383:ASN:ND2	1:A:386:VAL:HG23	2.20	0.53
1:A:90:ILE:HD11	1:A:115:ILE:CD1	2.39	0.52
1:B:392:LYS:HA	1:B:396:GLY:O	2.11	0.51
1:A:426:GLY:C	1:A:428:GLN:H	2.14	0.50
1:B:361:HIS:O	1:B:363:GLU:N	2.44	0.50
1:A:360:TYR:HA	1:A:368:LYS:O	2.12	0.50
1:A:400:GLY:O	1:A:405:SER:HB3	2.12	0.49
1:A:7:ILE:HD13	1:A:127:PHE:CD2	2.47	0.49
1:B:130:ASP:OD2	1:B:132:SER:OG	2.31	0.49
1:A:83:ALA:O	1:A:103:LEU:O	2.30	0.49
1:A:397:ASN:N	1:A:397:ASN:ND2	2.61	0.49
1:A:3:LYS:O	1:A:7:ILE:HG12	2.13	0.48
1:B:360:TYR:HA	1:B:368:LYS:O	2.13	0.48
1:A:118:ASN:C	1:A:118:ASN:HD22	2.16	0.48
1:A:377:LEU:HB3	1:A:382:LEU:HD12	1.96	0.47
1:A:388:MET:HG3	1:A:405:SER:OG	2.15	0.46
1:B:248:LEU:HA	1:B:251:ILE:HG22	1.97	0.46
1:B:371:ALA:H	1:B:405:SER:CB	2.28	0.46
1:A:208:TYR:HA	1:A:320:ILE:HG13	1.98	0.46
1:A:397:ASN:N	1:A:397:ASN:HD22	2.14	0.46
1:A:377:LEU:O	1:A:382:LEU:HB2	2.16	0.45
1:B:83:ALA:O	1:B:103:LEU:O	2.34	0.45
1:A:248:LEU:HA	1:A:251:ILE:HG22	1.98	0.44
1:A:401:HIS:HB2	1:A:404:ALA:O	2.17	0.44
1:A:400:GLY:O	1:A:405:SER:CB	2.65	0.44
1:B:397:ASN:O	1:B:407:ALA:O	2.36	0.44
1:A:90:ILE:HD11	1:A:115:ILE:HD13	1.99	0.44
1:B:136:THR:O	1:B:140:VAL:HG23	2.18	0.44
1:B:377:LEU:HD12	1:B:377:LEU:HA	1.81	0.43
1:A:366:ILE:CD1	1:A:408:VAL:CG1	2.98	0.42
1:B:210:THR:OG1	1:B:350:ILE:HD13	2.19	0.42
1:B:370:SER:HA	1:B:405:SER:CB	2.50	0.42
1:B:259:GLU:HG2	1:B:260:ASN:ND2	2.34	0.42
1:A:383:ASN:ND2	1:A:386:VAL:CG2	2.79	0.42
1:A:7:ILE:HG21	1:A:143:LEU:HD13	2.02	0.42
1:A:136:THR:O	1:A:140:VAL:HG23	2.19	0.41
1:A:87:SER:OG	1:A:121:GLN:NE2	2.50	0.41
1:B:111:LYS:HA	1:B:126:ILE:HG12	2.03	0.41
1:A:7:ILE:CD1	1:A:127:PHE:CD2	3.03	0.41
1:A:338:GLU:HB2	1:A:360:TYR:CZ	2.55	0.41
1:A:198:TYR:CE1	1:A:316:LYS:HE3	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:ILE:HD11	1:B:115:ILE:CD1	2.51	0.41
1:A:111:LYS:HA	1:A:126:ILE:HG12	2.03	0.41
1:B:212:PRO:CD	1:B:261:LEU:HD23	2.50	0.41
1:A:168:ASN:HA	1:A:169:PRO:HA	1.83	0.40
1:A:75:VAL:O	1:A:75:VAL:CG2	2.69	0.40
1:A:138:SER:HB2	1:A:157:ALA:HA	2.02	0.40
1:A:80:PHE:CD1	1:A:98:PHE:HD1	2.39	0.40
1:B:208:TYR:HA	1:B:320:ILE:HG13	2.03	0.40
1:A:399:GLY:N	1:A:406:GLY:O	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/432 (99%)	406 (95%)	15 (4%)	6 (1%)	11	34
1	B	426/432 (99%)	400 (94%)	21 (5%)	5 (1%)	13	39
All	All	853/864 (99%)	806 (94%)	36 (4%)	11 (1%)	12	36

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	130	ASP
1	A	427	GLU
1	B	76	ASN
1	B	362	ILE
1	A	76	ASN
1	A	332	ASN
1	A	401	HIS
1	B	332	ASN
1	B	169	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	408	VAL
1	A	425	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/373 (89%)	302 (91%)	31 (9%)	9	26
1	B	266/373 (71%)	251 (94%)	15 (6%)	21	51
All	All	599/746 (80%)	553 (92%)	46 (8%)	13	35

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	A	42	LEU
1	A	59	GLU
1	A	60	HIS
1	A	69	LEU
1	A	84	ASP
1	A	112	ASP
1	A	118	ASN
1	A	165	MET
1	A	185	ARG
1	A	228	LYS
1	A	232	ILE
1	A	236	LYS
1	A	258	ILE
1	A	263	ILE
1	A	282	LEU
1	A	309	ASN
1	A	334	ILE
1	A	346	ILE
1	A	349	SER
1	A	366	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	377	LEU
1	A	382	LEU
1	A	383	ASN
1	A	384	LEU
1	A	385	SER
1	A	390	VAL
1	A	397	ASN
1	A	410	SER
1	A	416	GLU
1	A	425	ILE
1	B	0	MET
1	B	1	MET
1	B	42	LEU
1	B	165	MET
1	B	185	ARG
1	B	260	ASN
1	B	282	LEU
1	B	309	ASN
1	B	310	GLN
1	B	333	ASN
1	B	349	SER
1	B	372	ARG
1	B	375	ARG
1	B	377	LEU
1	B	383	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	HIS
1	A	118	ASN
1	A	121	GLN
1	A	235	ASN
1	A	238	GLN
1	A	288	ASN
1	A	309	ASN
1	A	383	ASN
1	A	397	ASN
1	B	121	GLN
1	B	235	ASN
1	B	238	GLN
1	B	260	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	288	ASN
1	B	310	GLN
1	B	361	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 4 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	504	-	5,5,5	0.13	0	5,5,5	0.30	0
3	GOL	A	505	-	5,5,5	0.18	0	5,5,5	0.47	0
3	GOL	A	503	-	5,5,5	0.07	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	504	-	-	3/4/4/4	-
3	GOL	A	505	-	-	2/4/4/4	-
3	GOL	A	503	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	504	GOL	C1-C2-C3-O3
3	A	504	GOL	O2-C2-C3-O3
3	A	505	GOL	O2-C2-C3-O3
3	A	503	GOL	C1-C2-C3-O3
3	A	505	GOL	C1-C2-C3-O3
3	A	503	GOL	O2-C2-C3-O3
3	A	504	GOL	O1-C1-C2-O2
3	A	503	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	429/432 (99%)	-0.02	1 (0%) 95 94	33, 50, 70, 114	23 (5%)
1	B	428/432 (99%)	0.11	7 (1%) 72 66	39, 65, 95, 126	115 (26%)
All	All	857/864 (99%)	0.05	8 (0%) 84 80	33, 56, 89, 126	138 (16%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	403	VAL	4.0
1	B	407	ALA	3.6
1	B	364	GLY	3.3
1	B	367	ALA	3.0
1	B	389	ALA	2.5
1	B	404	ALA	2.1
1	B	426	GLY	2.1
1	A	109	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	505	6/6	0.78	0.27	65,71,75,76	0
3	GOL	A	504	6/6	0.81	0.22	83,87,90,91	5
3	GOL	A	503	6/6	0.86	0.19	80,89,90,93	4
2	MN	A	502	1/1	0.88	0.10	73,73,73,73	1
2	MN	A	501	1/1	0.95	0.04	57,57,57,57	0
2	MN	B	502	1/1	0.96	0.08	54,54,54,54	1
2	MN	B	501	1/1	0.98	0.05	59,59,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MN A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MN B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MN B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.