



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2025 – 08:52 PM EDT

PDB ID : 3U69 / pdb\_00003u69  
Title : Unliganded wild-type human thrombin  
Authors : Figueiredo, A.C.; Clement, C.C.; Philipp, M.; Barbosa Pereira, P.J.  
Deposited on : 2011-10-12  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1935 (1.56-1.56)
Clashscore	180529	2073 (1.56-1.56)
Ramachandran outliers	177936	2037 (1.56-1.56)
Sidechain outliers	177891	2034 (1.56-1.56)
RSRZ outliers	164620	1935 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	30	
2	H	259	

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5272 atoms, of which 2477 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prothrombin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	L	30	533	165	267	45	55	1	0	3	0

- Molecule 2 is a protein called Prothrombin.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	H	251	4269	1360	2156	371	366	16	0	14	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	H	1	14	8	1	5	0	0

- Molecule 4 is BICINE (CCD ID: BCN) (formula:  $C_6H_{13}NO_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	H	1	Total	C	H	N	O	0	0
			23	6	12	1	4		

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	H	1	Total	Na	0	0
			1	1		

- Molecule 6 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	1	Total	I	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	H	1	Total	Cl	0	0
			1	1		

- Molecule 8 is (4R)-2-METHYLPENTANE-2,4-DIOL (CCD ID: MRD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	H	1	Total	C	H	O	0	0
			22	6	14	2		
8	H	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 9 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	H	1	Total	C	H	O	0	0
			22	6	14	2		



- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	L	39	Total 39	O 39	0	0
10	H	325	Total 325	O 325	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Prothrombin



#### • Molecule 2: Prothrombin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.55Å 73.00Å 83.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.30 – 1.55 47.30 – 1.55	Depositor EDS
% Data completeness (in resolution range)	96.5 (47.30-1.55) 97.8 (47.30-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.25 (at 1.55Å)	Xtriage
Refinement program	PHENIX 1.7_650	Depositor
R, $R_{free}$	0.133 , 0.167 0.134 , 0.167	Depositor DCC
$R_{free}$ test set	2542 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.1	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 42.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5272	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MRD, CL, NAG, BCN, IOD, MPD, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.44	0/277	0.76	0/366
2	H	0.50	0/2207	0.72	0/2975
All	All	0.49	0/2484	0.73	0/3341

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	266	267	267	4	0
2	H	2113	2156	2152	36	0
3	H	14	0	13	0	0
4	H	11	12	11	1	0
5	H	1	0	0	0	0
6	H	1	0	0	1	0
7	H	1	0	0	0	0
8	H	16	28	28	8	0
9	H	8	14	14	0	0
10	H	325	0	0	10	0
10	L	39	0	0	1	0
All	All	2795	2477	2485	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:335[B]:GLN:NE2	10:H:108:HOH:O	2.04	0.87
2:H:398:ILE:O	8:H:700:MRD:H5C1	1.76	0.85
8:H:700:MRD:H1C1	8:H:700:MRD:H5C3	1.62	0.82
2:H:498:ARG:HA	2:H:498:ARG:HE	1.47	0.80
2:H:427:LYS:HG2	6:H:2:IOD:I	2.54	0.78
2:H:535:PHE:HB3	8:H:701:MRD:HMC1	1.74	0.70
2:H:380[A]:LEU:HD22	2:H:400[A]:MET:HE1	1.74	0.69
2:H:397:LYS:HG3	10:H:815:HOH:O	1.93	0.69
8:H:701:MRD:H5C3	8:H:701:MRD:H1C1	1.74	0.69
2:H:380[A]:LEU:CD2	2:H:400[A]:MET:HE1	2.23	0.68
2:H:498:ARG:HA	2:H:498:ARG:NE	2.09	0.68
2:H:380[A]:LEU:HD22	2:H:400[A]:MET:CE	2.23	0.67
2:H:494:LYS:CG	2:H:501[B]:ILE:HD11	2.24	0.67
8:H:700:MRD:H1C3	10:H:814:HOH:O	1.92	0.67
8:H:700:MRD:H5C3	8:H:700:MRD:C1	2.26	0.64
1:L:307:LYS:HG2	2:H:328:GLU:OE2	1.98	0.64
2:H:337[B]:MET:SD	2:H:385:LYS:HD3	2.41	0.61
2:H:337[B]:MET:HG3	2:H:461:TRP:CZ3	2.36	0.61
2:H:494:LYS:HG3	2:H:501[B]:ILE:HD11	1.83	0.60
2:H:395:ILE:HD12	10:H:803:HOH:O	2.02	0.60
2:H:494:LYS:HG2	2:H:501[B]:ILE:HD11	1.85	0.58
2:H:380[A]:LEU:CD2	2:H:400[A]:MET:CE	2.81	0.57
2:H:464:LEU:O	2:H:465:LYS:HD3	2.06	0.56
4:H:1500:BCN:H62	10:H:801:HOH:O	2.07	0.55
2:H:337[B]:MET:CG	2:H:461:TRP:CZ3	2.90	0.55
2:H:568:LYS:HB3	10:H:805:HOH:O	2.08	0.53
2:H:380[A]:LEU:CD2	2:H:400[A]:MET:SD	3.00	0.50
2:H:400[A]:MET:HE3	8:H:700:MRD:O4	2.13	0.48
2:H:548:GLY:HA3	10:H:818:HOH:O	2.13	0.47
2:H:337[B]:MET:HE3	2:H:346[B]:LEU:HD13	1.95	0.47
2:H:322:VAL:HG12	2:H:323:GLU:HG2	1.98	0.46
1:L:310[B]:ARG:HD2	10:L:151:HOH:O	2.15	0.46
2:H:494:LYS:HG2	2:H:501[B]:ILE:CD1	2.47	0.45
2:H:412:TRP:CZ3	2:H:413:ARG:HG2	2.52	0.45
2:H:516:LYS:HE3	10:H:659:HOH:O	2.16	0.45
2:H:380[A]:LEU:HD21	2:H:400[A]:MET:HE1	1.96	0.43
1:L:295:LEU:HB2	1:L:301:LYS:HE3	1.99	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:H:701:MRD:HMC2	10:H:255:HOH:O	2.17	0.43
1:L:305[B]:GLU:HG2	1:L:309:GLU:CG	2.49	0.43
2:H:543:GLY:HA2	2:H:561:THR:O	2.19	0.43
2:H:392:GLU:HB3	2:H:395:ILE:HG12	2.02	0.42
2:H:368:PRO:HG2	2:H:412:TRP:CZ2	2.55	0.42
2:H:450[B]:LEU:HA	2:H:450[B]:LEU:HD23	1.78	0.41
2:H:337[B]:MET:HG2	2:H:461:TRP:CZ3	2.57	0.40
2:H:568:LYS:CB	10:H:805:HOH:O	2.68	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	31/30 (103%)	30 (97%)	1 (3%)	0	100	100
2	H	261/259 (101%)	252 (97%)	9 (3%)	0	100	100
All	All	292/289 (101%)	282 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	30/27 (111%)	29 (97%)	1 (3%)	33	8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	233/225 (104%)	230 (99%)	3 (1%)	65	41
All	All	263/252 (104%)	259 (98%)	4 (2%)	58	34

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	301	LYS
2	H	390	ARG
2	H	465	LYS
2	H	498	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	451	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	H	1373	2	14,14,15	0.52	0	17,19,21	0.64	0
4	BCN	H	1500	-	10,10,10	0.84	0	11,11,11	1.71	4 (36%)
9	MPD	H	702	-	7,7,7	0.37	0	9,10,10	0.19	0
8	MRD	H	700	-	7,7,7	0.39	0	9,10,10	0.37	0
8	MRD	H	701	-	7,7,7	0.23	0	9,10,10	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	H	1373	2	-	0/6/23/26	0/1/1/1
4	BCN	H	1500	-	-	2/10/10/10	-
9	MPD	H	702	-	-	1/5/5/5	-
8	MRD	H	700	-	-	3/5/5/5	-
8	MRD	H	701	-	-	3/5/5/5	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1500	BCN	O4-C4-C3	2.83	122.88	111.22
4	H	1500	BCN	C4-C3-N1	2.40	121.84	113.44
4	H	1500	BCN	C5-N1-C3	2.32	116.96	111.44
4	H	1500	BCN	C1-N1-C5	-2.22	106.55	111.91

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	H	700	MRD	C2-C3-C4-O4
8	H	700	MRD	C2-C3-C4-C5
8	H	701	MRD	C2-C3-C4-O4
8	H	701	MRD	C2-C3-C4-C5
4	H	1500	BCN	C6-C5-N1-C3
8	H	700	MRD	O2-C2-C3-C4
8	H	701	MRD	O2-C2-C3-C4
9	H	702	MPD	O2-C2-C3-C4
4	H	1500	BCN	C6-C5-N1-C1

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	H	1500	BCN	1	0
8	H	700	MRD	5	0
8	H	701	MRD	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	30/30 (100%)	-0.13	1 (3%) 49 57	9, 20, 40, 43	3 (10%)
2	H	251/259 (96%)	-0.69	2 (0%) 82 88	6, 17, 37, 53	14 (5%)
All	All	281/289 (97%)	-0.63	3 (1%) 77 84	6, 18, 37, 53	17 (6%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	H	465	LYS	2.4
1	L	291	ALA	2.3
2	H	394	ASN	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	H	1373	14/15	0.64	0.17	49,56,61,61	0
8	MRD	H	700	8/8	0.83	0.18	24,34,69,81	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	MRD	H	701	8/8	0.83	0.15	25,41,59,65	0
9	MPD	H	702	8/8	0.84	0.17	40,56,69,69	0
4	BCN	H	1500	11/11	0.90	0.13	12,24,40,42	23
6	IOD	H	2	1/1	0.97	0.05	27,27,27,27	1
7	CL	H	3	1/1	0.99	0.02	20,20,20,20	0
5	NA	H	1	1/1	0.99	0.02	13,13,13,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.