



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 09:50 PM EDT

PDB ID : 5UDH  
Title : HHARI/ARIH1-UBCH7 Ubiquitin  
Authors : Miller, D.J.; Schulman, B.A.  
Deposited on : 2016-12-27  
Resolution : 3.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1619 (3.28-3.20)
Clashscore	141614	1755 (3.28-3.20)
Ramachandran outliers	138981	1728 (3.28-3.20)
Sidechain outliers	138945	1727 (3.28-3.20)
RSRZ outliers	127900	1567 (3.28-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	470	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>9%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	470	<div> <div>3%</div> <div> <div></div> <div>74%</div> <div>7%</div> <div>•</div> <div>19%</div> </div> </div>
2	C	156	<div> <div></div> <div> <div></div> <div>92%</div> <div>6%</div> <div>•</div> </div> </div>
2	D	156	<div> <div></div> <div> <div></div> <div>95%</div> <div>•</div> <div>•</div> </div> </div>
3	E	82	<div> <div>%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>11%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9178 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase ARIH1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	411	Total	C	N	O	S	0	0	0
			3243	2057	564	581	41			
1	B	381	Total	C	N	O	S	0	0	0
			2977	1879	519	539	40			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	88	GLY	-	expression tag	UNP Q9Y4X5
A	89	SER	-	expression tag	UNP Q9Y4X5
B	88	GLY	-	expression tag	UNP Q9Y4X5
B	89	SER	-	expression tag	UNP Q9Y4X5

- Molecule 2 is a protein called Ubiquitin-conjugating enzyme E2 L3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	152	Total	C	N	O	S	0	0	0
			1212	773	209	225	5			
2	D	155	Total	C	N	O	S	0	0	0
			1204	769	208	222	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	expression tag	UNP P68036
C	0	SER	-	expression tag	UNP P68036
C	86	LYS	CYS	engineered mutation	UNP P68036
D	-1	GLY	-	expression tag	UNP P68036
D	0	SER	-	expression tag	UNP P68036
D	86	LYS	CYS	engineered mutation	UNP P68036

- Molecule 3 is a protein called Ubiquitin C variant.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	73	Total	C	N	O	0	0	0
			530	335	88	107			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-5	HIS	-	expression tag	UNP Q59EM9
E	-4	HIS	-	expression tag	UNP Q59EM9
E	-3	HIS	-	expression tag	UNP Q59EM9
E	-2	HIS	-	expression tag	UNP Q59EM9
E	-1	HIS	-	expression tag	UNP Q59EM9
E	0	HIS	-	expression tag	UNP Q59EM9


- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	Zn	0	0
			6	6		
4	B	6	Total	Zn	0	0
			6	6		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 ubiquitin-protein ligase ARIH1



Chain D: 

95%



● Molecule 3: Ubiquitin C variant

Chain E: 

%

83%

6%

11%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	184.57Å 76.79Å 147.72Å 90.00° 107.33° 90.00°	Depositor
Resolution (Å)	141.01 – 3.24 48.46 – 3.24	Depositor EDS
% Data completeness (in resolution range)	98.9 (141.01-3.24) 99.0 (48.46-3.24)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 3.25Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.222 , 0.279 0.224 , 0.277	Depositor DCC
$R_{free}$ test set	1550 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	118.7	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 80.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9178	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	122.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.47	0/3322	0.69	0/4505
1	B	0.44	0/3047	0.65	0/4138
2	C	0.44	0/1242	0.65	0/1685
2	D	0.46	0/1235	0.69	0/1682
3	E	0.45	0/536	0.58	0/732
All	All	0.46	0/9382	0.66	0/12742

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	1
All	All	0	4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	221	MET	Peptide
1	A	298	LYS	Peptide
1	A	312	HIS	Peptide
1	B	298	LYS	Peptide



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3243	0	2968	22	0
1	B	2977	0	2709	15	0
2	C	1212	0	1182	6	0
2	D	1204	0	1146	3	0
3	E	530	0	503	2	0
4	A	6	0	0	0	0
4	B	6	0	0	0	0
All	All	9178	0	8508	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:476:TYR:OH	1:A:499:GLU:OE1	2.08	0.71
1:A:220:ILE:HD11	1:A:254:VAL:HG13	1.72	0.69
1:A:109:ILE:HG23	1:A:263:THR:HG21	1.75	0.69
1:A:431:GLU:OE1	1:A:469:ARG:NH1	2.31	0.63
1:A:106:ALA:HA	1:A:109:ILE:HG22	1.82	0.61

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/470 (85%)	371 (92%)	30 (8%)	0	100	100
1	B	371/470 (79%)	347 (94%)	24 (6%)	0	100	100
2	C	150/156 (96%)	146 (97%)	4 (3%)	0	100	100
2	D	153/156 (98%)	148 (97%)	5 (3%)	0	100	100
3	E	71/82 (87%)	65 (92%)	6 (8%)	0	100	100
All	All	1146/1334 (86%)	1077 (94%)	69 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/430 (78%)	323 (96%)	12 (4%)	35	66
1	B	309/430 (72%)	299 (97%)	10 (3%)	39	69
2	C	128/139 (92%)	127 (99%)	1 (1%)	81	91
2	D	122/139 (88%)	120 (98%)	2 (2%)	62	82
3	E	54/74 (73%)	53 (98%)	1 (2%)	57	79
All	All	948/1212 (78%)	922 (97%)	26 (3%)	44	73

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	236	CYS
1	B	301	ARG
2	D	151	ARG
1	B	239	LEU
1	B	418	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	358	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	411/470 (87%)	-0.04	6 (1%) 73 64	58, 103, 165, 204	0
1	B	381/470 (81%)	0.01	13 (3%) 45 33	89, 135, 182, 197	0
2	C	152/156 (97%)	-0.06	0 100 100	76, 108, 144, 176	3 (1%)
2	D	155/156 (99%)	-0.11	0 100 100	74, 114, 148, 177	3 (1%)
3	E	73/82 (89%)	-0.38	1 (1%) 75 66	117, 162, 190, 203	0
All	All	1172/1334 (87%)	-0.06	20 (1%) 70 60	58, 119, 177, 204	6 (0%)

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	352	GLU	4.2
1	A	410	ALA	4.0
1	B	222	GLU	3.6
1	A	224	GLY	3.5
1	B	384	SER	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	ZN	A	601	1/1	0.93	0.06	191,191,191,191	0
4	ZN	B	601	1/1	0.95	0.10	185,185,185,185	0
4	ZN	B	602	1/1	0.97	0.09	151,151,151,151	0
4	ZN	B	605	1/1	0.97	0.22	88,88,88,88	0
4	ZN	A	605	1/1	0.98	0.21	96,96,96,96	0
4	ZN	B	603	1/1	0.98	0.20	127,127,127,127	0
4	ZN	B	604	1/1	0.98	0.20	126,126,126,126	0
4	ZN	A	603	1/1	0.98	0.26	98,98,98,98	0
4	ZN	A	606	1/1	0.99	0.20	92,92,92,92	0
4	ZN	A	604	1/1	0.99	0.24	94,94,94,94	0
4	ZN	A	602	1/1	0.99	0.18	106,106,106,106	0
4	ZN	B	606	1/1	0.99	0.20	109,109,109,109	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.