



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 3, 2024 – 08:21 am GMT

PDB ID : 4UXE
Title : Crystal structure of the carboxy-terminal region of the bacteriophage T4 proximal long tail fibre protein gp34, P21 selenomethionine crystal
Authors : Granell, M.; Alvira, S.; Garcia-Doval, C.; Singh, A.K.; van Raaij, M.J.
Deposited on : 2014-08-22
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>6%</div> </div> <div></div> </div>
1	B	410	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>7%</div> </div> <div></div> </div>
1	C	410	<div> <div>5%</div> <div> <div></div> <div>90%</div> <div>7%</div> </div> <div></div> </div>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10810 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LARGE TAIL FIBER PROTEIN P34.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	396	Total	C	N	O	Se	0	3	0
			3042	1901	536	599	6			
1	B	397	Total	C	N	O	Se	0	6	0
			3059	1912	534	605	8			
1	C	396	Total	C	N	O	Se	0	5	0
			3048	1906	534	600	8			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	880	MSE	-	expression tag	UNP P18771
A	881	GLY	-	expression tag	UNP P18771
A	882	SER	-	expression tag	UNP P18771
A	883	SER	-	expression tag	UNP P18771
A	884	HIS	-	expression tag	UNP P18771
A	885	HIS	-	expression tag	UNP P18771
A	886	HIS	-	expression tag	UNP P18771
A	887	HIS	-	expression tag	UNP P18771
A	888	HIS	-	expression tag	UNP P18771
A	889	HIS	-	expression tag	UNP P18771
A	890	SER	-	expression tag	UNP P18771
A	891	GLN	-	expression tag	UNP P18771
A	892	ASP	-	expression tag	UNP P18771
A	893	PRO	-	expression tag	UNP P18771
B	880	MSE	-	expression tag	UNP P18771
B	881	GLY	-	expression tag	UNP P18771
B	882	SER	-	expression tag	UNP P18771
B	883	SER	-	expression tag	UNP P18771
B	884	HIS	-	expression tag	UNP P18771
B	885	HIS	-	expression tag	UNP P18771
B	886	HIS	-	expression tag	UNP P18771
B	887	HIS	-	expression tag	UNP P18771
B	888	HIS	-	expression tag	UNP P18771

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	889	HIS	-	expression tag	UNP P18771
B	890	SER	-	expression tag	UNP P18771
B	891	GLN	-	expression tag	UNP P18771
B	892	ASP	-	expression tag	UNP P18771
B	893	PRO	-	expression tag	UNP P18771
C	880	MSE	-	expression tag	UNP P18771
C	881	GLY	-	expression tag	UNP P18771
C	882	SER	-	expression tag	UNP P18771
C	883	SER	-	expression tag	UNP P18771
C	884	HIS	-	expression tag	UNP P18771
C	885	HIS	-	expression tag	UNP P18771
C	886	HIS	-	expression tag	UNP P18771
C	887	HIS	-	expression tag	UNP P18771
C	888	HIS	-	expression tag	UNP P18771
C	889	HIS	-	expression tag	UNP P18771
C	890	SER	-	expression tag	UNP P18771
C	891	GLN	-	expression tag	UNP P18771
C	892	ASP	-	expression tag	UNP P18771
C	893	PRO	-	expression tag	UNP P18771

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	661	Total	O	0	0
			661	661		
3	B	558	Total	O	0	0
			558	558		
3	C	400	Total	O	0	0
			400	400		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: LARGE TAIL FIBER PROTEIN P34



• Molecule 1: LARGE TAIL FIBER PROTEIN P34



• Molecule 1: LARGE TAIL FIBER PROTEIN P34



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	92.84Å 76.33Å 116.96Å 90.00° 99.34° 90.00°	Depositor
Resolution (Å)	29.62 – 2.00 29.62 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.62-2.00) 99.8 (29.62-2.00)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.94 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.140 , 0.175 0.147 , 0.181	Depositor DCC
R_{free} test set	2205 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.162	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10810	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/3114	0.67	2/4227 (0.0%)
1	B	0.67	0/3142	0.66	0/4264
1	C	0.65	0/3123	0.65	0/4239
All	All	0.66	0/9379	0.66	2/12730 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	937	ARG	NE-CZ-NH1	6.57	123.58	120.30
1	A	937	ARG	NE-CZ-NH2	-5.40	117.60	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3042	0	2950	29	0
1	B	3059	0	2965	39	0
1	C	3048	0	2957	34	0
2	A	12	0	16	0	0
2	B	12	0	16	0	0
2	C	18	0	24	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	661	0	0	7	0
3	B	558	0	0	4	0
3	C	400	0	0	4	0
All	All	10810	0	8928	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1026[A]:MSE:HE1	1:C:1026[A]:MSE:CE	1.55	1.33
1:B:1026[A]:MSE:HE1	1:C:1026[A]:MSE:HE1	1.19	1.18
1:B:1026[A]:MSE:CE	1:C:1026[A]:MSE:CE	2.30	1.09
1:B:1166[A]:MSE:CE	1:B:1242:PRO:HD2	1.86	1.05
1:A:1031[A]:THR:HG22	3:A:3196:HOH:O	1.58	1.01

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/410 (97%)	383 (96%)	13 (3%)	1 (0%)	37	35
1	B	401/410 (98%)	390 (97%)	11 (3%)	0	100	100
1	C	399/410 (97%)	383 (96%)	15 (4%)	1 (0%)	37	35
All	All	1197/1230 (97%)	1156 (97%)	39 (3%)	2 (0%)	44	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	1067	ALA
1	A	1067	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/337 (99%)	328 (98%)	5 (2%)	60	66
1	B	337/337 (100%)	336 (100%)	1 (0%)	91	94
1	C	334/337 (99%)	332 (99%)	2 (1%)	84	88
All	All	1004/1011 (99%)	996 (99%)	8 (1%)	79	84

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	1123	ARG
1	C	901	ILE
1	A	1281	LYS
1	A	1280	ASN
1	B	1123	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	916	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	2290	-	5,5,5	0.25	0	5,5,5	0.63	0
2	GOL	C	2289	-	5,5,5	0.44	0	5,5,5	0.38	0
2	GOL	A	2289	-	5,5,5	0.46	0	5,5,5	0.39	0
2	GOL	B	2291	-	5,5,5	0.36	0	5,5,5	0.56	0
2	GOL	B	2290	-	5,5,5	0.49	0	5,5,5	0.20	0
2	GOL	C	2291	-	5,5,5	0.41	0	5,5,5	0.39	0
2	GOL	A	2290	-	5,5,5	0.46	0	5,5,5	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	2290	-	-	2/4/4/4	-
2	GOL	C	2289	-	-	0/4/4/4	-
2	GOL	A	2289	-	-	0/4/4/4	-
2	GOL	B	2291	-	-	0/4/4/4	-
2	GOL	B	2290	-	-	0/4/4/4	-
2	GOL	C	2291	-	-	0/4/4/4	-
2	GOL	A	2290	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2290	GOL	O1-C1-C2-C3
2	C	2290	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	2290	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	391/410 (95%)	-0.68	20 (5%) 34 32	4, 13, 60, 87	2 (0%)
1	B	392/410 (95%)	-0.75	12 (3%) 51 49	4, 13, 47, 89	3 (0%)
1	C	391/410 (95%)	-0.66	20 (5%) 34 32	4, 13, 53, 94	2 (0%)
All	All	1174/1230 (95%)	-0.70	52 (4%) 39 38	4, 13, 55, 94	7 (0%)

The worst 5 of 52 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	897	ILE	5.6
1	A	897	ILE	5.2
1	B	893	PRO	5.0
1	C	1279	VAL	4.8
1	A	893	PRO	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	A	2289	6/6	0.93	0.08	18,25,26,29	0
2	GOL	A	2290	6/6	0.94	0.07	23,32,34,36	0
2	GOL	B	2291	6/6	0.94	0.09	23,32,35,41	0
2	GOL	C	2290	6/6	0.95	0.07	21,23,27,29	0
2	GOL	C	2289	6/6	0.96	0.08	17,25,27,29	0
2	GOL	B	2290	6/6	0.97	0.05	17,20,25,28	0
2	GOL	C	2291	6/6	0.97	0.07	19,30,34,34	0

6.5 Other polymers [i](#)

There are no such residues in this entry.