



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 25, 2024 – 05:19 PM EST

PDB ID : 2VDR
Title : Integrin AlphaIIbBeta3 Headpiece Bound to a chimeric Fibrinogen Gamma chain peptide, LGGAKQRGDV
Authors : Springer, T.A.; Zhu, J.; Xiao, T.
Deposited on : 2007-10-10
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	452	<div><div>3%</div><div>96%</div><div>.</div></div>
2	B	461	<div><div>22%</div><div>94%</div><div>5%</div><div>.</div></div>
3	C	10	<div><div>40%</div><div>70%</div><div>30%</div></div>
4	H	221	<div><div>6%</div><div>94%</div><div>5%</div><div>.</div></div>
5	L	214	<div><div>%</div><div>94%</div><div>6%</div></div>
6	D	5	<div><div>60%</div><div>40%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
7	E	7	<div><div></div><div>29%</div><div>71%</div></div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	MAN	D	3	X	-	-	-
7	MAN	E	3	X	-	-	-

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 11884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTEGRIN ALPHA-IIB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	452	Total	C	N	O	S	0	9	0
			3539	2250	616	665	8			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	282	GLY	ALA	conflict	UNP P08514

- Molecule 2 is a protein called INTEGRIN BETA-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	455	Total	C	N	O	S	0	10	0
			3580	2236	607	704	33			

- Molecule 3 is a protein called FIBRINOGEN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	7	Total	C	N	O	0	0	0
			54	31	12	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	408	ARG	ALA	engineered mutation	UNP Q53Y18


- Molecule 4 is a protein called MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	219	Total	C	N	O	S	0	3	0
			1678	1064	270	338	6			

- Molecule 5 is a protein called MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	L	214	Total	C	N	O	S	0	9	0
			1686	1051	273	351	11			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	D	5	Total	C	N	O		0	0	0
			61	34	2	25				

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	E	7	Total	C	N	O		0	0	0
			83	46	2	35				

- Molecule 8 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			6	3	3		
8	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	4	Total	Ca	0	0
			4	4		
9	B	2	Total	Ca	0	0
			2	2		

- Molecule 10 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	A	1	Total	C	N	O	0	0
			14	8	1	5		
10	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 11 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	B	1	Total	Mg	0	0
			1	1		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	412	Total	O	0	0
			412	412		
12	B	256	Total	O	0	0
			256	256		
12	C	12	Total	O	0	0
			12	12		
12	H	214	Total	O	0	0
			214	214		
12	L	248	Total	O	0	0
			248	248		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: INTEGRIN ALPHA-IIB




- Molecule 2: INTEGRIN BETA-3




- Molecule 3: FIBRINOGEN




- Molecule 4: MONOCLONAL ANTIBODY 10E5 HEAVY CHAIN




- Molecule 5: MONOCLONAL ANTIBODY 10E5 LIGHT CHAIN



- Molecule 6: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 7: α -D-mannopyranose-(1-3)-[α -D-mannopyranose-(1-6)] α -D-mannopyranose-(1-6)-[α -D-mannopyranose-(1-3)] α -D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	148.33Å 148.33Å 176.79Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.44 – 2.40 43.44 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.5 (43.44-2.40) 96.1 (43.44-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.148 , 0.193 0.154 , 0.198	Depositor DCC
R_{free} test set	4276 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	43.2	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 53.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	11884	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, MAN, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/3656	0.57	0/4980
2	B	0.34	0/3661	0.50	0/4960
3	C	0.41	0/53	0.61	0/67
4	H	0.35	0/1730	0.53	0/2369
5	L	0.36	0/1754	0.56	0/2377
All	All	0.36	0/10854	0.54	0/14753

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3539	0	3385	12	0
2	B	3580	0	3523	12	0
3	C	54	0	54	0	0
4	H	1678	0	1637	13	0
5	L	1686	0	1614	14	0
6	D	61	0	52	0	0
7	E	83	0	70	0	0
8	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	6	0	7	0	0
9	A	4	0	0	0	0
9	B	2	0	0	0	0
10	A	28	0	26	0	0
10	B	14	0	13	0	0
11	B	1	0	0	0	0
12	A	412	0	0	1	0
12	B	256	0	0	4	0
12	C	12	0	0	0	0
12	H	214	0	0	2	0
12	L	248	0	0	0	0
All	All	11884	0	10389	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:HD11	1.46	1.15
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:CD1	2.07	1.01
1:A:270:LEU:HD23	1:A:276[A]:ARG:HA	1.81	0.63
4:H:177[B]:GLN:CD	5:L:160[B]:LEU:CD1	2.68	0.61
1:A:270:LEU:HD23	1:A:276[B]:ARG:HA	1.82	0.60
2:B:201[A]:THR:HG22	12:B:4097:HOH:O	2.02	0.59
4:H:177[B]:GLN:CD	5:L:160[B]:LEU:HD11	2.21	0.59
5:L:48:ILE:HD13	5:L:54:LEU:HD12	1.85	0.58
5:L:136:LEU:HD21	5:L:146:VAL:HG22	1.84	0.58
4:H:177[B]:GLN:OE1	5:L:160[B]:LEU:HD12	1.99	0.57
4:H:177[A]:GLN:NE2	12:H:4197:HOH:O	2.32	0.54
4:H:30:LYS:NZ	12:H:4046:HOH:O	2.41	0.52
1:A:280:LEU:CD1	1:A:306:LEU:HD23	2.41	0.51
5:L:122:SER:O	5:L:126:THR:HG23	2.10	0.51
5:L:48:ILE:CD1	5:L:54:LEU:HD12	2.41	0.50
2:B:173:LEU:HD12	12:B:4068:HOH:O	2.11	0.50
2:B:88[B]:ILE:HD11	2:B:427:VAL:HG22	1.93	0.50
5:L:136:LEU:HD12	5:L:136:LEU:N	2.27	0.48
4:H:133:VAL:HG12	4:H:134:CYS:N	2.29	0.48
1:A:280:LEU:CD1	1:A:306:LEU:CD2	2.93	0.47
1:A:122:ALA:O	1:A:123:GLU:HB2	2.14	0.47
2:B:141:GLN:NE2	12:B:4053:HOH:O	2.46	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:308:PHE:CE2	2:B:328:THR:HG21	2.50	0.46
2:B:201[A]:THR:HG23	12:B:4095:HOH:O	2.15	0.46
5:L:37:GLN:HB2	5:L:47:LEU:HD11	1.98	0.46
1:A:278[A]:HIS:CD2	1:A:339:ALA:HB1	2.51	0.46
5:L:105[A]:GLU:HG3	5:L:173:TYR:OH	2.17	0.45
1:A:235:TRP:CZ2	1:A:270:LEU:HD11	2.52	0.45
2:B:223:PHE:CZ	2:B:254:THR:HG21	2.52	0.44
1:A:90[A]:ARG:CZ	12:A:4053:HOH:O	2.64	0.44
4:H:194:TRP:CG	4:H:195:PRO:HA	2.52	0.44
1:A:5:PRO:HA	1:A:8:LEU:HD21	2.00	0.44
4:H:208:SER:O	4:H:210:THR:HG23	2.18	0.43
2:B:443:PRO:HA	2:B:454:THR:HG22	2.01	0.43
1:A:278[A]:HIS:NE2	1:A:339:ALA:HB1	2.34	0.43
2:B:245:LEU:HD23	2:B:305:ASN:HB2	2.00	0.43
4:H:172:PHE:CD1	5:L:164:THR:HG23	2.54	0.42
1:A:297:ASP:O	1:A:372:ASN:HB2	2.20	0.42
1:A:262:TRP:HB3	2:B:317:LEU:HD13	2.01	0.42
2:B:15:GLN:O	2:B:19:VAL:HG23	2.20	0.41
4:H:177[B]:GLN:HB3	5:L:160[B]:LEU:HD11	2.02	0.41
4:H:35:HIS:O	4:H:96:CYS:HA	2.21	0.41
2:B:176:PRO:HB2	2:B:214:ARG:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	459/452 (102%)	450 (98%)	8 (2%)	1 (0%)	44	59
2	B	460/461 (100%)	440 (96%)	17 (4%)	3 (1%)	19	29
3	C	5/10 (50%)	4 (80%)	1 (20%)	0	100	100
4	H	218/221 (99%)	213 (98%)	5 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	L	223/214 (104%)	218 (98%)	5 (2%)	0	100	100
All	All	1365/1358 (100%)	1325 (97%)	36 (3%)	4 (0%)	37	51

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	GLU
2	B	9	GLY
2	B	10	VAL
2	B	80	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	369/360 (102%)	365 (99%)	4 (1%)	70	84
2	B	414/409 (101%)	412 (100%)	2 (0%)	86	94
3	C	5/6 (83%)	5 (100%)	0	100	100
4	H	192/190 (101%)	192 (100%)	0	100	100
5	L	199/188 (106%)	199 (100%)	0	100	100
All	All	1179/1153 (102%)	1173 (100%)	6 (0%)	86	94

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LEU
1	A	166	TYR
1	A	190	TYR
1	A	288	TYR
2	B	215	ASN
2	B	277	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	316	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	D	1	2,6	14,14,15	0.57	0	17,19,21	0.90	1 (5%)
6	NAG	D	2	6	14,14,15	0.68	0	17,19,21	0.80	0
6	MAN	D	3	6	11,11,12	0.52	0	15,15,17	0.58	0
6	MAN	D	4	6	11,11,12	0.53	0	15,15,17	0.60	0
6	MAN	D	5	6	11,11,12	0.61	0	15,15,17	1.07	2 (13%)
7	NAG	E	1	2,7	14,14,15	0.71	0	17,19,21	1.02	1 (5%)
7	NAG	E	2	7	14,14,15	0.56	0	17,19,21	0.82	0
7	MAN	E	3	7	11,11,12	0.64	0	15,15,17	0.65	0
7	MAN	E	4	7	11,11,12	0.48	0	15,15,17	1.16	1 (6%)
7	MAN	E	5	7	11,11,12	0.51	0	15,15,17	0.93	1 (6%)
7	MAN	E	6	7	11,11,12	0.57	0	15,15,17	0.72	1 (6%)
7	MAN	E	7	7	11,11,12	0.57	0	15,15,17	0.76	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	D	1	2,6	-	0/6/23/26	0/1/1/1
6	NAG	D	2	6	-	0/6/23/26	0/1/1/1
6	MAN	D	3	6	1/1/4/5	0/2/19/22	0/1/1/1
6	MAN	D	4	6	-	2/2/19/22	0/1/1/1
6	MAN	D	5	6	-	2/2/19/22	0/1/1/1
7	NAG	E	1	2,7	-	0/6/23/26	0/1/1/1
7	NAG	E	2	7	-	2/6/23/26	0/1/1/1
7	MAN	E	3	7	1/1/4/5	0/2/19/22	0/1/1/1
7	MAN	E	4	7	-	0/2/19/22	0/1/1/1
7	MAN	E	5	7	-	0/2/19/22	0/1/1/1
7	MAN	E	6	7	-	2/2/19/22	0/1/1/1
7	MAN	E	7	7	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	E	4	MAN	C1-O5-C5	3.64	117.06	112.19
6	D	5	MAN	C1-C2-C3	2.80	113.72	109.64
7	E	5	MAN	C1-O5-C5	2.59	115.66	112.19
7	E	7	MAN	C1-O5-C5	2.18	115.11	112.19
7	E	6	MAN	C1-O5-C5	2.09	114.99	112.19
6	D	1	NAG	O5-C1-C2	-2.09	108.06	111.29
6	D	5	MAN	C1-O5-C5	2.04	114.92	112.19
7	E	1	NAG	O5-C1-C2	-2.03	108.16	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	D	3	MAN	C1
7	E	3	MAN	C1

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	D	5	MAN	O5-C5-C6-O6
7	E	6	MAN	O5-C5-C6-O6
7	E	6	MAN	C4-C5-C6-O6
7	E	2	NAG	O5-C5-C6-O6
6	D	5	MAN	C4-C5-C6-O6
6	D	4	MAN	C4-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
7	E	2	NAG	C4-C5-C6-O6
6	D	4	MAN	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	NAG	A	3249	1	14,14,15	0.48	0	17,19,21	0.97	1 (5%)
10	NAG	A	3015	1	14,14,15	0.47	0	17,19,21	0.74	0
8	GOL	B	1462	9	5,5,5	0.41	0	5,5,5	0.35	0
8	GOL	A	1453	-	5,5,5	0.44	0	5,5,5	0.14	0
10	NAG	B	3099	2	14,14,15	0.49	0	17,19,21	0.87	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	NAG	A	3249	1	-	0/6/23/26	0/1/1/1
10	NAG	A	3015	1	-	0/6/23/26	0/1/1/1
8	GOL	B	1462	9	-	2/4/4/4	-
8	GOL	A	1453	-	-	0/4/4/4	-
10	NAG	B	3099	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	3249	NAG	O5-C1-C2	-2.37	107.63	111.29
10	B	3099	NAG	O5-C1-C2	-2.15	107.96	111.29

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	B	3099	NAG	C4-C5-C6-O6
10	B	3099	NAG	O5-C5-C6-O6
8	B	1462	GOL	O1-C1-C2-C3
8	B	1462	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

Warning: The R factor obtained from EDS is 0.2719, which does not match the depositor's R factor of 0.148. Please interpret the results in this section carefully.

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	452/452 (100%)	0.57	15 (3%)	49	46	23, 48, 66, 90	9 (1%)
2	B	455/461 (98%)	1.27	103 (22%)	3	3	23, 55, 95, 109	10 (2%)
3	C	7/10 (70%)	2.37	4 (57%)	0	0	39, 49, 108, 110	0
4	H	219/221 (99%)	0.58	14 (6%)	27	24	31, 48, 76, 106	3 (1%)
5	L	214/214 (100%)	0.40	3 (1%)	73	70	22, 47, 61, 99	9 (4%)
All	All	1347/1358 (99%)	0.79	139 (10%)	13	11	22, 50, 87, 110	31 (2%)

All (139) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	456	GLU	7.8
2	B	457	CYS	7.5
2	B	441	ALA	6.6
2	B	72	LYS	5.9
2	B	461	ARG	5.5
2	B	458	GLY	5.3
2	B	444	ASN	5.2
2	B	442	GLU	5.0
2	B	1	GLY	4.8
3	C	406	LYS	4.4
2	B	379	VAL	4.3
2	B	455	PHE	4.2
2	B	9	GLY	4.2
4	H	138	THR	4.0
2	B	10	VAL	3.9
3	C	405	ALA	3.9
2	B	395	VAL	3.8
2	B	405	GLY	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	H	137	THR	3.8
2	B	447	ARG	3.8
2	B	453	GLY	3.7
2	B	450	ASN	3.7
1	A	1	LEU	3.6
2	B	451	GLY	3.4
2	B	11	SER	3.4
2	B	24	ALA	3.4
2	B	422	LYS	3.3
2	B	452	ASN	3.3
4	H	221	PRO	3.3
2	B	460	CYS	3.3
2	B	51	PRO	3.2
2	B	437	CYS	3.2
4	H	134	CYS	3.2
2	B	397	PHE	3.2
2	B	436	ALA	3.2
2	B	443	PRO	3.2
2	B	91	ARG	3.2
2	B	446	HIS	3.1
2	B	49	CYS	3.1
2	B	393	ASP	3.0
2	B	392	GLY	3.0
2	B	22	MET	3.0
3	C	407	GLN	3.0
2	B	40	LEU	2.9
2	B	123	SER	2.9
2	B	34	GLY	2.9
4	H	195	PRO	2.9
2	B	459	VAL	2.9
2	B	32	PRO	2.9
2	B	372	ALA	2.9
2	B	454	THR	2.9
2	B	366	LEU	2.8
5	L	121	SER	2.8
4	H	133	VAL	2.8
2	B	380	ILE	2.8
2	B	61	ALA	2.8
2	B	33	LEU	2.8
2	B	374	CYS	2.8
2	B	56	PHE	2.8
2	B	35	SER	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	435	CYS	2.8
2	B	83	VAL	2.8
2	B	375	LEU	2.7
2	B	430	THR	2.7
2	B	31	LEU	2.7
2	B	433	CYS	2.7
2	B	79	GLN	2.7
2	B	59	SER	2.7
1	A	328	VAL	2.7
1	A	8	LEU	2.7
2	B	52	GLU	2.7
4	H	194	TRP	2.7
2	B	8	ARG	2.6
2	B	387[A]	MET	2.6
2	B	69	LEU	2.6
1	A	326	GLY	2.6
2	B	403	VAL	2.6
1	A	337	PRO	2.6
2	B	382	GLY	2.6
2	B	373	THR	2.6
5	L	213	GLU	2.6
2	B	53	SER	2.5
2	B	396	SER	2.5
2	B	445	SER	2.5
1	A	325	VAL	2.5
2	B	36	PRO	2.5
2	B	58	VAL	2.5
1	A	335[A]	ARG	2.5
2	B	253	LYS	2.5
2	B	92	LEU	2.5
2	B	121	SER	2.5
4	H	177[A]	GLN	2.5
2	B	449	ASN	2.4
1	A	323	ALA	2.4
1	A	248	LEU	2.4
2	B	67	ARG	2.4
4	H	193	THR	2.4
2	B	80	VAL	2.4
1	A	324	GLU	2.4
2	B	398	SER	2.4
2	B	23	CYS	2.3
5	L	212	ASN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	409	GLU	2.3
2	B	65	GLU	2.3
3	C	411	VAL	2.3
2	B	391	ILE	2.3
1	A	67	SER	2.3
4	H	196	SER	2.3
2	B	46	LYS	2.3
2	B	390	LYS	2.2
2	B	394	THR	2.2
2	B	419	VAL	2.2
2	B	30	ALA	2.2
4	H	220	GLY	2.2
4	H	186	SER	2.2
2	B	2	PRO	2.2
2	B	82	GLN	2.2
1	A	400	ARG	2.2
2	B	448	CYS	2.2
2	B	28	ASP	2.2
2	B	39	ASP	2.2
2	B	241	ASP	2.2
4	H	144	LEU	2.1
2	B	81	THR	2.1
2	B	319[A]	GLN	2.1
2	B	124	MET	2.1
2	B	84	SER	2.1
2	B	406	CYS	2.1
2	B	440	GLN	2.1
2	B	48	ASN	2.1
1	A	162[A]	TRP	2.1
2	B	45	LEU	2.1
1	A	327	ARG	2.1
1	A	130	CYS	2.1
4	H	175	VAL	2.0
2	B	412[A]	LYS	2.0
2	B	5	CYS	2.0
2	B	62	ARG	2.0
2	B	215	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

SUGAR-RSR INFOmissingINFO

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
10	NAG	A	3249	14/15	0.42	0.25	91,93,94,94	14
10	NAG	A	3015	14/15	0.51	0.28	89,91,93,94	14
10	NAG	B	3099	14/15	0.65	0.21	103,108,110,111	0
8	GOL	B	1462	6/6	0.87	0.20	62,62,63,65	0
8	GOL	A	1453	6/6	0.87	0.21	60,68,72,75	0
11	MG	B	2001	1/1	0.98	0.14	28,28,28,28	0
9	CA	B	2002	1/1	0.99	0.17	40,40,40,40	0
9	CA	A	2004	1/1	0.99	0.14	55,55,55,55	0
9	CA	A	2005	1/1	0.99	0.16	51,51,51,51	0
9	CA	A	2006	1/1	0.99	0.13	61,61,61,61	0
9	CA	A	2007	1/1	0.99	0.14	57,57,57,57	0
9	CA	B	2003	1/1	1.00	0.19	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.