



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2024 – 09:14 AM EDT

PDB ID : 3WL4
Title : N,N'-diacetylchitobiose deacetylase (Se-derivative) from *Pyrococcus furiosus*
Authors : Nakamura, T.; Niiyama, M.; Hashimoto, W.; Uegaki, K.
Deposited on : 2013-11-07
Resolution : 1.54 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3511 (1.56-1.52)
Clashscore	180529	3784 (1.56-1.52)
Ramachandran outliers	177936	3720 (1.56-1.52)
Sidechain outliers	177891	3717 (1.56-1.52)
RSRZ outliers	164620	3510 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 73% 24% .
1	B	267	 72% 24% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TAM	A	308	-	X	X	-
5	TAM	A	309	-	X	X	-
5	TAM	B	308	-	X	-	-
5	TAM	B	309	-	X	X	-
6	HEZ	A	316	-	-	X	-
6	HEZ	B	313	-	-	X	-
6	HEZ	B	320	-	-	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	Se	0	0	0
			2190	1416	360	404	3	7			
1	B	267	Total	C	N	O	S	Se	0	0	0
			2190	1416	360	404	3	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	expression tag	UNP Q8U3V1
B	1	MSE	-	expression tag	UNP Q8U3V1

- Molecule 2 is CADMIUM ION (three-letter code: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cd	0	0
			2	2		
2	B	2	Total	Cd	0	0
			2	2		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		
3	B	1	Total	Ca	0	0
			1	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Cl	0	0
			4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	4	Total	Cl	0	0
			4	4		

- Molecule 5 is TRIS(HYDROXYETHYL)AMINOMETHANE (three-letter code: TAM) (formula: $C_7H_{17}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			11	7	1	3		
5	A	1	Total	C	N	O	0	0
			11	7	1	3		
5	B	1	Total	C	N	O	0	0
			11	7	1	3		
5	B	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 6 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	A	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		
6	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 7 is HEXAN-1-OL (three-letter code: HE2) (formula: $C_6H_{14}O$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	6	1		
7	B	1	Total	C	O	0	0
			7	6	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	165	Total 165	O 165	0	0
8	B	156	Total 156	O 156	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	113.89Å 113.89Å 201.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.40 – 1.54 26.40 – 1.54	Depositor EDS
% Data completeness (in resolution range)	99.8 (26.40-1.54) 99.8 (26.40-1.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.63 (at 1.54Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.154 , 0.169 0.153 , 0.168	Depositor DCC
R_{free} test set	7197 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 29.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.476 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	4941	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEZ, CD, TAM, CL, HE2, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.74	25/2242 (1.1%)	1.74	51/3023 (1.7%)
1	B	1.77	31/2242 (1.4%)	1.70	41/3023 (1.4%)
All	All	1.76	56/4484 (1.2%)	1.72	92/6046 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

The worst 5 of 56 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	GLU	CG-CD	10.14	1.67	1.51
1	A	94	GLU	CD-OE1	9.15	1.35	1.25
1	B	166	GLN	CB-CG	-8.81	1.28	1.52
1	A	225	GLU	CG-CD	7.69	1.63	1.51
1	B	225	GLU	CG-CD	7.49	1.63	1.51

The worst 5 of 92 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	170	PHE	CB-CG-CD1	-11.50	112.75	120.80
1	A	232	ARG	NE-CZ-NH2	-10.42	115.09	120.30
1	A	153	ARG	NE-CZ-NH1	10.41	125.51	120.30
1	B	232	ARG	NE-CZ-NH2	-10.31	115.14	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	TYR	CB-CG-CD2	-10.18	114.89	121.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	121	ARG	Sidechain
1	B	121	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2190	0	2161	38	0
1	B	2190	0	2161	39	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0
5	A	22	0	31	14	0
5	B	22	0	32	11	0
6	A	72	0	125	25	0
6	B	96	0	168	27	0
7	A	7	0	14	0	0
7	B	7	0	13	4	0
8	A	165	0	0	9	0
8	B	156	0	0	7	0
All	All	4941	0	4705	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 107 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:313:HEZ:H62	8:B:536:HOH:O	1.25	1.29
5:A:309:TAM:H52	5:A:309:TAM:C4	1.74	1.16
6:A:316:HEZ:C2	8:A:532:HOH:O	1.95	1.12
5:A:309:TAM:C4	5:A:309:TAM:C5	2.29	1.09
5:A:309:TAM:H52	5:A:309:TAM:H42	1.11	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/267 (99%)	259 (98%)	5 (2%)	1 (0%)	30	13
1	B	265/267 (99%)	260 (98%)	4 (2%)	1 (0%)	30	13
All	All	530/534 (99%)	519 (98%)	9 (2%)	2 (0%)	30	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	ASP
1	B	42	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/232 (103%)	237 (99%)	2 (1%)	79	62
1	B	239/232 (103%)	238 (100%)	1 (0%)	89	81
All	All	478/464 (103%)	475 (99%)	3 (1%)	84	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	196	ASN
1	B	196	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	148	HIS
1	B	169	ASN
1	B	218	GLN
1	B	172	ASN
1	A	172	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 41 ligands modelled in this entry, 14 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	HEZ	B	313	-	7,7,7	1.93	2 (28%)	6,6,6	1.22	1 (16%)
6	HEZ	A	317	-	7,7,7	1.28	1 (14%)	6,6,6	1.99	2 (33%)
6	HEZ	A	312	-	7,7,7	0.28	0	6,6,6	1.07	0
6	HEZ	A	316	-	7,7,7	0.95	0	6,6,6	0.99	0
6	HEZ	A	318	-	7,7,7	1.35	0	6,6,6	1.21	0
6	HEZ	B	320	-	7,7,7	0.83	0	6,6,6	1.96	2 (33%)
5	TAM	A	309	2	10,10,10	3.19	9 (90%)	12,12,12	6.03	8 (66%)
6	HEZ	B	311	-	7,7,7	0.70	0	6,6,6	1.27	1 (16%)
6	HEZ	B	310	-	7,7,7	1.00	0	6,6,6	0.85	0
7	HE2	A	313	-	6,6,6	0.99	1 (16%)	5,5,5	0.85	0
6	HEZ	B	319	-	7,7,7	1.32	1 (14%)	6,6,6	0.55	0
5	TAM	B	309	2	10,10,10	3.07	7 (70%)	12,12,12	6.27	7 (58%)
6	HEZ	A	315	-	7,7,7	0.55	0	6,6,6	0.96	0
6	HEZ	B	318	-	7,7,7	1.06	0	6,6,6	2.00	3 (50%)
6	HEZ	B	312	-	7,7,7	1.05	0	6,6,6	0.95	0
6	HEZ	A	310	-	7,7,7	1.12	0	6,6,6	1.03	0
6	HEZ	B	314	-	7,7,7	1.34	1 (14%)	6,6,6	0.84	0
6	HEZ	B	322	-	7,7,7	0.30	0	6,6,6	1.52	1 (16%)
6	HEZ	A	314	-	7,7,7	0.99	0	6,6,6	2.00	2 (33%)
5	TAM	B	308	-	10,10,10	1.72	1 (10%)	12,12,12	4.33	9 (75%)
6	HEZ	A	311	-	7,7,7	0.29	0	6,6,6	1.76	2 (33%)
6	HEZ	B	315	-	7,7,7	0.48	0	6,6,6	1.20	1 (16%)
6	HEZ	B	317	-	7,7,7	1.19	1 (14%)	6,6,6	0.97	0
6	HEZ	B	321	-	7,7,7	0.58	0	6,6,6	1.14	0
7	HE2	B	316	-	6,6,6	1.11	0	5,5,5	1.42	1 (20%)
6	HEZ	A	319	-	7,7,7	0.78	0	6,6,6	1.60	2 (33%)
5	TAM	A	308	-	10,10,10	1.38	2 (20%)	12,12,12	6.09	8 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HEZ	B	313	-	-	4/5/5/5	-
6	HEZ	A	317	-	-	4/5/5/5	-
6	HEZ	A	312	-	-	0/5/5/5	-
6	HEZ	A	316	-	-	4/5/5/5	-
6	HEZ	A	318	-	-	3/5/5/5	-
6	HEZ	B	320	-	-	4/5/5/5	-
5	TAM	A	309	2	-	3/12/12/12	-
6	HEZ	B	311	-	-	3/5/5/5	-
6	HEZ	B	310	-	-	0/5/5/5	-
7	HE2	A	313	-	-	2/4/4/4	-
6	HEZ	B	319	-	-	3/5/5/5	-
5	TAM	B	309	2	-	8/12/12/12	-
6	HEZ	A	315	-	-	3/5/5/5	-
6	HEZ	B	318	-	-	2/5/5/5	-
6	HEZ	B	312	-	-	0/5/5/5	-
6	HEZ	A	310	-	-	0/5/5/5	-
6	HEZ	B	314	-	-	2/5/5/5	-
6	HEZ	B	322	-	-	2/5/5/5	-
6	HEZ	A	314	-	-	3/5/5/5	-
5	TAM	B	308	-	-	7/12/12/12	-
6	HEZ	A	311	-	-	2/5/5/5	-
6	HEZ	B	315	-	-	1/5/5/5	-
6	HEZ	B	317	-	-	4/5/5/5	-
6	HEZ	B	321	-	-	4/5/5/5	-
7	HE2	B	316	-	-	2/4/4/4	-
6	HEZ	A	319	-	-	1/5/5/5	-
5	TAM	A	308	-	-	4/12/12/12	-

The worst 5 of 26 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	309	TAM	C2-C5	-4.75	1.43	1.52
6	B	313	HEZ	C4-C3	4.37	1.73	1.51
5	B	309	TAM	C3-C6	-4.17	1.44	1.52
5	A	309	TAM	C3-C6	-4.16	1.44	1.52
5	A	309	TAM	C2-C	-4.15	1.47	1.53

The worst 5 of 50 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	309	TAM	C5-C2-C	-12.67	101.02	115.97
5	B	309	TAM	O4-C4-C1	-10.77	81.10	111.33
5	A	308	TAM	C6-C3-C	-10.62	103.44	115.97
5	B	309	TAM	C6-C3-C	-10.35	103.77	115.97
5	B	309	TAM	C4-C1-C	10.21	128.01	115.97

There are no chirality outliers.

5 of 75 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	308	TAM	C1-C-C2-C5
5	A	308	TAM	C3-C-C2-C5
5	A	308	TAM	N-C-C2-C5
5	A	308	TAM	C-C3-C6-O6
5	A	309	TAM	C2-C-C1-C4

There are no ring outliers.

16 monomers are involved in 81 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	313	HEZ	6	0
6	A	317	HEZ	1	0
6	A	316	HEZ	12	0
6	A	318	HEZ	5	0
6	B	320	HEZ	14	0
5	A	309	TAM	7	0
6	B	319	HEZ	4	0
5	B	309	TAM	6	0
6	A	315	HEZ	1	0
6	B	318	HEZ	3	0
6	A	314	HEZ	4	0
5	B	308	TAM	5	0
6	A	311	HEZ	1	0
7	B	316	HE2	4	0
6	A	319	HEZ	2	0
5	A	308	TAM	7	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/267 (97%)	-1.23	0 100 100	13, 19, 37, 52	0
1	B	260/267 (97%)	-1.24	0 100 100	13, 19, 36, 53	0
All	All	520/534 (97%)	-1.24	0 100 100	13, 19, 36, 53	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	HEZ	B	315	8/8	0.97	0.07	28,32,37,41	0
7	HE2	A	313	7/7	0.97	0.07	27,30,38,40	0
6	HEZ	A	315	8/8	0.98	0.07	40,56,69,71	0
6	HEZ	A	316	8/8	0.98	0.06	32,51,58,60	0
6	HEZ	B	311	8/8	0.98	0.07	25,41,52,54	0
6	HEZ	B	313	8/8	0.98	0.08	16,38,49,52	0
5	TAM	A	309	11/11	0.98	0.06	15,19,41,48	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	HEZ	B	317	8/8	0.98	0.07	39,48,51,56	0
6	HEZ	B	319	8/8	0.98	0.07	30,40,71,74	0
6	HEZ	B	321	8/8	0.98	0.09	45,62,72,73	0
6	HEZ	A	314	8/8	0.98	0.07	16,35,40,40	0
7	HE2	B	316	7/7	0.98	0.05	21,27,34,34	0
6	HEZ	A	318	8/8	0.99	0.06	31,40,48,51	0
6	HEZ	A	319	8/8	0.99	0.04	33,39,41,42	0
6	HEZ	B	310	8/8	0.99	0.04	22,30,32,34	0
5	TAM	B	309	11/11	0.99	0.06	16,22,33,44	0
6	HEZ	B	312	8/8	0.99	0.04	22,26,32,33	0
6	HEZ	A	310	8/8	0.99	0.04	23,26,33,34	0
6	HEZ	B	314	8/8	0.99	0.05	33,41,43,50	0
6	HEZ	A	311	8/8	0.99	0.05	25,44,48,49	0
6	HEZ	A	312	8/8	0.99	0.06	30,45,54,57	0
6	HEZ	B	318	8/8	0.99	0.06	17,34,47,48	0
5	TAM	A	308	11/11	0.99	0.04	14,19,21,27	11
6	HEZ	B	320	8/8	0.99	0.06	18,28,43,47	0
4	CL	B	306	1/1	0.99	0.03	23,23,23,23	0
6	HEZ	B	322	8/8	0.99	0.04	31,43,56,56	0
5	TAM	B	308	11/11	0.99	0.04	13,17,24,24	11
6	HEZ	A	317	8/8	0.99	0.05	22,35,41,42	0
2	CD	B	301	1/1	1.00	0.01	16,16,16,16	0
2	CD	B	302	1/1	1.00	0.02	20,20,20,20	0
3	CA	A	303	1/1	1.00	0.01	17,17,17,17	0
3	CA	B	303	1/1	1.00	0.01	16,16,16,16	0
4	CL	A	304	1/1	1.00	0.05	19,19,19,19	0
4	CL	A	305	1/1	1.00	0.02	20,20,20,20	0
4	CL	A	306	1/1	1.00	0.04	22,22,22,22	0
4	CL	A	307	1/1	1.00	0.03	23,23,23,23	0
4	CL	B	304	1/1	1.00	0.04	19,19,19,19	0
4	CL	B	305	1/1	1.00	0.03	22,22,22,22	0
2	CD	A	301	1/1	1.00	0.01	16,16,16,16	0
4	CL	B	307	1/1	1.00	0.02	20,20,20,20	0
2	CD	A	302	1/1	1.00	0.02	20,20,20,20	0

6.5 Other polymers ⓘ

There are no such residues in this entry.