



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 6, 2024 – 05:40 PM JST

PDB ID : 4WL2
Title : Structure of penicillin V acylase from *Pectobacterium atrosepticum*
Authors : Ramasamy, S.; Avinash, V.S.; Pundle, A.V.; Suresh, C.G.
Deposited on : 2014-10-06
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5504 (2.50-2.50)
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	355	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>...</div> </div> </div>
1	B	355	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>8%</div> <div>5%</div> <div>...</div> </div> </div>
1	C	355	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>...</div> </div> </div>
1	D	355	<div> <div>4%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>...</div> </div> </div>
1	E	355	<div> <div>7%</div> <div> <div></div> <div>82%</div> <div>13%</div> <div>...</div> </div> </div>
1	F	355	<div> <div>8%</div> <div> <div></div> <div>86%</div> <div>8%</div> <div>...</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	355	<div> <div>5%</div> <div>85%</div> <div>9%</div> <div>• •</div> </div>
1	H	355	<div> <div>10%</div> <div>82%</div> <div>12%</div> <div>• • •</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 21734 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative exported choloylglycine hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	B	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	C	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	D	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	E	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	F	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	G	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			
1	H	347	Total	C	N	O	S	0	0	0
			2711	1730	462	513	6			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	348	LEU	-	expression tag	UNP Q6D291
A	349	GLU	-	expression tag	UNP Q6D291
A	350	HIS	-	expression tag	UNP Q6D291
A	351	HIS	-	expression tag	UNP Q6D291
A	352	HIS	-	expression tag	UNP Q6D291
A	353	HIS	-	expression tag	UNP Q6D291
A	354	HIS	-	expression tag	UNP Q6D291
A	355	HIS	-	expression tag	UNP Q6D291
B	348	LEU	-	expression tag	UNP Q6D291
B	349	GLU	-	expression tag	UNP Q6D291
B	350	HIS	-	expression tag	UNP Q6D291
B	351	HIS	-	expression tag	UNP Q6D291
B	352	HIS	-	expression tag	UNP Q6D291

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	353	HIS	-	expression tag	UNP Q6D291
B	354	HIS	-	expression tag	UNP Q6D291
B	355	HIS	-	expression tag	UNP Q6D291
C	348	LEU	-	expression tag	UNP Q6D291
C	349	GLU	-	expression tag	UNP Q6D291
C	350	HIS	-	expression tag	UNP Q6D291
C	351	HIS	-	expression tag	UNP Q6D291
C	352	HIS	-	expression tag	UNP Q6D291
C	353	HIS	-	expression tag	UNP Q6D291
C	354	HIS	-	expression tag	UNP Q6D291
C	355	HIS	-	expression tag	UNP Q6D291
D	348	LEU	-	expression tag	UNP Q6D291
D	349	GLU	-	expression tag	UNP Q6D291
D	350	HIS	-	expression tag	UNP Q6D291
D	351	HIS	-	expression tag	UNP Q6D291
D	352	HIS	-	expression tag	UNP Q6D291
D	353	HIS	-	expression tag	UNP Q6D291
D	354	HIS	-	expression tag	UNP Q6D291
D	355	HIS	-	expression tag	UNP Q6D291
E	348	LEU	-	expression tag	UNP Q6D291
E	349	GLU	-	expression tag	UNP Q6D291
E	350	HIS	-	expression tag	UNP Q6D291
E	351	HIS	-	expression tag	UNP Q6D291
E	352	HIS	-	expression tag	UNP Q6D291
E	353	HIS	-	expression tag	UNP Q6D291
E	354	HIS	-	expression tag	UNP Q6D291
E	355	HIS	-	expression tag	UNP Q6D291
F	348	LEU	-	expression tag	UNP Q6D291
F	349	GLU	-	expression tag	UNP Q6D291
F	350	HIS	-	expression tag	UNP Q6D291
F	351	HIS	-	expression tag	UNP Q6D291
F	352	HIS	-	expression tag	UNP Q6D291
F	353	HIS	-	expression tag	UNP Q6D291
F	354	HIS	-	expression tag	UNP Q6D291
F	355	HIS	-	expression tag	UNP Q6D291
G	348	LEU	-	expression tag	UNP Q6D291
G	349	GLU	-	expression tag	UNP Q6D291
G	350	HIS	-	expression tag	UNP Q6D291
G	351	HIS	-	expression tag	UNP Q6D291
G	352	HIS	-	expression tag	UNP Q6D291
G	353	HIS	-	expression tag	UNP Q6D291
G	354	HIS	-	expression tag	UNP Q6D291

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	355	HIS	-	expression tag	UNP Q6D291
H	348	LEU	-	expression tag	UNP Q6D291
H	349	GLU	-	expression tag	UNP Q6D291
H	350	HIS	-	expression tag	UNP Q6D291
H	351	HIS	-	expression tag	UNP Q6D291
H	352	HIS	-	expression tag	UNP Q6D291
H	353	HIS	-	expression tag	UNP Q6D291
H	354	HIS	-	expression tag	UNP Q6D291
H	355	HIS	-	expression tag	UNP Q6D291


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	11	Total O 11 11	0	0
2	B	3	Total O 3 3	0	0
2	C	5	Total O 5 5	0	0
2	D	7	Total O 7 7	0	0
2	E	5	Total O 5 5	0	0
2	F	2	Total O 2 2	0	0
2	G	9	Total O 9 9	0	0
2	H	4	Total O 4 4	0	0



i

- Molecule 1: Putative exported choloylglycine hydrolase









- Molecule 1: Putative exported choloylglycine hydrolase





- Molecule 1: Putative exported choloylglycine hydrolase



- Molecule 1: Putative exported choloylglycine hydrolase



- Molecule 1: Putative exported choloylglycine hydrolase



HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	118.45Å 150.21Å 185.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	63.43 – 2.50 63.43 – 2.50	Depositor EDS
% Data completeness (in resolution range)	88.0 (63.43-2.50) 88.0 (63.43-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.218 , 0.248 0.221 , 0.249	Depositor DCC
R_{free} test set	5077 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	39.2	Xtriage
Anisotropy	0.136	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21734	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2781	0.92	17/3779 (0.4%)
1	B	0.66	0/2781	0.83	7/3779 (0.2%)
1	C	0.73	2/2781 (0.1%)	0.86	10/3779 (0.3%)
1	D	0.67	1/2781 (0.0%)	0.86	10/3779 (0.3%)
1	E	0.70	2/2781 (0.1%)	0.89	10/3779 (0.3%)
1	F	0.65	1/2781 (0.0%)	0.84	9/3779 (0.2%)
1	G	0.62	0/2781	0.82	4/3779 (0.1%)
1	H	0.74	3/2781 (0.1%)	0.90	12/3779 (0.3%)
All	All	0.68	9/22248 (0.0%)	0.87	79/30232 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	2
1	E	0	1
1	F	0	2
1	H	0	2
All	All	0	10

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	233	PRO	CA-C	10.92	1.74	1.52
1	C	235	VAL	N-CA	10.70	1.67	1.46
1	H	233	PRO	N-CA	8.77	1.62	1.47
1	E	234	GLY	N-CA	6.78	1.56	1.46
1	E	184	SER	CA-CB	-6.53	1.43	1.52

The worst 5 of 79 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	ARG	NE-CZ-NH2	17.32	128.96	120.30
1	E	184	SER	C-N-CD	11.51	152.56	128.40
1	G	150	HIS	CB-CA-C	-11.24	87.91	110.40
1	F	184	SER	C-N-CD	10.76	150.99	128.40
1	D	276	ARG	NE-CZ-NH1	8.69	124.64	120.30

There are no chirality outliers.

5 of 10 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	346	GLY	Peptide
1	B	137	THR	Peptide
1	C	346	GLY	Peptide
1	D	1	CYS	Peptide
1	D	139	GLN	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2711	0	2660	72	0
1	B	2711	0	2660	43	0
1	C	2711	0	2660	30	18
1	D	2711	0	2658	44	18
1	E	2711	0	2660	27	13
1	F	2711	0	2659	22	0
1	G	2711	0	2660	27	0
1	H	2711	0	2660	39	13
2	A	11	0	0	2	0
2	B	3	0	0	0	0
2	C	5	0	0	2	0
2	D	7	0	0	4	0
2	E	5	0	0	0	0
2	F	2	0	0	0	0
2	G	9	0	0	1	0
2	H	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	21734	0	21277	291	31

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:235:VAL:N	1:C:235:VAL:CA	1.67	1.56
1:H:233:PRO:C	1:H:233:PRO:CA	1.74	1.54
1:D:68:GLY:O	1:D:138:LYS:NZ	1.57	1.34
1:A:3:ARG:NH1	1:A:181:MET:SD	2.03	1.31
1:B:209:THR:HG22	1:B:214:ASP:OD2	1.21	1.28

The worst 5 of 31 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:235:VAL:N	1:D:234:GLY:C[4_555]	0.89	1.31
1:C:235:VAL:N	1:D:234:GLY:CA[4_555]	1.01	1.19
1:C:234:GLY:O	1:D:234:GLY:O[4_555]	1.05	1.15
1:E:233:PRO:C	1:H:233:PRO:N[4_455]	1.06	1.14
1:E:233:PRO:O	1:H:233:PRO:N[4_455]	1.13	1.07

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	345/355 (97%)	334 (97%)	8 (2%)	3 (1%)	14	28
1	B	345/355 (97%)	336 (97%)	7 (2%)	2 (1%)	22	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	345/355 (97%)	338 (98%)	6 (2%)	1 (0%)	37	56
1	D	345/355 (97%)	338 (98%)	7 (2%)	0	100	100
1	E	345/355 (97%)	337 (98%)	6 (2%)	2 (1%)	22	39
1	F	345/355 (97%)	336 (97%)	7 (2%)	2 (1%)	22	39
1	G	345/355 (97%)	337 (98%)	7 (2%)	1 (0%)	37	56
1	H	345/355 (97%)	332 (96%)	11 (3%)	2 (1%)	22	39
All	All	2760/2840 (97%)	2688 (97%)	59 (2%)	13 (0%)	25	44

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	147	ALA
1	H	138	LYS
1	H	44	ALA
1	A	143	GLN
1	A	210	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/297 (97%)	272 (94%)	17 (6%)	16	33
1	B	289/297 (97%)	268 (93%)	21 (7%)	11	24
1	C	289/297 (97%)	272 (94%)	17 (6%)	16	33
1	D	289/297 (97%)	270 (93%)	19 (7%)	14	28
1	E	289/297 (97%)	264 (91%)	25 (9%)	8	17
1	F	289/297 (97%)	270 (93%)	19 (7%)	14	28
1	G	289/297 (97%)	271 (94%)	18 (6%)	15	31
1	H	289/297 (97%)	271 (94%)	18 (6%)	15	31
All	All	2312/2376 (97%)	2158 (93%)	154 (7%)	13	28

5 of 154 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	276	ARG
1	H	97	LYS
1	G	19	ARG
1	G	143	GLN
1	H	329	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	340	GLN
1	F	210	ASN
1	H	331	GLN
1	H	36	GLN
1	H	179	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	347/355 (97%)	0.13	16 (4%) 38 35	19, 34, 75, 172	0
1	B	347/355 (97%)	0.22	17 (4%) 36 33	18, 36, 77, 175	0
1	C	347/355 (97%)	0.08	18 (5%) 34 31	20, 34, 73, 201	0
1	D	347/355 (97%)	0.08	13 (3%) 45 42	18, 33, 69, 128	0
1	E	347/355 (97%)	0.19	25 (7%) 23 21	18, 34, 81, 182	0
1	F	347/355 (97%)	0.40	28 (8%) 19 18	20, 39, 83, 163	0
1	G	347/355 (97%)	0.21	18 (5%) 34 31	21, 39, 75, 199	0
1	H	347/355 (97%)	0.48	37 (10%) 12 12	21, 41, 86, 149	0
All	All	2776/2840 (97%)	0.22	172 (6%) 28 26	18, 36, 80, 201	0

The worst 5 of 172 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	347	LEU	8.5
1	E	347	LEU	8.4
1	A	347	LEU	8.0
1	D	140	VAL	8.0
1	A	147	ALA	7.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.