



wwPDB EM Validation Summary Report ⓘ

Jun 10, 2025 – 02:12 PM JST

PDB ID : 7WLK / pdb_00007wlk
EMDB ID : EMD-32586
Title : CryoEM structure of human low-voltage activated T-type calcium channel Cav3.3 in complex with Otilonium Bromide(OB)
Authors : He, L.; Yu, Z.; Dong, Y.; Chen, Q.; Zhao, Y.
Deposited on : 2022-01-13
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0rc1
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	2223	 35% 16% 49%

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Voltage-dependent T-type calcium channel subunit alpha-1I.


Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1135	Total	C	N	O	S	1	0
			9111	5979	1499	1555	78		

- Molecule 2 is 2-[diethyl(methyl)- λ^4 -azanyl]ethyl 4-[(2-octoxyphenyl)carbonylamino]benzoate (CCD ID: 7TB) (formula: $C_{29}H_{43}N_2O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				AltConf
2	A	1	Total	C	N	O	0
			35	29	2	4	

- Molecule 3 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: $C_{41}H_{82}NO_8P$).



Mol	Chain	Residues	Atoms				AltConf
3	A	1	Total	C	O	P	0
			33	24	8	1	
3	A	1	Total	C			0
			9	9			
3	A	1	Total	C			0
			11	11			
3	A	1	Total	C			0
			12	12			
3	A	1	Total	C			0
			12	12			
3	A	1	Total	C			0
			12	12			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
4	A	2	Total	Ca	0
			2	2	

- Molecule 5 is CHOLESTEROL HEMISUCCINATE (CCD ID: Y01) (formula: C₃₁H₅₀O₄).



- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				AltConf
6	A	1	Total	C	N	O	0
			14	8	1	5	
6	A	1	Total	C	N	O	0
			14	8	1	5	

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Voltage-dependent T-type calcium channel subunit alpha-1I





GLY	GLY	ALA	THR
GLU	GLY	ARG	THR
LEU	SER	LYS	LEU
GLU	THR	PHE	GLU
PRO	SER	SER	ASP
GLY	SER	SER	LEU
ASP	THR	THR	GLY
ALA	CYS	SER	THR
ALA	THR	ALA	LEU
SER	HIS	SER	SER
LYS	HIS	ALA	ASP
ARG	HIS	ALA	SER
LYS	ASP	PRO	PRO
ARG	SER	GLY	PRO
	MET	ARG	ARG
	ASP	ARG	ALA
	PRO	PRO	LEU
	SER	HIS	GLY
	LEU	ALA	GLY
	GLY	ALA	GLU
	ASP	ALA	PRO
	SER	ALA	PRO
	LEU	ALA	ALA
	GLY	ALA	ALA
	LEU	GLY	ALA
	ALA	LEU	GLY
	ALA	ARG	GLY
	GLY	SER	GLY
	PRO	PRO	LEU
	SER	SER	ARG
	TRP	THR	ARG
	ALA	ALA	ARG
	ALA	LYS	ARG
	ASP	ASP	LEU
	ARG	PRO	LEU
	SER	GLY	ARG
	THR	ARG	GLY
	LEU	ALA	GLY
	LEU	PRO	ARG
	LEU	LEU	GLY
	PRO	PRO	LEU
	PRO	PRO	ARG
	PRO	ALA	ARG
	PRO	ALA	GLY
	GLN	PRO	GLY
	PRO	PRO	HIS
	LEU	LEU	SER
	PRO	PRO	THR

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	1684039	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	9.6	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.101	Depositor
Minimum map value	-0.059	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.015	Depositor
Map size (\AA)	266.24, 266.24, 266.24	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.04, 1.04, 1.04	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, Y01, 3PE, NAG, 7TB

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/9314	0.46	1/12625 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	176	ALA	N-CA-C	-5.57	106.32	114.39

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9111	0	9291	260	0
2	A	35	0	0	0	0
3	A	89	0	131	2	0
4	A	2	0	0	0	0
5	A	196	0	279	23	0
6	A	28	0	26	1	0
All	All	9461	0	9727	276	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 276 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:861:LEU:CD1	1:A:1423:LEU:CD2	1.85	1.51
1:A:861:LEU:HD12	1:A:1423:LEU:CD2	1.03	1.49
1:A:861:LEU:CD1	1:A:1423:LEU:HD22	1.51	1.35
1:A:861:LEU:CD1	1:A:1423:LEU:HD21	1.70	1.03
1:A:861:LEU:HD11	1:A:1423:LEU:HD22	1.35	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1128/2223 (51%)	1053 (93%)	75 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1006/1879 (54%)	1005 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1379	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1511	GLN
1	A	1718	GLN
1	A	1353	ASN
1	A	1362	ASN
1	A	1487	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	Y01	A	2316	-	38,38,38	0.62	0	57,57,57	1.69	12 (21%)
3	3PE	A	2306	-	11,11,50	1.50	0	10,10,55	0.89	0
3	3PE	A	2305	-	10,10,50	1.50	0	9,9,55	0.85	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	3PE	A	2315	-	11,11,50	1.51	0	10,10,55	0.85	0
5	Y01	A	2310	-	38,38,38	0.75	1 (2%)	57,57,57	1.62	12 (21%)
3	3PE	A	2303	-	8,8,50	1.47	0	7,7,55	0.76	0
3	3PE	A	2307	-	11,11,50	1.51	0	10,10,55	0.84	0
6	NAG	A	2311	1	14,14,15	0.38	0	17,19,21	1.29	1 (5%)
6	NAG	A	2313	1	14,14,15	0.27	0	17,19,21	0.50	0
5	Y01	A	2314	-	32,32,38	0.81	2 (6%)	49,49,57	1.63	12 (24%)
3	3PE	A	2302	-	32,32,50	2.36	11 (34%)	36,37,55	1.27	2 (5%)
5	Y01	A	2308	-	34,34,38	0.73	1 (2%)	52,52,57	1.64	9 (17%)
2	7TB	A	2301	-	36,36,36	1.64	8 (22%)	46,46,46	1.29	5 (10%)
5	Y01	A	2312	-	38,38,38	0.71	0	57,57,57	1.72	13 (22%)
5	Y01	A	2309	-	34,34,38	0.78	1 (2%)	52,52,57	1.72	10 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	Y01	A	2316	-	-	11/19/77/77	0/4/4/4
3	3PE	A	2306	-	-	4/9/9/54	-
3	3PE	A	2305	-	-	3/8/8/54	-
3	3PE	A	2315	-	-	4/9/9/54	-
5	Y01	A	2310	-	-	13/19/77/77	0/4/4/4
3	3PE	A	2303	-	-	0/6/6/54	-
3	3PE	A	2307	-	-	4/9/9/54	-
6	NAG	A	2311	1	-	5/6/23/26	0/1/1/1
6	NAG	A	2313	1	-	3/6/23/26	0/1/1/1
5	Y01	A	2314	-	-	8/12/70/77	0/4/4/4
3	3PE	A	2302	-	-	20/34/34/54	-
5	Y01	A	2308	-	-	6/14/72/77	0/4/4/4
2	7TB	A	2301	-	-	10/34/34/34	0/2/2/2
5	Y01	A	2312	-	-	13/19/77/77	0/4/4/4
5	Y01	A	2309	-	-	11/14/72/77	0/4/4/4

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2301	7TB	C32-N07	5.32	1.49	1.35
3	A	2302	3PE	P-O13	4.76	1.73	1.54
3	A	2302	3PE	O31-C31	4.71	1.47	1.33
3	A	2302	3PE	O21-C21	4.33	1.46	1.34
3	A	2302	3PE	P-O11	4.13	1.73	1.60

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	2312	Y01	CBI-CBE-CBB	-5.40	111.03	119.49
5	A	2309	Y01	OAW-CAY-CAM	5.01	120.30	111.09
5	A	2309	Y01	CBI-CBE-CBB	-4.97	111.71	119.49
5	A	2308	Y01	OAW-CAY-CAM	4.86	120.03	111.09
5	A	2316	Y01	CBI-CBE-CBB	-4.75	112.05	119.49

There are no chirality outliers.

5 of 115 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2302	3PE	C1-O11-P-O12
3	A	2302	3PE	C1-O11-P-O13
5	A	2309	Y01	CAC-CBB-CBE-CBI
5	A	2309	Y01	OAG-CAY-OAW-CBC
5	A	2309	Y01	CAM-CAY-OAW-CBC


There are no ring outliers.




9 monomers are involved in 26 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	2316	Y01	6	0
3	A	2315	3PE	1	0
5	A	2310	Y01	3	0
6	A	2311	NAG	1	0
5	A	2314	Y01	7	0
3	A	2302	3PE	1	0
5	A	2308	Y01	3	0
5	A	2312	Y01	3	0
5	A	2309	Y01	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will


also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32586. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 128



Y Index: 128




Z Index: 128


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 143



Y Index: 135




Z Index: 141


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 71 nm^3 ; this corresponds to an approximate mass of 64 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-32586 and PDB model 7WLK. Per-residue inclusion information can be found in [section 3](#) on [page 7](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).

9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7440	<div></div> 0.4380
A	<div></div> 0.7440	<div></div> 0.4380

