



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 2, 2025 – 02:03 am BST

PDB ID : 2WR1 / pdb\_00002wr1  
Title : structure of influenza H2 hemagglutinin with human receptor  
Authors : Liu, J.; Stevens, D.J.; Haire, L.F.; Walker, P.A.; Coombs, P.J.; Russell, R.J.;  
Gamblin, S.J.; Skehel, J.J.  
Deposited on : 2009-08-29  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	509	<div> <div>3%</div> <div>75%</div> <div>18%</div> <div>• •</div> </div>
1	B	509	<div> <div>5%</div> <div>72%</div> <div>21%</div> <div>• • •</div> </div>
1	C	509	<div> <div>3%</div> <div>73%</div> <div>19%</div> <div>• 5%</div> </div>
2	D	3	<div> <div>100%</div> </div>
2	G	3	<div> <div>100%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	E	2	 100%
4	F	4	 50% 25% 25%
5	H	4	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	F	1	-	-	X	-

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 13094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMAGGLUTININ.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	490	Total	C	N	O	S	4	0	0
			3882	2436	671	753	22			
1	B	490	Total	C	N	O	S	0	0	0
			3879	2434	670	753	22			
1	C	485	Total	C	N	O	S	0	0	0
			3844	2413	664	745	22			

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			46	25	2	19			
2	G	3	Total	C	N	O	0	0	0
			46	25	2	19			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-3)-beta-D-mannopyranose

-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	4	Total	C	N	O	0	0	1
			40	23	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	C	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	433	Total	O	0	0
			433	433		
7	B	406	Total	O	0	0
			406	406		
7	C	398	Total	O	0	0
			398	398		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: HEMAGGLUTININ




#### • Molecule 1: HEMAGGLUTININ



- Molecule 1: HEMAGGLUTININ







RES1
RES2
RES3
RES4

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.56Å 141.46Å 199.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.10 30.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.10) 99.8 (30.00-2.10)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 2.10Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.196 , 0.230 0.194 , 0.226	Depositor DCC
$R_{free}$ test set	5783 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.5	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 58.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13094	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SIA, GAL, BMA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.02	2/3968 (0.1%)	0.79	2/5372 (0.0%)
1	B	1.04	5/3962 (0.1%)	0.82	6/5361 (0.1%)
1	C	0.94	2/3927 (0.1%)	0.77	3/5313 (0.1%)
All	All	1.00	9/11857 (0.1%)	0.79	11/16046 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	400	SER	CB-OG	6.17	1.50	1.42
1	B	464	ASN	CG-ND2	6.06	1.48	1.32
1	C	434	GLU	CD-OE1	5.84	1.32	1.25
1	B	102	TYR	CE2-CZ	5.42	1.45	1.38
1	A	160	TYR	CE2-CZ	5.32	1.45	1.38

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	211	ARG	NE-CZ-NH1	-9.17	115.71	120.30
1	A	404	ARG	NE-CZ-NH1	-7.38	116.61	120.30
1	B	211	ARG	NE-CZ-NH1	-7.28	116.66	120.30
1	B	462	LEU	CB-CG-CD2	-7.04	99.03	111.00
1	C	23	ILE	CG1-CB-CG2	-6.26	97.62	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	492	GLU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3882	0	3742	104	0
1	B	3879	0	3749	130	0
1	C	3844	0	3718	126	0
2	D	46	0	40	5	0
2	G	46	0	40	6	0
3	E	28	0	25	7	0
4	F	40	0	33	7	0
5	H	50	0	43	6	0
6	A	28	0	26	6	0
6	C	14	0	13	3	0
7	A	433	0	0	21	0
7	B	406	0	0	27	0
7	C	398	0	0	29	0
All	All	13094	0	11429	362	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 362 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:483:ASN:HD21	5:H:1:NAG:C1	1.51	1.21
1:B:168:ASN:HD21	3:E:1:NAG:C1	1.56	1.19
1:C:168:ASN:HD21	6:C:1500:NAG:C1	1.58	1.17
1:B:477:CYS:SG	7:B:2394:HOH:O	2.04	1.13
1:A:148:ARG:HG2	1:A:148:ARG:HH11	1.03	1.09

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/509 (96%)	465 (96%)	17 (4%)	4 (1%)	16	13
1	B	484/509 (95%)	462 (96%)	18 (4%)	4 (1%)	16	13
1	C	479/509 (94%)	452 (94%)	24 (5%)	3 (1%)	22	19
All	All	1449/1527 (95%)	1379 (95%)	59 (4%)	11 (1%)	16	13

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	ASP
1	A	494	GLU
1	B	464	ASN
1	B	494	GLU
1	C	495	SER

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/447 (96%)	409 (96%)	18 (4%)	25	26
1	B	428/447 (96%)	413 (96%)	15 (4%)	31	34
1	C	425/447 (95%)	408 (96%)	17 (4%)	27	28
All	All	1280/1341 (96%)	1230 (96%)	50 (4%)	27	29

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	390	THR
1	C	131	THR
1	C	494	GLU
1	B	437	LEU
1	B	494	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 38 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	168	ASN
1	C	454	GLN
1	C	182	HIS
1	C	249	ASN
1	C	483	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

Of 16 monosaccharides modelled in this entry, 15 were used for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	D	1	2	15,15,15	1.10	1 (6%)	21,21,21	1.97	5 (23%)
2	GAL	D	2	2	11,11,12	0.54	0	15,15,17	1.67	4 (26%)
2	SIA	D	3	1,2	20,20,21	1.34	1 (5%)	24,28,31	1.29	3 (12%)
3	NAG	E	1	3	14,14,15	0.67	0	17,19,21	1.71	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	E	2	3	14,14,15	0.75	0	17,19,21	1.26	3 (17%)
4	NAG	F	1	4	14,14,15	0.68	0	17,19,21	2.00	5 (29%)
4	NAG	F	2	4	14,14,15	0.52	0	17,19,21	0.86	0
4	BMA	F	3	4	11,11,12	0.75	0	15,15,17	0.62	0
2	NAG	G	1	2	15,15,15	1.55	3 (20%)	21,21,21	3.82	15 (71%)
2	GAL	G	2	2	11,11,12	0.77	0	15,15,17	3.10	8 (53%)
2	SIA	G	3	2	20,20,21	1.36	3 (15%)	24,28,31	1.52	5 (20%)
5	NAG	H	1	5	14,14,15	0.55	0	17,19,21	1.62	3 (17%)
5	NAG	H	2	5	14,14,15	0.62	0	17,19,21	0.98	1 (5%)
5	BMA	H	3	5	11,11,12	0.79	0	15,15,17	0.96	1 (6%)
5	MAN	H	4	5	11,11,12	0.82	1 (9%)	15,15,17	1.84	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2	-	0/6/26/26	0/1/1/1
2	GAL	D	2	2	-	0/2/19/22	0/1/1/1
2	SIA	D	3	1,2	-	2/18/34/38	0/1/1/1
3	NAG	E	1	3	-	4/6/23/26	0/1/1/1
3	NAG	E	2	3	-	4/6/23/26	0/1/1/1
4	NAG	F	1	4	-	5/6/23/26	0/1/1/1
4	NAG	F	2	4	-	4/6/23/26	0/1/1/1
4	BMA	F	3	4	-	1/2/19/22	0/1/1/1
2	NAG	G	1	2	-	0/6/26/26	0/1/1/1
2	GAL	G	2	2	-	0/2/19/22	0/1/1/1
2	SIA	G	3	2	-	2/18/34/38	0/1/1/1
5	NAG	H	1	5	-	2/6/23/26	0/1/1/1
5	NAG	H	2	5	-	2/6/23/26	0/1/1/1
5	BMA	H	3	5	-	0/2/19/22	0/1/1/1
5	MAN	H	4	5	-	2/2/19/22	0/1/1/1

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	SIA	C2-C1	4.69	1.56	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	1	NAG	C2-N2	4.20	1.52	1.45
2	G	3	SIA	C4-C5	3.08	1.55	1.53
2	G	3	SIA	O6-C6	-2.80	1.39	1.44
2	G	1	NAG	C8-C7	2.60	1.55	1.50

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	1	NAG	C6-C5-C4	-9.22	91.40	113.00
2	G	2	GAL	C1-O5-C5	-7.10	102.57	112.19
2	G	1	NAG	O5-C1-C2	-6.96	102.52	109.52
2	G	1	NAG	C1-C2-N2	-6.51	103.19	110.73
2	D	1	NAG	O5-C5-C6	5.19	119.33	106.44

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	1	NAG	C4-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6
5	H	4	MAN	C4-C5-C6-O6
5	H	4	MAN	O5-C5-C6-O6


There are no ring outliers.

12 monomers are involved in 31 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	4	MAN	1	0
2	D	1	NAG	1	0
4	F	2	NAG	1	0
2	G	3	SIA	4	0
5	H	1	NAG	5	0
4	F	1	NAG	7	0
2	G	2	GAL	1	0
2	G	1	NAG	2	0
2	D	2	GAL	1	0
3	E	1	NAG	6	0
2	D	3	SIA	4	0
3	E	2	NAG	1	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.




## Oligosaccharide Chain G




## Oligosaccharide Chain E



## Oligosaccharide Chain F





## 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	NAG	A	1500	-	14,14,15	0.69	0	17,19,21	0.73	0
6	NAG	C	1500	-	14,14,15	0.65	0	17,19,21	1.10	1 (5%)
6	NAG	A	1499	-	14,14,15	0.55	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	1500	-	-	5/6/23/26	0/1/1/1
6	NAG	C	1500	-	-	2/6/23/26	0/1/1/1
6	NAG	A	1499	-	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	1500	NAG	C6-C5-C4	-2.35	107.49	113.00
6	A	1499	NAG	C1-O5-C5	2.21	115.19	112.19

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1499	NAG	C1-C2-N2-C7
6	A	1499	NAG	C8-C7-N2-C2
6	A	1499	NAG	O7-C7-N2-C2
6	A	1500	NAG	C1-C2-N2-C7
6	A	1500	NAG	C8-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1500	NAG	5	0
6	C	1500	NAG	3	0
6	A	1499	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	490/509 (96%)	0.00	17 (3%) 47 49	21, 36, 65, 153	1 (0%)
1	B	490/509 (96%)	0.12	26 (5%) 33 35	24, 38, 66, 139	0
1	C	485/509 (95%)	0.08	17 (3%) 47 49	25, 39, 64, 145	0
All	All	1465/1527 (95%)	0.07	60 (4%) 42 44	21, 38, 66, 153	1 (0%)

The worst 5 of 60 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	453	MET	5.7
1	A	74	LEU	5.3
1	B	74	LEU	5.1
1	B	463	GLY	4.9
1	A	459	VAL	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	BMA	F	4	1/12	0.26	0.29	89,89,89,89	0
4	BMA	F	3	11/12	0.63	0.10	88,88,89,89	0
2	NAG	D	1	15/15	0.64	0.13	87,90,92,95	0

*Continued on next page...*

*Continued from previous page...*




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	F	1	14/15	0.64	0.17	86,88,88,88	0
3	NAG	E	2	14/15	0.67	0.12	75,76,76,76	0
2	NAG	G	1	15/15	0.68	0.17	62,65,66,69	0
4	NAG	F	2	14/15	0.77	0.12	87,87,88,88	0
2	GAL	G	2	11/12	0.78	0.11	47,55,58,58	0
5	MAN	H	4	11/12	0.78	0.14	56,56,56,57	0
5	BMA	H	3	11/12	0.79	0.10	55,56,57,57	0
3	NAG	E	1	14/15	0.79	0.15	74,75,75,76	0
5	NAG	H	2	14/15	0.80	0.11	55,56,56,56	0
2	GAL	D	2	11/12	0.81	0.10	73,80,83,83	0
2	SIA	D	3	20/21	0.86	0.12	60,65,72,73	0
5	NAG	H	1	14/15	0.90	0.11	56,57,57,58	0
2	SIA	G	3	20/21	0.93	0.09	35,40,46,48	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



**Electron density around Chain G:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	A	1499	14/15	0.75	0.14	92,94,95,95	0
6	NAG	A	1500	14/15	0.82	0.13	59,61,62,62	0
6	NAG	C	1500	14/15	0.82	0.14	61,63,63,63	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.