



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 08:17 AM EDT

PDB ID : 2X1U  
Title : Crystallographic binding studies with an engineered monomeric variant of triosephosphate isomerase  
Authors : Salin, M.; Kapetanidou, E.G.; Vaismaa, M.; Lajunen, M.; Casteleijn, M.G.; Neubauer, P.; Salmon, L.; Wierenga, R.  
Deposited on : 2010-01-04  
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4003 (1.86-1.82)
Clashscore	141614	4233 (1.86-1.82)
Ramachandran outliers	138981	4185 (1.86-1.82)
Sidechain outliers	138945	4186 (1.86-1.82)
RSRZ outliers	127900	3957 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4061 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called TRIOSEPHOSPHATE ISOMERASE, GLYCOSOMAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	232	Total	C	N	O	S	0	2	0
			1785	1141	312	327	5			
1	B	238	Total	C	N	O	S	0	1	0
			1812	1152	318	338	4			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	15	SER	ASN	engineered mutation	UNP P04789
A	18	PRO	GLN	engineered mutation	UNP P04789
A	19	ASP	GLN	engineered mutation	UNP P04789
A	68	GLY	ILE	engineered mutation	UNP P04789
A	69	ASN	ALA	engineered mutation	UNP P04789
A	70	ALA	LYS	engineered mutation	UNP P04789
A	71	ASP	SER	engineered mutation	UNP P04789
A	72	ALA	GLY	engineered mutation	UNP P04789
A	81	ALA	PRO	engineered mutation	UNP P04789
A	82	SER	ILE	engineered mutation	UNP P04789
A	100	TRP	ALA	engineered mutation	UNP P04789
A	233	ALA	VAL	engineered mutation	UNP P04789
B	15	SER	ASN	engineered mutation	UNP P04789
B	18	PRO	GLN	engineered mutation	UNP P04789
B	19	ASP	GLN	engineered mutation	UNP P04789
B	68	GLY	ILE	engineered mutation	UNP P04789
B	69	ASN	ALA	engineered mutation	UNP P04789
B	70	ALA	LYS	engineered mutation	UNP P04789
B	71	ASP	SER	engineered mutation	UNP P04789
B	72	ALA	GLY	engineered mutation	UNP P04789
B	81	ALA	PRO	engineered mutation	UNP P04789
B	82	SER	ILE	engineered mutation	UNP P04789
B	100	TRP	ALA	engineered mutation	UNP P04789
B	233	ALA	VAL	engineered mutation	UNP P04789

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	196	Total	O	0	0
			196	196		
3	B	238	Total	O	0	0
			238	238		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.67Å 85.38Å 56.30Å 90.00° 98.85° 90.00°	Depositor
Resolution (Å)	8.22 – 1.84 8.22 – 1.84	Depositor EDS
% Data completeness (in resolution range)	99.7 (8.22-1.84) 99.7 (8.22-1.84)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.27 (at 1.84Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.5_2)	Depositor
R, $R_{free}$	0.183 , 0.237 0.179 , 0.233	Depositor DCC
$R_{free}$ test set	1814 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.1	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.49 , 63.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4061	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.15% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.33	0/1826	0.51	0/2478
1	B	0.35	0/1850	0.51	0/2511
All	All	0.34	0/3676	0.51	0/4989

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1785	0	1809	28	0
1	B	1812	0	1833	23	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
3	A	196	0	0	3	0
3	B	238	0	0	5	0
All	All	4061	0	3642	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:12:TRP:HH2	1:A:21:LEU:HD22	1.44	0.82
1:B:13:LYS:HE2	3:B:2009:HOH:O	1.97	0.64
1:A:84:LYS:HZ3	1:A:89:ASN:HD22	1.54	0.55
1:B:51:THR:CG2	1:B:62:ILE:HD13	2.37	0.55
1:A:215:ASN:C	1:A:215:ASN:HD22	2.10	0.55
1:B:215:ASN:ND2	1:B:218:ASN:H	2.03	0.55
1:B:28:PHE:HE2	1:B:238:LEU:HD11	1.72	0.55
1:B:215:ASN:HD22	1:B:215:ASN:C	2.09	0.54
1:A:12:TRP:HB2	1:A:43:SER:HA	1.90	0.54
1:A:215:ASN:ND2	1:A:218:ASN:H	2.05	0.53
1:A:84:LYS:NZ	1:A:89:ASN:HD22	2.06	0.53
1:A:24:LEU:HD23	1:A:238:LEU:HG	1.91	0.53
1:A:12:TRP:CZ2	1:A:47:HIS:CD2	2.98	0.52
1:A:12:TRP:CH2	1:A:21:LEU:HD22	2.35	0.52
1:B:2:SER:HA	3:B:2002:HOH:O	2.09	0.51
1:A:50[A]:MET:HG3	1:A:51:THR:N	2.26	0.50
1:A:84:LYS:NZ	1:A:89:ASN:ND2	2.60	0.49
1:B:84:LYS:HG2	1:B:121:PHE:CE1	2.47	0.49
1:A:141:VAL:HG23	3:A:2118:HOH:O	2.13	0.47
1:B:33:ILE:HB	1:B:59:LYS:HD2	1.97	0.47
1:A:215:ASN:ND2	1:A:217:LYS:HG3	2.29	0.47
1:A:215:ASN:HD21	1:A:218:ASN:H	1.62	0.47
1:B:239:LYS:HB2	1:B:240:PRO:HD2	1.97	0.46
1:A:44:THR:OG1	1:A:47:HIS:HD2	1.98	0.46
1:B:10:ALA:CB	1:B:238:LEU:HD13	2.45	0.46
1:A:167:GLU:HG2	1:A:211:GLY:HA3	1.97	0.46
1:A:12:TRP:CE3	1:A:43:SER:HB2	2.52	0.45
1:A:201:ASP:N	1:A:201:ASP:OD1	2.48	0.45
1:B:10:ALA:HA	1:B:233:ALA:O	2.17	0.45
1:A:10:ALA:HA	1:A:233:ALA:O	2.17	0.45
1:B:201:ASP:O	1:B:205:GLU:HG2	2.17	0.45
1:A:58:PRO:HD3	1:B:59:LYS:HG2	1.98	0.44
1:B:46:VAL:HG13	3:B:2053:HOH:O	2.17	0.44
1:B:215:ASN:HD21	1:B:218:ASN:H	1.66	0.44
1:B:132:GLN:NE2	3:B:2152:HOH:O	2.50	0.44
1:B:10:ALA:HB1	1:B:238:LEU:HD13	1.99	0.44
1:A:193:TRP:CE2	1:A:197:LYS:HG3	2.53	0.44
1:B:127:ILE:HD12	1:B:143:VAL:HG13	1.99	0.43
1:B:239:LYS:HB2	1:B:240:PRO:CD	2.48	0.43
1:A:167:GLU:HG2	1:A:211:GLY:CA	2.48	0.43
1:A:84:LYS:HZ2	1:A:89:ASN:ND2	2.16	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:24:LEU:HD12	1:A:24:LEU:HA	1.87	0.42
1:B:51:THR:HG22	1:B:62:ILE:HD13	2.00	0.42
1:A:180:PRO:HA	1:A:222:LEU:HD23	2.01	0.42
3:A:2043:HOH:O	1:B:201:ASP:HB3	2.20	0.41
1:A:217:LYS:HG3	1:A:217:LYS:H	1.48	0.41
1:B:98:ARG:HA	1:B:98:ARG:HD3	1.86	0.41
1:A:98:ARG:HA	1:A:98:ARG:HD3	1.91	0.41
1:B:250:GLN:NE2	3:B:2235:HOH:O	2.46	0.40
1:A:84:LYS:HD3	3:A:2074:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/238 (97%)	225 (98%)	5 (2%)	0	100	100
1	B	237/238 (100%)	234 (99%)	3 (1%)	0	100	100
All	All	467/476 (98%)	459 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/187 (98%)	182 (99%)	2 (1%)	73	64
1	B	188/187 (100%)	187 (100%)	1 (0%)	88	85
All	All	372/374 (100%)	369 (99%)	3 (1%)	81	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	ASN
1	A	217	LYS
1	B	215	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	HIS
1	A	65	GLN
1	A	66	ASN
1	A	89	ASN
1	A	215	ASN
1	A	224	GLN
1	A	250	GLN
1	B	65	GLN
1	B	89	ASN
1	B	132	GLN
1	B	215	ASN
1	B	250	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1252	-	4,4,4	0.26	0	6,6,6	0.15	0
2	SO4	B	1253	-	4,4,4	0.28	0	6,6,6	0.13	0
2	SO4	B	1252	-	4,4,4	0.18	0	6,6,6	0.24	0
2	SO4	A	1251	-	4,4,4	0.20	0	6,6,6	0.21	0
2	SO4	A	1253	-	4,4,4	0.28	0	6,6,6	0.12	0
2	SO4	B	1251	-	4,4,4	0.33	0	6,6,6	0.07	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	232/238 (97%)	-0.29	5 (2%) 62 60	12, 19, 34, 48	0
1	B	238/238 (100%)	-0.33	4 (1%) 70 69	10, 18, 34, 50	0
All	All	470/476 (98%)	-0.31	9 (1%) 66 65	10, 19, 34, 50	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	SER	6.0
1	B	15	SER	4.9
1	B	2	SER	4.2
1	B	16	GLY	2.8
1	B	12	TRP	2.7
1	A	217	LYS	2.5
1	A	46	VAL	2.1
1	A	45	PHE	2.1
1	A	50[A]	MET	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	B	1253	5/5	0.94	0.23	40,49,53,53	0
2	SO4	A	1253	5/5	0.95	0.17	25,30,35,36	0
2	SO4	A	1252	5/5	0.95	0.11	24,30,34,37	0
2	SO4	A	1251	5/5	0.98	0.06	21,23,25,25	0
2	SO4	B	1252	5/5	0.99	0.04	15,16,20,23	0
2	SO4	B	1251	5/5	0.99	0.07	16,20,22,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.