



wwPDB EM Validation Summary Report ⓘ

Jun 20, 2024 – 07:16 AM JST

PDB ID : 7X6Q
EMDB ID : EMD-33025
Title : cryo-EM structure of human TRiC-ATP-closed state
Authors : Cong, Y.; Liu, C.X.
Deposited on : 2022-03-08
Resolution : 4.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.










Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	 9% 96% ..
1	a	556	 9% 96% ..
2	B	535	 7% 97% .
2	L	535	 6% 97% .
3	E	539	 10% 99% .
3	e	539	 7% 99% .
4	G	545	 9% 97% .
4	N	545	 9% 97% .
5	H	543	 13% 97% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	O	543	
6	I	539	
6	M	539	
7	J	548	
7	P	548	
8	K	531	
8	z	531	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 64472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called T-complex protein 1 subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	537	Total	C	N	O	S	0	0
			4077	2553	713	788	23		
1	a	537	Total	C	N	O	S	0	0
			4077	2553	713	788	23		

- Molecule 2 is a protein called T-complex protein 1 subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	523	Total	C	N	O	S	0	0
			3937	2463	696	759	19		
2	L	523	Total	C	N	O	S	0	0
			3937	2463	696	759	19		

- Molecule 3 is a protein called T-complex protein 1 subunit epsilon.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	535	Total	C	N	O	S	0	0
			4126	2583	718	795	30		
3	e	535	Total	C	N	O	S	0	0
			4126	2583	718	795	30		

- Molecule 4 is a protein called T-complex protein 1 subunit gamma.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	G	526	Total	C	N	O	S	0	0
			4094	2552	727	785	30		
4	N	526	Total	C	N	O	S	0	0
			4094	2552	727	785	30		

- Molecule 5 is a protein called T-complex protein 1 subunit eta.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	H	526	Total	C	N	O	S	0	0
			4037	2549	697	766	25		
5	O	526	Total	C	N	O	S	0	0
			4037	2549	697	766	25		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	290	SER	LEU	engineered mutation	UNP Q99832
O	290	SER	LEU	engineered mutation	UNP Q99832

- Molecule 6 is a protein called T-complex protein 1 subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	I	521	Total	C	N	O	S	0	0
			3935	2459	687	766	23		
6	M	521	Total	C	N	O	S	0	0
			3935	2459	687	766	23		

- Molecule 7 is a protein called T-complex protein 1 subunit theta.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	J	527	Total	C	N	O	S	0	0
			4008	2529	682	770	27		
7	P	527	Total	C	N	O	S	0	0
			4008	2529	682	770	27		


- Molecule 8 is a protein called T-complex protein 1 subunit zeta.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	K	525	Total	C	N	O	S	0	0
			4022	2528	704	769	21		
8	z	525	Total	C	N	O	S	0	0
			4022	2528	704	769	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: T-complex protein 1 subunit alpha



- Molecule 1: T-complex protein 1 subunit alpha




- Molecule 2: T-complex protein 1 subunit beta




- Molecule 2: T-complex protein 1 subunit beta







• Molecule 3: T-complex protein 1 subunit epsilon




• Molecule 3: T-complex protein 1 subunit epsilon




• Molecule 4: T-complex protein 1 subunit gamma




• Molecule 4: T-complex protein 1 subunit gamma







- Molecule 5: T-complex protein 1 subunit eta




- Molecule 5: T-complex protein 1 subunit eta




- Molecule 6: T-complex protein 1 subunit delta




- Molecule 6: T-complex protein 1 subunit delta







• Molecule 7: T-complex protein 1 subunit theta



• Molecule 7: T-complex protein 1 subunit theta




• Molecule 8: T-complex protein 1 subunit zeta



• Molecule 8: T-complex protein 1 subunit zeta





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	41646	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	57	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.055	Depositor
Minimum map value	-0.022	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	320.4, 320.4, 320.4	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.89, 0.89, 0.89	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/4117	0.48	1/5558 (0.0%)
1	a	0.68	2/4117 (0.0%)	0.48	0/5558
2	B	0.68	0/3980	0.47	0/5365
2	L	0.67	0/3980	0.47	0/5365
3	E	0.71	1/4176 (0.0%)	0.46	0/5626
3	e	0.69	0/4176	0.45	0/5626
4	G	0.69	0/4141	0.47	0/5585
4	N	0.70	0/4141	0.46	0/5585
5	H	0.68	1/4094 (0.0%)	0.49	1/5526 (0.0%)
5	O	0.67	3/4094 (0.1%)	0.48	1/5526 (0.0%)
6	I	0.67	2/3967 (0.1%)	0.47	0/5352
6	M	0.67	3/3967 (0.1%)	0.48	1/5352 (0.0%)
7	J	0.69	0/4066	0.48	0/5498
7	P	0.70	1/4066 (0.0%)	0.49	0/5498
8	K	0.70	2/4069 (0.0%)	0.47	0/5486
8	z	0.70	0/4069	0.47	0/5486
All	All	0.68	15/65220 (0.0%)	0.47	4/87992 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	L	0	1
5	O	0	1
6	M	0	1
7	J	0	1
7	P	0	1
All	All	0	6

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	M	106	THR	C-O	-6.96	1.10	1.23
8	K	157	ARG	C-O	-6.55	1.10	1.23
5	O	94	THR	C-O	-6.48	1.11	1.23
6	M	106	THR	CA-C	-6.36	1.36	1.52
1	a	87	GLY	C-O	-6.20	1.13	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	M	105	GLY	C-N-CA	-6.65	105.07	121.70
5	H	91	GLY	N-CA-C	5.81	127.63	113.10
1	A	87	GLY	O-C-N	-5.36	114.12	122.70
5	O	93	GLY	C-N-CA	-5.35	108.33	121.70

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	249	ILE	Peptide
7	J	499	ASP	Peptide
2	L	249	ILE	Peptide
6	M	105	GLY	Mainchain
5	O	93	GLY	Mainchain

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	535/556 (96%)	510 (95%)	23 (4%)	2 (0%)	34 72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	a	535/556 (96%)	511 (96%)	22 (4%)	2 (0%)	34	72
2	B	521/535 (97%)	501 (96%)	19 (4%)	1 (0%)	47	81
2	L	521/535 (97%)	502 (96%)	17 (3%)	2 (0%)	34	72
3	E	533/539 (99%)	513 (96%)	20 (4%)	0	100	100
3	e	533/539 (99%)	513 (96%)	20 (4%)	0	100	100
4	G	524/545 (96%)	515 (98%)	9 (2%)	0	100	100
4	N	524/545 (96%)	514 (98%)	10 (2%)	0	100	100
5	H	524/543 (96%)	506 (97%)	18 (3%)	0	100	100
5	O	524/543 (96%)	505 (96%)	19 (4%)	0	100	100
6	I	519/539 (96%)	510 (98%)	8 (2%)	1 (0%)	47	81
6	M	519/539 (96%)	511 (98%)	7 (1%)	1 (0%)	47	81
7	J	525/548 (96%)	502 (96%)	22 (4%)	1 (0%)	47	81
7	P	525/548 (96%)	501 (95%)	24 (5%)	0	100	100
8	K	523/531 (98%)	507 (97%)	15 (3%)	1 (0%)	47	81
8	z	523/531 (98%)	508 (97%)	13 (2%)	2 (0%)	34	72
All	All	8408/8672 (97%)	8129 (97%)	266 (3%)	13 (0%)	50	81

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	ASN
2	L	21	ALA
2	B	21	ALA
6	I	534	ASP
1	a	90	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	448/463 (97%)	448 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	a	448/463 (97%)	448 (100%)	0	100	100
2	B	416/427 (97%)	416 (100%)	0	100	100
2	L	416/427 (97%)	416 (100%)	0	100	100
3	E	452/455 (99%)	452 (100%)	0	100	100
3	e	452/455 (99%)	452 (100%)	0	100	100
4	G	457/469 (97%)	457 (100%)	0	100	100
4	N	457/469 (97%)	457 (100%)	0	100	100
5	H	433/443 (98%)	433 (100%)	0	100	100
5	O	433/443 (98%)	432 (100%)	1 (0%)	93	96
6	I	443/452 (98%)	443 (100%)	0	100	100
6	M	443/452 (98%)	443 (100%)	0	100	100
7	J	433/452 (96%)	433 (100%)	0	100	100
7	P	433/452 (96%)	433 (100%)	0	100	100
8	K	437/442 (99%)	437 (100%)	0	100	100
8	z	437/442 (99%)	437 (100%)	0	100	100
All	All	7038/7206 (98%)	7037 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	O	32	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
8	K	84	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33025. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


6.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 132




Y Index: 141




Z Index: 160


6.3.2 Raw map



X Index: 132



Y Index: 159




Z Index: 161


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 470 nm³; this corresponds to an approximate mass of 424 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.222 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.222 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.50	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	6.17	7.97	6.65

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 6.17 differs from the reported value 4.5 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33025 and PDB model 7X6Q. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 84% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6470	 0.3040
A	 0.6330	 0.3120
B	 0.6770	 0.3190
E	 0.6570	 0.3130
G	 0.6480	 0.3100
H	 0.6250	 0.2850
I	 0.6650	 0.3220
J	 0.6190	 0.2810
K	 0.6180	 0.2820
L	 0.6890	 0.3230
M	 0.6740	 0.3190
N	 0.6650	 0.3190
O	 0.6360	 0.2860
P	 0.6170	 0.2880
a	 0.6420	 0.3080
e	 0.6650	 0.3140
z	 0.6190	 0.2800

