



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2025 – 12:08 AM JST

PDB ID : 5XD0 / pdb_00005xd0
Title : Apo Structure of Beta-1,3-1,4-glucanase from Paenibacillus sp.X4
Authors : Baek, S.C.; Ho, T.-H.; Kang, L.-W.; Kim, H.
Deposited on : 2017-03-24
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	409	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 85%, green 8%, grey 8%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 85% 7% 8% </div> </div>
1	B	409	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, orange 0%, yellow 84%, green 8%, grey 8%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 84% 8% 8% </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6163 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2930	1841	488	588	13			
1	B	378	Total	C	N	O	S	0	0	0
			2930	1841	488	588	13			

- Molecule 2 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	135	Total	O	0	0
			135	135		
4	B	151	Total	O	0	0
			151	151		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glucanase



• Molecule 1: Glucanase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.72Å 64.16Å 70.50Å 71.04° 77.07° 75.60°	Depositor
Resolution (Å)	65.86 – 1.79 65.86 – 1.79	Depositor EDS
% Data completeness (in resolution range)	96.4 (65.86-1.79) 96.4 (65.86-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.151 , 0.188 0.163 , 0.195	Depositor DCC
R_{free} test set	3509 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 37.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6163	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.02	0/3007	0.97	9/4090 (0.2%)
1	B	1.01	2/3007 (0.1%)	0.97	8/4090 (0.2%)
All	All	1.01	2/6014 (0.0%)	0.97	17/8180 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	56	SER	CB-OG	-5.71	1.34	1.42
1	B	391	GLU	CD-OE1	5.44	1.31	1.25

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	289	ASP	CB-CG-OD1	8.88	126.29	118.30
1	A	204	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	B	211	ASP	CB-CG-OD1	6.41	124.07	118.30
1	B	203	LEU	CB-CG-CD1	6.17	121.50	111.00
1	A	158	ASP	CB-CG-OD2	-6.09	112.82	118.30
1	B	302	ARG	NE-CZ-NH2	-6.01	117.30	120.30
1	B	167	LEU	CB-CG-CD1	5.95	121.12	111.00
1	A	239	ARG	NE-CZ-NH2	-5.71	117.44	120.30
1	A	223	PHE	CB-CG-CD1	5.47	124.63	120.80
1	A	119	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	167	LEU	CB-CG-CD1	5.20	119.84	111.00
1	B	195	ASP	CB-CG-OD1	5.19	122.97	118.30
1	A	204	ARG	NE-CZ-NH2	-5.17	117.71	120.30
1	A	289	ASP	CB-CG-OD2	-5.16	113.66	118.30
1	B	204	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	B	195	ASP	CB-CG-OD2	-5.03	113.77	118.30
1	B	231	PHE	CB-CG-CD2	-5.03	117.28	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2930	0	2746	9	0
1	B	2930	0	2746	12	0
2	A	10	0	14	0	0
3	B	7	0	10	0	0
4	A	135	0	0	0	1
4	B	151	0	0	1	2
All	All	6163	0	5516	21	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:280:SER:HB3	4:B:729:HOH:O	1.41	1.20
1:A:35:LYS:H	1:A:171:GLN:NE2	1.98	0.60
1:B:254:LEU:HD11	1:B:269:VAL:HG21	1.85	0.58
1:B:35:LYS:H	1:B:171:GLN:NE2	2.02	0.57
1:B:254:LEU:HD11	1:B:269:VAL:CG2	2.36	0.56
1:B:40:HIS:HE1	1:B:59:ASP:OD2	1.89	0.56
1:A:35:LYS:H	1:A:171:GLN:HE22	1.57	0.53
1:B:35:LYS:H	1:B:171:GLN:HE22	1.59	0.50
1:B:330:VAL:HB	1:B:336:ASN:HB2	1.96	0.47
1:A:40:HIS:HE1	1:A:59:ASP:OD2	1.98	0.47
1:B:225:LEU:HD13	1:B:248:TYR:CZ	2.51	0.45
1:A:330:VAL:HB	1:A:336:ASN:HB2	1.99	0.45
1:A:225:LEU:HD13	1:A:248:TYR:CZ	2.52	0.45
1:B:262:THR:O	1:B:322:MET:HA	2.17	0.44
1:A:100:GLY:O	1:A:104:THR:HG23	2.19	0.43
1:A:40:HIS:H	1:A:405:ASN:HD22	1.66	0.42
1:A:40:HIS:H	1:A:405:ASN:ND2	2.17	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:THR:O	1:A:322:MET:HA	2.21	0.41
1:B:191:ILE:HG23	1:B:195:ASP:HB2	2.03	0.41
1:B:284:LEU:HD22	1:B:285:GLU:HG3	2.02	0.41
1:B:85:LYS:HG3	1:B:90:GLY:HA2	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:722:HOH:O	4:B:611:HOH:O[1_465]	1.49	0.71
4:B:733:HOH:O	4:B:738:HOH:O[1_455]	1.88	0.32

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/409 (92%)	371 (99%)	5 (1%)	0	100	100
1	B	376/409 (92%)	374 (100%)	2 (0%)	0	100	100
All	All	752/818 (92%)	745 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/339 (93%)	302 (96%)	12 (4%)	28	16
1	B	314/339 (93%)	305 (97%)	9 (3%)	37	26
All	All	628/678 (93%)	607 (97%)	21 (3%)	33	21

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	TYR
1	A	86	TYR
1	A	102	LEU
1	A	125	TYR
1	A	126	LYS
1	A	167	LEU
1	A	203	LEU
1	A	220	PRO
1	A	272	SER
1	A	275	THR
1	A	318	GLN
1	A	389	TYR
1	B	79	THR
1	B	86	TYR
1	B	102	LEU
1	B	167	LEU
1	B	203	LEU
1	B	272	SER
1	B	275	THR
1	B	318	GLN
1	B	389	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	HIS
1	A	132	ASN
1	A	171	GLN
1	A	198	GLN
1	A	200	GLN
1	A	242	ASN
1	A	252	ASN
1	A	288	ASN
1	A	317	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	318	GLN
1	A	336	ASN
1	A	370	ASN
1	A	386	ASN
1	A	405	ASN
1	B	40	HIS
1	B	132	ASN
1	B	171	GLN
1	B	198	GLN
1	B	200	GLN
1	B	226	ASN
1	B	242	ASN
1	B	252	ASN
1	B	318	GLN
1	B	336	ASN
1	B	386	ASN
1	B	405	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PEG	B	501	-	6,6,6	0.51	0	5,5,5	0.57	0
2	PGE	A	501	-	9,9,9	0.80	0	8,8,8	1.39	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	B	501	-	-	4/4/4/4	-
2	PGE	A	501	-	-	2/7/7/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PGE	O3-C5-C6	-2.84	97.60	110.07

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PGE	O3-C5-C6-O4
3	B	501	PEG	C1-C2-O2-C3
3	B	501	PEG	O1-C1-C2-O2
2	A	501	PGE	C4-C3-O2-C2
3	B	501	PEG	O2-C3-C4-O4
3	B	501	PEG	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/409 (92%)	-0.37	4 (1%) 77 77	8, 14, 24, 40	0
1	B	378/409 (92%)	-0.37	1 (0%) 90 90	8, 13, 23, 44	0
All	All	756/818 (92%)	-0.37	5 (0%) 84 84	8, 14, 23, 44	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	32	ALA	5.7
1	A	32	ALA	3.0
1	A	272	SER	2.8
1	A	273	GLY	2.7
1	A	274	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PGE	A	501	10/10	0.82	0.16	21,34,40,42	0
3	PEG	B	501	7/7	0.87	0.14	19,28,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.