



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 14, 2025 – 06:13 PM EDT

PDB ID : 1XT8 / pdb_00001xt8
Title : Crystal Structure of Cysteine-Binding Protein from *Campylobacter jejuni* at 2.0 Å Resolution
Authors : Muller, A.; Thomas, G.H.; Horler, R.; Brannigan, J.A.; Blagova, E.; Levdikov, V.M.; Fogg, M.J.; Wilson, K.S.; Wilkinson, A.J.; Structural Proteomics in Europe (SPINE)
Deposited on : 2004-10-21
Resolution : 2.00 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	292	 76% 8% • 15%
1	B	292	 78% 5% •• 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called putative amino-acid transporter periplasmic solute-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1961	1253	327	377	4			
1	B	251	Total	C	N	O	S	0	0	0
			1983	1266	331	382	4			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	CYS	engineered mutation	GB 15792309
A	261	LEU	-	cloning artifact	GB 15792309
A	262	GLU	-	cloning artifact	GB 15792309
A	263	HIS	-	expression tag	GB 15792309
A	264	HIS	-	expression tag	GB 15792309
A	265	HIS	-	expression tag	GB 15792309
A	266	HIS	-	expression tag	GB 15792309
A	267	HIS	-	expression tag	GB 15792309
A	268	HIS	-	expression tag	GB 15792309
B	1	SER	CYS	engineered mutation	GB 15792309
B	261	LEU	-	cloning artifact	GB 15792309
B	262	GLU	-	cloning artifact	GB 15792309
B	263	HIS	-	expression tag	GB 15792309
B	264	HIS	-	expression tag	GB 15792309
B	265	HIS	-	expression tag	GB 15792309
B	266	HIS	-	expression tag	GB 15792309
B	267	HIS	-	expression tag	GB 15792309
B	268	HIS	-	expression tag	GB 15792309

- Molecule 2 is CYSTEINE (CCD ID: CYS) (formula: C₃H₇NO₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			7	3	1	2	1		
2	A	1	Total	C	N	O	S	0	0
			7	3	1	2	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	163	Total 163	O 163	0	0
4	B	161	Total 161	O 161	0	0

- Molecule 1: putative amino-acid transporter periplasmic solute-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.86Å 88.79Å 72.11Å 90.00° 107.81° 90.00°	Depositor
Resolution (Å)	19.92 – 2.00 19.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.92-2.00) 78.7 (19.92-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 2.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.177 , 0.230 0.222 , 0.265	Depositor DCC
R_{free} test set	1518 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4288	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1996	0.63	0/2692
1	B	0.52	0/2018	0.65	3/2721 (0.1%)
All	All	0.53	0/4014	0.64	3/5413 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	246	ASP	CB-CG-OD2	5.26	123.03	118.30
1	B	247	ASP	CB-CG-OD2	5.25	123.02	118.30
1	B	177	ASP	CB-CG-OD1	5.19	122.97	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	245	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1961	0	1963	22	0
1	B	1983	0	1988	21	0
2	A	14	0	8	2	0
3	B	6	0	8	0	0
4	A	163	0	0	5	0
4	B	161	0	0	1	0
All	All	4288	0	3967	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:GLY:O	1:B:248:VAL:HG13	1.43	1.16
1:B:30:LYS:HZ3	1:B:89:ASN:HD22	1.17	0.88
1:A:197:GLU:OE1	1:A:201:LYS:HE3	1.73	0.87
1:A:64:VAL:HG23	4:A:564:HOH:O	1.79	0.83
1:B:30:LYS:NZ	1:B:89:ASN:HD22	1.82	0.77
1:B:244:PHE:HB3	1:B:248:VAL:HG11	1.69	0.73
1:B:246:ASP:O	1:B:247:ASP:CG	2.30	0.69
1:B:246:ASP:O	1:B:247:ASP:CB	2.43	0.66
1:B:245:GLY:O	1:B:248:VAL:CG1	2.34	0.65
1:A:79:LYS:NZ	1:A:99:GLN:HE22	1.96	0.63
1:A:159:ASN:OD1	1:A:176:HIS:HE1	1.83	0.62
1:B:247:ASP:OD1	1:B:247:ASP:C	2.40	0.60
1:A:21:VAL:HB	4:A:563:HOH:O	2.03	0.59
1:B:247:ASP:OD1	1:B:248:VAL:N	2.36	0.58
1:B:159:ASN:OD1	1:B:176:HIS:HE1	1.87	0.58
1:B:136:LYS:HE2	1:B:157:ASP:OD1	2.06	0.56
2:A:402:CYS:N	1:B:91:THR:HG1	2.03	0.55
1:B:30:LYS:HZ3	1:B:89:ASN:ND2	1.97	0.54
1:A:79:LYS:NZ	1:A:99:GLN:NE2	2.55	0.54
1:A:30:LYS:NZ	1:A:89:ASN:HD22	2.05	0.54
1:B:176:HIS:CD2	1:B:177:ASP:H	2.24	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:176:HIS:CD2	1:A:177:ASP:H	2.29	0.51
1:A:64:VAL:CG2	4:A:564:HOH:O	2.45	0.51
1:A:238:GLU:HB3	4:A:561:HOH:O	2.10	0.51
1:A:79:LYS:HZ1	1:A:99:GLN:HE22	1.59	0.50
1:A:88:ALA:O	1:A:89:ASN:C	2.50	0.50
1:A:91:THR:HG1	2:A:401:CYS:N	2.10	0.50
1:A:103:CYS:HA	1:A:216:LYS:HZ3	1.77	0.49
1:B:117:LYS:HD2	1:B:190:ASP:HA	1.93	0.49
1:A:117:LYS:HD3	1:A:190:ASP:HA	1.95	0.49
1:B:167:MET:HE1	4:B:535:HOH:O	2.13	0.48
1:B:176:HIS:HD2	1:B:177:ASP:H	1.60	0.48
1:A:176:HIS:HD2	1:A:177:ASP:H	1.63	0.47
1:A:23:ARG:HE	1:A:65:GLN:NE2	2.14	0.46
1:B:136:LYS:HB2	1:B:136:LYS:NZ	2.31	0.45
1:B:76:GLU:OE1	1:B:82:LYS:NZ	2.50	0.45
1:B:246:ASP:O	1:B:247:ASP:HB3	2.16	0.44
1:A:30:LYS:HZ3	1:A:89:ASN:HB3	1.83	0.43
1:B:136:LYS:HB2	1:B:136:LYS:HZ3	1.85	0.42
1:A:64:VAL:CB	4:A:564:HOH:O	2.67	0.41
1:A:79:LYS:HZ2	1:A:99:GLN:HE22	1.66	0.41
1:A:142:ALA:O	1:A:146:GLN:HG3	2.19	0.41
1:A:216:LYS:NZ	1:A:220:ASP:OD2	2.48	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/292 (84%)	240 (98%)	5 (2%)	1 (0%)	30	27
1	B	249/292 (85%)	240 (96%)	7 (3%)	2 (1%)	16	12
All	All	495/584 (85%)	480 (97%)	12 (2%)	3 (1%)	22	17

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	89	ASN
1	B	89	ASN
1	B	247	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/244 (87%)	208 (98%)	4 (2%)	52	57
1	B	215/244 (88%)	209 (97%)	6 (3%)	38	40
All	All	427/488 (88%)	417 (98%)	10 (2%)	45	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	VAL
1	A	64	VAL
1	A	95	GLN
1	A	170	ARG
1	B	21	VAL
1	B	95	GLN
1	B	146	GLN
1	B	166	LEU
1	B	246	ASP
1	B	248	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	GLN
1	A	89	ASN
1	A	95	GLN
1	A	99	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	135	ASN
1	A	176	HIS
1	A	188	HIS
1	A	221	ASN
1	A	230	GLN
1	A	233	HIS
1	B	89	ASN
1	B	135	ASN
1	B	150	ASN
1	B	158	GLN
1	B	176	HIS
1	B	188	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	CYS	A	402	-	5,6,6	1.20	1 (20%)	3,7,7	1.73	1 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	CYS	A	401	-	5,6,6	1.01	1 (20%)	3,7,7	1.61	1 (33%)
3	GOL	B	403	-	5,5,5	0.34	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CYS	A	402	-	-	1/6/6/6	-
2	CYS	A	401	-	-	0/6/6/6	-
3	GOL	B	403	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	CYS	OXT-C	-2.57	1.22	1.30
2	A	401	CYS	OXT-C	-2.06	1.24	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	CYS	OXT-C-O	-2.75	117.85	124.08
2	A	402	CYS	OXT-C-O	-2.47	118.47	124.08

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	403	GOL	O1-C1-C2-C3
3	B	403	GOL	O1-C1-C2-O2
2	A	402	CYS	O-C-CA-N

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	402	CYS	1	0
2	A	401	CYS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/292 (84%)	0.24	2 (0%) 82 82	27, 36, 39, 44	10 (4%)
1	B	251/292 (85%)	0.32	8 (3%) 50 48	22, 36, 40, 43	12 (4%)
All	All	499/584 (85%)	0.28	10 (2%) 64 63	22, 36, 40, 44	22 (4%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	247	ASP	4.4
1	B	257	GLY	3.7
1	A	247	ASP	3.1
1	B	249	LYS	3.0
1	A	246	ASP	2.8
1	B	246	ASP	2.5
1	B	252	ASP	2.5
1	B	236	TYR	2.4
1	B	38	GLU	2.2
1	B	254	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	403	6/6	0.83	0.14	60,60,61,61	0
2	CYS	A	402	7/7	0.95	0.07	30,31,32,35	0
2	CYS	A	401	7/7	0.96	0.07	30,30,31,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.