



wwPDB EM Validation Summary Report ⓘ

Nov 4, 2024 – 10:33 AM JST

PDB ID : 7XW5
EMDB ID : EMD-33491
Title : TSHR-thyroid stimulating hormone-Gs-ML109 complex
Authors : Duan, J.; Xu, P.; Luan, X.; Ji, Y.; Yuan, Q.; He, X.; Ye, J.; Cheng, X.; Jiang, H.; Zhang, S.; Jiang, Y.; Xu, H.E.
Deposited on : 2022-05-26
Resolution : 2.96 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev113
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	
2	B	350	
3	G	71	
4	N	134	
5	R	702	
6	X	92	
7	Y	118	

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 12618 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(s) subunit alpha isoforms short.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	235	Total	C	N	O	S	0	0
			1964	1237	358	362	7		

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	226	ALA	GLY	conflict	UNP P63094
A	249	ASP	ALA	conflict	UNP P63094
A	252	ASP	SER	conflict	UNP P63094
A	?	-	MET	deletion	UNP P63094
A	?	-	VAL	deletion	UNP P63094
A	?	-	ILE	deletion	UNP P63094
A	?	-	ARG	deletion	UNP P63094
A	?	-	GLU	deletion	UNP P63094
A	?	-	ASP	deletion	UNP P63094
A	?	-	ASN	deletion	UNP P63094
A	?	-	GLN	deletion	UNP P63094
A	?	-	THR	deletion	UNP P63094
A	?	-	ASN	deletion	UNP P63094
A	356	SER	ALA	conflict	UNP P63094
A	362	ALA	ILE	conflict	UNP P63094
A	365	ILE	VAL	conflict	UNP P63094

- Molecule 2 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	338	Total	C	N	O	S	0	0
			2601	1604	467	509	21		

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	HIS	-	expression tag	UNP P62873
B	-8	HIS	-	expression tag	UNP P62873
B	-7	HIS	-	expression tag	UNP P62873
B	-6	HIS	-	expression tag	UNP P62873
B	-5	HIS	-	expression tag	UNP P62873
B	-4	HIS	-	expression tag	UNP P62873
B	-3	GLY	-	expression tag	UNP P62873
B	-2	SER	-	expression tag	UNP P62873
B	-1	LEU	-	expression tag	UNP P62873
B	0	LEU	-	expression tag	UNP P62873
B	1	GLN	-	expression tag	UNP P62873

- Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	57	Total	C	N	O	S	0	0
			436	273	77	83	3		

- Molecule 4 is a protein called Nanobody35.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	N	128	Total	C	N	O	S	0	0
			973	605	170	192	6		

- Molecule 5 is a protein called Thyrotropin receptor.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	R	572	Total	C	N	O	S	0	0
			4517	2964	725	797	31		

There are 62 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	281	ILE	SER	variant	UNP P16473
R	?	-	ALA	deletion	UNP P16473
R	?	-	LEU	deletion	UNP P16473
R	?	-	ASN	deletion	UNP P16473
R	?	-	SER	deletion	UNP P16473
R	?	-	PRO	deletion	UNP P16473
R	?	-	LEU	deletion	UNP P16473
R	?	-	HIS	deletion	UNP P16473
R	?	-	GLN	deletion	UNP P16473

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	?	-	GLU	deletion	UNP P16473
R	?	-	TYR	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	ASN	deletion	UNP P16473
R	?	-	LEU	deletion	UNP P16473
R	?	-	GLY	deletion	UNP P16473
R	?	-	ASP	deletion	UNP P16473
R	?	-	SER	deletion	UNP P16473
R	?	-	ILE	deletion	UNP P16473
R	?	-	VAL	deletion	UNP P16473
R	?	-	GLY	deletion	UNP P16473
R	?	-	TYR	deletion	UNP P16473
R	?	-	LYS	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	LYS	deletion	UNP P16473
R	?	-	SER	deletion	UNP P16473
R	?	-	LYS	deletion	UNP P16473
R	?	-	PHE	deletion	UNP P16473
R	?	-	GLN	deletion	UNP P16473
R	?	-	ASP	deletion	UNP P16473
R	?	-	THR	deletion	UNP P16473
R	?	-	HIS	deletion	UNP P16473
R	?	-	ASN	deletion	UNP P16473
R	?	-	ASN	deletion	UNP P16473
R	?	-	ALA	deletion	UNP P16473
R	?	-	HIS	deletion	UNP P16473
R	?	-	TYR	deletion	UNP P16473
R	?	-	TYR	deletion	UNP P16473
R	?	-	VAL	deletion	UNP P16473
R	?	-	PHE	deletion	UNP P16473
R	?	-	PHE	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	GLN	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	ASP	deletion	UNP P16473
R	?	-	GLU	deletion	UNP P16473
R	?	-	ILE	deletion	UNP P16473
R	?	-	ILE	deletion	UNP P16473
R	?	-	GLY	deletion	UNP P16473
R	?	-	PHE	deletion	UNP P16473

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
R	367	ASN	GLY	conflict	UNP P16473
R	368	LYS	GLN	conflict	UNP P16473
R	369	THR	GLU	conflict	UNP P16473
R	765	HIS	-	expression tag	UNP P16473
R	766	HIS	-	expression tag	UNP P16473
R	767	HIS	-	expression tag	UNP P16473
R	768	HIS	-	expression tag	UNP P16473
R	769	HIS	-	expression tag	UNP P16473
R	770	HIS	-	expression tag	UNP P16473
R	771	HIS	-	expression tag	UNP P16473
R	772	HIS	-	expression tag	UNP P16473


- Molecule 6 is a protein called Glycoprotein hormones alpha chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	X	87	Total	C	N	O	S	0	0
			670	415	116	126	13		

- Molecule 7 is a protein called Thyrotropin subunit beta.


Mol	Chain	Residues	Atoms					AltConf	Trace
7	Y	105	Total	C	N	O	S	0	0
			838	534	136	154	14		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				AltConf
8	R	1	Total	C	N	O	0
			14	8	1	5	
8	R	1	Total	C	N	O	0
			14	8	1	5	
8	R	1	Total	C	N	O	0
			14	8	1	5	
8	R	1	Total	C	N	O	0
			14	8	1	5	
8	X	1	Total	C	N	O	0
			14	8	1	5	
8	X	1	Total	C	N	O	0
			14	8	1	5	
8	Y	1	Total	C	N	O	0
			14	8	1	5	

- Molecule 9 is {N}-[4-[[2-methoxy-5-[(2 {S})-5-oxidanyl-4-oxidanylidene-3-(phenylmethyl)-1,2-dihydroquinazolin-2-yl]phenyl]methoxy]phenyl]ethanamide (three-letter code: HOI) (formula: C₃₁H₂₉N₃O₅) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
9	R	1	Total	C	N	O	0
			39	31	3	5	

- Molecule 10 is CHOLESTEROL (three-letter code: CLR) (formula: C₂₇H₄₆O).



Mol	Chain	Residues	Atoms			AltConf
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	
10	R	1	Total	C	O	0
			28	27	1	

- Molecule 11 is PALMITIC ACID (three-letter code: PLM) (formula: C₁₆H₃₂O₂).




Mol	Chain	Residues	Atoms			AltConf
11	R	1	Total	C	O	0
			18	16	2	
11	R	1	Total	C	O	0
			18	16	2	
11	R	1	Total	C	O	0
			18	16	2	
11	R	1	Total	C	O	0
			18	16	2	
11	R	1	Total	C	O	0
			18	16	2	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide-binding protein G(s) subunit alpha isoforms short




- Molecule 2: Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1



- Molecule 3: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



- Molecule 4: Nanobody35



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	565098	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	60	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.753	Depositor
Minimum map value	-3.959	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.093	Depositor
Recommended contour level	0.6	Depositor
Map size (\AA)	321.3, 321.3, 321.3	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.071, 1.071, 1.071	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLM, NAG, HOI, CLR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2002	1.06	8/2691 (0.3%)
2	B	0.72	4/2648 (0.2%)	1.14	14/3589 (0.4%)
3	G	0.61	0/442	0.98	1/597 (0.2%)
4	N	0.71	0/993	1.04	2/1345 (0.1%)
5	R	0.63	0/4625	0.97	6/6295 (0.1%)
6	X	0.67	0/686	0.97	1/926 (0.1%)
7	Y	0.72	0/860	0.97	0/1169
All	All	0.67	4/12256 (0.0%)	1.03	32/16612 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
2	B	0	1
6	X	0	1
All	All	0	4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	105	TYR	CA-C	-7.16	1.34	1.52
2	B	104	ALA	CA-CB	-5.58	1.40	1.52
2	B	104	ALA	C-O	-5.32	1.13	1.23
2	B	105	TYR	CE2-CZ	-5.09	1.31	1.38

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	27	ARG	NE-CZ-NH2	8.73	124.66	120.30
1	A	332	ARG	NE-CZ-NH2	8.58	124.59	120.30
2	B	48	ARG	NE-CZ-NH2	8.45	124.53	120.30
5	R	274	ARG	NE-CZ-NH2	8.29	124.44	120.30
2	B	219	ARG	NE-CZ-NH2	8.15	124.38	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	232	ARG	Sidechain
1	A	348	TYR	Sidechain
2	B	314	ARG	Sidechain
6	X	54	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1964	0	1940	13	0
2	B	2601	0	2505	15	0
3	G	436	0	448	0	0
4	N	973	0	938	0	0
5	R	4517	0	4579	25	0
6	X	670	0	637	0	0
7	Y	838	0	798	2	0
8	R	56	0	52	0	0
8	X	28	0	26	0	0
8	Y	14	0	13	0	0
9	R	39	0	0	0	0
10	R	392	0	644	16	0
11	R	90	0	155	6	0
All	All	12618	0	12735	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:R:811:CLR:H212	10:R:811:CLR:H183	1.37	1.04
1:A:63:LEU:HD13	1:A:214:VAL:HG22	1.47	0.94
1:A:40:THR:HG22	1:A:218:ASN:OD1	1.67	0.92
5:R:144:LEU:CD1	5:R:156:LEU:HD21	2.00	0.92
5:R:144:LEU:HD13	5:R:156:LEU:HD21	1.53	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/249 (93%)	226 (98%)	5 (2%)	0	100	100
2	B	336/350 (96%)	319 (95%)	17 (5%)	0	100	100
3	G	55/71 (78%)	55 (100%)	0	0	100	100
4	N	126/134 (94%)	123 (98%)	3 (2%)	0	100	100
5	R	566/702 (81%)	535 (94%)	31 (6%)	0	100	100
6	X	85/92 (92%)	83 (98%)	2 (2%)	0	100	100
7	Y	103/118 (87%)	92 (89%)	11 (11%)	0	100	100
All	All	1502/1716 (88%)	1433 (95%)	69 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/221 (96%)	213 (100%)	0	100	100
2	B	281/292 (96%)	278 (99%)	3 (1%)	70	83
3	G	46/58 (79%)	46 (100%)	0	100	100
4	N	106/112 (95%)	106 (100%)	0	100	100
5	R	504/637 (79%)	504 (100%)	0	100	100
6	X	79/83 (95%)	79 (100%)	0	100	100
7	Y	95/107 (89%)	95 (100%)	0	100	100
All	All	1324/1510 (88%)	1321 (100%)	3 (0%)	91	96

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	59	TYR
2	B	105	TYR
2	B	292	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
10	CLR	R	812	-	31,31,31	1.16	2 (6%)	48,48,48	0.59	0
11	PLM	R	821	-	17,17,17	0.95	1 (5%)	17,17,17	0.59	0
11	PLM	R	824	-	17,17,17	0.88	1 (5%)	17,17,17	0.62	0
10	CLR	R	809	-	31,31,31	1.22	2 (6%)	48,48,48	0.78	0
10	CLR	R	814	-	31,31,31	1.12	1 (3%)	48,48,48	0.84	1 (2%)
10	CLR	R	810	-	31,31,31	1.13	2 (6%)	48,48,48	0.70	1 (2%)
8	NAG	X	201	6	14,14,15	1.23	1 (7%)	17,19,21	0.80	0
9	HOI	R	805	-	42,43,43	0.90	0	53,60,60	0.95	6 (11%)
8	NAG	R	802	5	14,14,15	1.22	1 (7%)	17,19,21	0.85	0
10	CLR	R	815	-	31,31,31	1.12	1 (3%)	48,48,48	0.43	0
8	NAG	R	801	5	14,14,15	1.23	2 (14%)	17,19,21	0.77	0
8	NAG	Y	201	7	14,14,15	1.33	3 (21%)	17,19,21	0.89	1 (5%)
10	CLR	R	816	-	31,31,31	0.80	1 (3%)	48,48,48	1.50	5 (10%)
8	NAG	R	803	5	14,14,15	1.35	3 (21%)	17,19,21	0.79	0
10	CLR	R	818	-	31,31,31	1.15	1 (3%)	48,48,48	0.46	0
11	PLM	R	823	-	17,17,17	0.60	1 (5%)	17,17,17	0.88	0
11	PLM	R	822	-	17,17,17	0.96	1 (5%)	17,17,17	0.53	0
10	CLR	R	806	-	31,31,31	1.05	1 (3%)	48,48,48	0.64	0
10	CLR	R	811	-	31,31,31	0.78	1 (3%)	48,48,48	1.43	7 (14%)
11	PLM	R	820	-	17,17,17	0.90	1 (5%)	17,17,17	0.63	0
10	CLR	R	808	-	31,31,31	1.20	2 (6%)	48,48,48	0.72	1 (2%)
10	CLR	R	813	-	31,31,31	1.18	2 (6%)	48,48,48	0.62	0
8	NAG	X	202	6	14,14,15	1.22	2 (14%)	17,19,21	0.82	1 (5%)
10	CLR	R	819	-	31,31,31	1.21	3 (9%)	48,48,48	0.60	0
10	CLR	R	817	-	31,31,31	1.17	1 (3%)	48,48,48	0.60	0
8	NAG	R	804	5	14,14,15	1.19	2 (14%)	17,19,21	0.78	0
10	CLR	R	807	-	31,31,31	1.07	2 (6%)	48,48,48	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	CLR	R	812	-	-	1/10/68/68	0/4/4/4
11	PLM	R	821	-	-	6/15/15/15	-
11	PLM	R	824	-	-	7/15/15/15	-
10	CLR	R	809	-	-	0/10/68/68	0/4/4/4
10	CLR	R	814	-	-	0/10/68/68	0/4/4/4
10	CLR	R	810	-	-	4/10/68/68	0/4/4/4
8	NAG	X	201	6	-	0/6/23/26	0/1/1/1
9	HOI	R	805	-	-	4/19/35/35	0/5/5/5
8	NAG	R	802	5	-	0/6/23/26	0/1/1/1
10	CLR	R	815	-	-	0/10/68/68	0/4/4/4
8	NAG	R	801	5	-	0/6/23/26	0/1/1/1
8	NAG	Y	201	7	-	0/6/23/26	0/1/1/1
10	CLR	R	816	-	-	3/10/68/68	0/4/4/4
8	NAG	R	803	5	-	0/6/23/26	0/1/1/1
10	CLR	R	818	-	-	0/10/68/68	0/4/4/4
11	PLM	R	823	-	-	9/15/15/15	-
11	PLM	R	822	-	-	4/15/15/15	-
10	CLR	R	806	-	-	2/10/68/68	0/4/4/4
10	CLR	R	811	-	-	7/10/68/68	0/4/4/4
11	PLM	R	820	-	-	4/15/15/15	-
10	CLR	R	808	-	-	2/10/68/68	0/4/4/4
10	CLR	R	813	-	-	2/10/68/68	0/4/4/4
8	NAG	X	202	6	-	0/6/23/26	0/1/1/1
10	CLR	R	819	-	-	0/10/68/68	0/4/4/4
10	CLR	R	817	-	-	3/10/68/68	0/4/4/4
8	NAG	R	804	5	-	0/6/23/26	0/1/1/1
10	CLR	R	807	-	-	0/10/68/68	0/4/4/4

The worst 5 of 41 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	R	808	CLR	C20-C17	2.79	1.59	1.54
8	R	804	NAG	O5-C5	2.79	1.49	1.43
8	X	201	NAG	O5-C5	2.77	1.49	1.43
8	R	803	NAG	O5-C5	2.75	1.49	1.43
10	R	809	CLR	C11-C9	2.67	1.58	1.53

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	R	816	CLR	C13-C17-C20	-4.72	112.09	119.49
10	R	811	CLR	C13-C17-C20	-4.11	113.05	119.49
10	R	816	CLR	C13-C14-C8	-3.93	108.56	114.38
10	R	811	CLR	C13-C14-C8	-3.90	108.61	114.38
10	R	816	CLR	C17-C13-C14	3.36	104.05	100.07

There are no chirality outliers.

5 of 58 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	R	811	CLR	C17-C20-C22-C23
11	R	823	PLM	C9-CA-CB-CC
10	R	817	CLR	C17-C20-C22-C23
11	R	823	PLM	C4-C5-C6-C7
10	R	808	CLR	C20-C22-C23-C24

There are no ring outliers.


5 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	R	810	CLR	2	0
10	R	816	CLR	6	0
11	R	823	PLM	6	0
10	R	811	CLR	10	0
10	R	817	CLR	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand CLR R 809




Ligand CLR R 814





Ligand CLR R 810






Ligand HOI R 805












Ligand CLR R 813




Ligand CLR R 819



Ligand CLR R 817





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-33491. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 160




Y Index: 159




Z Index: 106


6.3.2 Raw map



X Index: 151



Y Index: 149




Z Index: 154


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.6. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 68 nm^3 ; this corresponds to an approximate mass of 61 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.338 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.338 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.96	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.44	3.90	3.51

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.44 differs from the reported value 2.96 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-33491 and PDB model 7XW5. Per-residue inclusion information can be found in [section 3](#) on [page 10](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.6 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.6).

9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.6) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.6810	<div></div> 0.4820
A	<div></div> 0.7680	<div></div> 0.5100
B	<div></div> 0.7940	<div></div> 0.5130
G	<div></div> 0.4870	<div></div> 0.4210
N	<div></div> 0.2950	<div></div> 0.4190
R	<div></div> 0.7580	<div></div> 0.5170
X	<div></div> 0.4160	<div></div> 0.3270
Y	<div></div> 0.4400	<div></div> 0.3440

1.0

0.0

<0.0