



## wwPDB EM Validation Summary Report ⓘ

Jul 2, 2024 – 10:17 PM JST

PDB ID : 8XXM  
EMDB ID : EMD-38753  
Title : Cryo-EM structure of the human 40S ribosome with PDCD4 and eIF3G  
Authors : Ye, X.; Huang, Z.; Li, Y.; Wang, M.; Cheng, J.  
Deposited on : 2024-01-18  
Resolution : 3.20 Å(reported)  
Based on initial model : 6ZVJ

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Ln	25	 96%
2	S2	1869	 53% 32% 6% 8%
3	SA	295	 63% 12% 25%
4	SB	264	 64% 17% 19%
5	SD	243	 75% 18% 7%
6	SE	263	 82% 18% 7%
7	SF	204	 71% 22% 7%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	SH	194	
9	SI	208	
10	SK	165	
11	SL	158	
12	SP	145	
13	SQ	146	
14	SR	135	
15	SS	152	
16	ST	145	
17	SU	119	
18	SV	83	
19	SX	143	
20	Sa	115	
21	Sc	69	
22	Sd	56	
23	Sg	317	
24	SC	293	
25	SG	249	
26	SJ	194	
27	SM	132	
28	SN	151	
29	SO	151	
30	SW	130	
31	SY	133	
32	SZ	125	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
33	Sb	84	<div><div></div><div>99%</div><div></div></div>
34	Se	59	<div><div></div><div>98%</div><div></div></div>
35	Sf	156	<div><div><div>24%</div><div>43%</div></div><div>57%</div><div></div></div>
36	CD	469	<div><div><div>10%</div><div></div></div><div>90%</div><div></div></div>
37	3G	320	<div><div><div></div><div>22%</div></div><div>74%</div><div></div></div>

## 2 Entry composition

There are 39 unique types of molecules in this entry. The entry contains 76934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 60S ribosomal protein L41.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Ln	24	Total	C	N	O	S	0	0
			230	139	62	26	3		

- Molecule 2 is a RNA chain called 18S rRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	S2	1723	Total	C	N	O	P	0	0
			36538	16298	6533	11984	1723		

- Molecule 3 is a protein called 40S ribosomal protein SA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	SA	221	Total	C	N	O	S	0	0
			1741	1106	305	322	8		

- Molecule 4 is a protein called 40S ribosomal protein S3a.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	SB	214	Total	C	N	O	S	0	0
			1738	1103	310	311	14		

- Molecule 5 is a protein called 40S ribosomal protein S3.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	SD	227	Total	C	N	O	S	0	0
			1765	1125	317	315	8		

- Molecule 6 is a protein called 40S ribosomal protein S4, X isoform.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	SE	262	Total	C	N	O	S	0	0
			2076	1324	386	358	8		

- Molecule 7 is a protein called 40S ribosomal protein S5.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	SF	189	Total	C	N	O	S	0	0
			1495	934	284	270	7		

- Molecule 8 is a protein called 40S ribosomal protein S7.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	SH	186	Total	C	N	O	S	0	0
			1497	956	274	266	1		

- Molecule 9 is a protein called 40S ribosomal protein S8.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	SI	206	Total	C	N	O	S	0	0
			1686	1058	332	291	5		

- Molecule 10 is a protein called 40S ribosomal protein S10.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	SK	98	Total	C	N	O	S	0	0
			827	539	148	134	6		

- Molecule 11 is a protein called 40S ribosomal protein S11.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	SL	153	Total	C	N	O	S	0	0
			1247	793	234	214	6		

- Molecule 12 is a protein called 40S ribosomal protein S15.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	SP	121	Total	C	N	O	S	0	0
			985	623	185	170	7		

- Molecule 13 is a protein called 40S ribosomal protein S16.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	SQ	144	Total	C	N	O	S	0	0
			1142	726	216	197	3		

- Molecule 14 is a protein called 40S ribosomal protein S17.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	SR	135	Total	C	N	O	S	0	0
			1090	685	202	198	5		

- Molecule 15 is a protein called 40S ribosomal protein S18.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	SS	145	Total	C	N	O	S	0	0
			1198	751	242	203	2		

- Molecule 16 is a protein called 40S ribosomal protein S19.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	ST	143	Total	C	N	O	S	0	0
			1112	697	214	198	3		

- Molecule 17 is a protein called 40S ribosomal protein S20.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	SU	104	Total	C	N	O	S	0	0
			821	514	155	148	4		

- Molecule 18 is a protein called 40S ribosomal protein S21.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	SV	83	Total	C	N	O	S	0	0
			636	393	117	121	5		

- Molecule 19 is a protein called 40S ribosomal protein S23.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	SX	141	Total	C	N	O	S	0	0
			1098	693	219	183	3		

- Molecule 20 is a protein called 40S ribosomal protein S26.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	Sa	102	Total	C	N	O	S	0	0
			821	512	171	133	5		

- Molecule 21 is a protein called 40S ribosomal protein S28.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	Sc	64	Total	C	N	O	S	0	0
			506	308	102	94	2		

- Molecule 22 is a protein called 40S ribosomal protein S29.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	Sd	55	Total	C	N	O	S	0	0
			459	286	94	74	5		

- Molecule 23 is a protein called Receptor of activated protein C kinase 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	Sg	313	Total	C	N	O	S	0	0
			2436	1535	424	465	12		

- Molecule 24 is a protein called 40S ribosomal protein S2.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	SC	222	Total	C	N	O	S	0	0
			1725	1115	298	302	10		

- Molecule 25 is a protein called 40S ribosomal protein S6.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	SG	237	Total	C	N	O	S	0	0
			1923	1200	387	329	7		

- Molecule 26 is a protein called 40S ribosomal protein S9.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	SJ	185	Total	C	N	O	S	0	0
			1525	969	306	248	2		

- Molecule 27 is a protein called 40S ribosomal protein S12.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	SM	122	Total	C	N	O	S	0	0
			940	590	164	177	9		

- Molecule 28 is a protein called 40S ribosomal protein S13.



Mol	Chain	Residues	Atoms					AltConf	Trace
28	SN	150	Total	C	N	O	S	0	0
			1208	773	229	205	1		

- Molecule 29 is a protein called 40S ribosomal protein S14.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	SO	140	Total	C	N	O	S	0	0
			1049	642	204	197	6		

- Molecule 30 is a protein called 40S ribosomal protein S15a.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	SW	129	Total	C	N	O	S	0	0
			1034	659	193	176	6		

- Molecule 31 is a protein called 40S ribosomal protein S24.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	SY	131	Total	C	N	O	S	0	0
			1065	673	209	178	5		

- Molecule 32 is a protein called 40S ribosomal protein S25.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	SZ	75	Total	C	N	O	S	0	0
			598	382	111	104	1		

- Molecule 33 is a protein called 40S ribosomal protein S27.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	Sb	83	Total	C	N	O	S	0	0
			651	408	121	115	7		

- Molecule 34 is a protein called 40S ribosomal protein S30.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	Se	58	Total	C	N	O	S	0	0
			459	284	100	74	1		

- Molecule 35 is a protein called Ubiquitin-40S ribosomal protein S27a.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	Sf	67	Total	C	N	O	S	0	0
			548	346	102	93	7		

- Molecule 36 is a protein called Programmed cell death protein 4.

Mol	Chain	Residues	Atoms				AltConf	Trace
36	CD	49	Total	C	N	O	0	0
			372	225	70	77		

- Molecule 37 is a protein called Eukaryotic translation initiation factor 3 subunit G.

Mol	Chain	Residues	Atoms				AltConf	Trace
37	3G	84	Total	C	N	O	0	0
			667	418	120	129		

- Molecule 38 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
38	S2	22	Total	Mg	0
			22	22	
38	SG	1	Total	Mg	0
			1	1	

- Molecule 39 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
39	Sa	1	Total	Zn	0
			1	1	
39	Sd	1	Total	Zn	0
			1	1	
39	Sf	1	Total	Zn	0
			1	1	

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 60S ribosomal protein L41

Chain Ln:  96%




- Molecule 2: 18S rRNA

Chain S2:  53% 32% 6% 8%









- Molecule 4: 40S ribosomal protein S3a

Chain SB: 64% 17% 19%




- Molecule 5: 40S ribosomal protein S3

Chain SD: 75% 18% 7%



- Molecule 6: 40S ribosomal protein S4, X isoform


Chain SE: 82% 18%





- Molecule 7: 40S ribosomal protein S5

Chain SF: 71% 22% 7%








• Molecule 8: 40S ribosomal protein S7




• Molecule 9: 40S ribosomal protein S8




• Molecule 10: 40S ribosomal protein S10



• Molecule 11: 40S ribosomal protein S11




• Molecule 12: 40S ribosomal protein S15




PHE  
ILE  
PRO  
LEU  
LYS

- Molecule 13: 40S ribosomal protein S16

Chain SQ:  82% 16% ..



- Molecule 14: 40S ribosomal protein S17

Chain SR:  79% 20% .




- Molecule 15: 40S ribosomal protein S18

Chain SS:  73% 22% • 5%




GLY  
VAL  
SER  
LYS  
LYS  
LYS


- Molecule 16: 40S ribosomal protein S19

Chain ST:  78% 21% .




- Molecule 17: 40S ribosomal protein S20

Chain SU:  77% 10% 13%



- Molecule 18: 40S ribosomal protein S21


Chain SV:  80% 20%




- Molecule 19: 40S ribosomal protein S23

MIT	G2	R11	R14	Q20	L52	K60	N63	K68	Q73	N77	G78	K79	L91	V100	F105	G106	R107	H110	D114	K121	K124	V125	A126	N127	V128	S129	K138	R142	SFR
-----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-----

- Chain Sa:  86% 11%




- Chain Sc:  91% 7%




- Chain Sd:  96%



- Chain Sg:  98%




- Chain SC: 




- WORLDWIDE  
**PDB**  
PROTEIN DATA BANK




Chain SG:  71% 23% 5%




- Molecule 26: 40S ribosomal protein S9

Chain SJ:  81% 14% 5%




- Molecule 27: 40S ribosomal protein S12

Chain SM:  30% 67% 23% 8%





- Molecule 28: 40S ribosomal protein S13

Chain SN:  86% 13% 1%




- Molecule 29: 40S ribosomal protein S14

Chain SO:  75% 18% 7%



- Molecule 30: 40S ribosomal protein S15a

Chain SW:  87% 12% 1%





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	68929	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	58	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.397	Depositor
Minimum map value	-0.173	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	446.88, 446.88, 446.88	wwPDB
Map dimensions	420, 420, 420	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.064, 1.064, 1.064	Depositor

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	Ln	0.28	0/231	0.74	0/294
2	S2	0.51	0/40843	0.99	127/63639 (0.2%)
3	SA	0.33	0/1778	0.62	2/2416 (0.1%)
4	SB	0.32	0/1765	0.61	1/2362 (0.0%)
5	SD	0.34	0/1793	0.62	0/2414
6	SE	0.33	0/2118	0.62	1/2849 (0.0%)
7	SF	0.34	0/1516	0.71	2/2037 (0.1%)
8	SH	0.30	0/1519	0.65	0/2033
9	SI	0.34	0/1715	0.70	1/2287 (0.0%)
10	SK	0.33	0/851	0.64	0/1147
11	SL	0.35	0/1268	0.58	0/1696
12	SP	0.30	0/1003	0.68	0/1342
13	SQ	0.36	0/1160	0.73	1/1553 (0.1%)
14	SR	0.29	0/1105	0.66	1/1484 (0.1%)
15	SS	0.29	0/1216	0.68	1/1628 (0.1%)
16	ST	0.31	0/1131	0.58	0/1515
17	SU	0.30	0/831	0.69	0/1115
18	SV	0.34	0/643	0.63	0/860
19	SX	0.38	0/1116	0.64	0/1490
20	Sa	0.36	0/836	0.70	2/1121 (0.2%)
21	Sc	0.32	0/508	0.87	1/680 (0.1%)
22	Sd	0.37	0/470	0.67	0/623
23	Sg	0.28	0/2493	0.67	1/3394 (0.0%)
24	SC	0.38	0/1762	0.64	0/2381
25	SG	0.30	0/1946	0.70	2/2590 (0.1%)
26	SJ	0.35	0/1550	0.63	0/2069
27	SM	0.29	0/950	0.63	0/1275
28	SN	0.31	0/1232	0.57	0/1656
29	SO	0.32	0/1062	0.63	0/1425
30	SW	0.35	0/1051	0.65	1/1406 (0.1%)
31	SY	0.35	0/1083	0.65	0/1438
32	SZ	0.29	0/604	0.78	0/810

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
33	Sb	0.30	0/665	0.62	0/891
34	Se	0.30	0/465	0.61	0/612
35	Sf	0.29	0/560	0.65	0/745
36	CD	0.28	0/378	0.60	0/506
37	3G	0.28	0/680	0.69	0/916
All	All	0.43	0/81897	0.85	144/118699 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
14	SR	0	1
19	SX	0	1
26	SJ	0	1
32	SZ	0	1
All	All	0	4

There are no bond length outliers.

The worst 5 of 144 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S2	1772	C	N1-C2-O2	13.51	127.01	118.90
2	S2	1772	C	N3-C2-O2	-12.94	112.84	121.90
2	S2	501	C	N1-C2-O2	12.07	126.14	118.90
2	S2	501	C	C2-N1-C1'	11.93	131.92	118.80
2	S2	293	C	N1-C2-O2	11.78	125.97	118.90

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
26	SJ	137	VAL	Peptide
14	SR	67	ARG	Peptide
19	SX	126	ALA	Peptide
32	SZ	46	ASN	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Ln	230	0	276	0	0
2	S2	36538	0	18415	329	0
3	SA	1741	0	1746	25	0
4	SB	1738	0	1809	29	0
5	SD	1765	0	1865	30	0
6	SE	2076	0	2177	27	0
7	SF	1495	0	1549	29	0
8	SH	1497	0	1590	17	0
9	SI	1686	0	1772	35	0
10	SK	827	0	854	14	0
11	SL	1247	0	1323	15	0
12	SP	985	0	1031	15	0
13	SQ	1142	0	1213	18	0
14	SR	1090	0	1149	16	0
15	SS	1198	0	1261	24	0
16	ST	1112	0	1146	20	0
17	SU	821	0	883	7	0
18	SV	636	0	637	14	0
19	SX	1098	0	1167	15	0
20	Sa	821	0	870	0	0
21	Sc	506	0	536	0	0
22	Sd	459	0	448	0	0
23	Sg	2436	0	2393	0	0
24	SC	1725	0	1813	31	0
25	SG	1923	0	2088	43	0
26	SJ	1525	0	1640	17	0
27	SM	940	0	965	19	0
28	SN	1208	0	1294	15	0
29	SO	1049	0	1073	24	0
30	SW	1034	0	1080	10	0
31	SY	1065	0	1142	20	0
32	SZ	598	0	656	11	0
33	Sb	651	0	672	0	0
34	Se	459	0	503	0	0
35	Sf	548	0	555	0	0
36	CD	372	0	352	5	0
37	3G	667	0	647	13	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
38	S2	22	0	0	0	0
38	SG	1	0	0	0	0
39	Sa	1	0	0	0	0
39	Sd	1	0	0	0	0
39	Sf	1	0	0	0	0
All	All	76934	0	60590	746	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 746 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S2:886:A:N6	2:S2:901:G:C4	2.22	1.08
2:S2:1756:C:N4	2:S2:1775:U:H3	1.63	0.96
2:S2:543:C:H1'	37:3G:274:LYS:NZ	1.83	0.93
2:S2:140:C:N4	2:S2:313:A:H61	1.66	0.92
2:S2:885:U:H3	2:S2:901:G:H1	1.12	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Ln	22/25 (88%)	22 (100%)	0	0	100	100
3	SA	219/295 (74%)	205 (94%)	14 (6%)	0	100	100
4	SB	212/264 (80%)	195 (92%)	17 (8%)	0	100	100
5	SD	225/243 (93%)	212 (94%)	12 (5%)	1 (0%)	34	69
6	SE	260/263 (99%)	250 (96%)	10 (4%)	0	100	100

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
7	SF	187/204 (92%)	169 (90%)	17 (9%)	1 (0%)	29	67
8	SH	182/194 (94%)	163 (90%)	17 (9%)	2 (1%)	14	51
9	SI	204/208 (98%)	185 (91%)	18 (9%)	1 (0%)	29	67
10	SK	96/165 (58%)	88 (92%)	8 (8%)	0	100	100
11	SL	151/158 (96%)	141 (93%)	10 (7%)	0	100	100
12	SP	119/145 (82%)	117 (98%)	2 (2%)	0	100	100
13	SQ	142/146 (97%)	130 (92%)	10 (7%)	2 (1%)	11	46
14	SR	133/135 (98%)	122 (92%)	10 (8%)	1 (1%)	19	58
15	SS	143/152 (94%)	134 (94%)	9 (6%)	0	100	100
16	ST	141/145 (97%)	134 (95%)	7 (5%)	0	100	100
17	SU	102/119 (86%)	92 (90%)	10 (10%)	0	100	100
18	SV	81/83 (98%)	76 (94%)	5 (6%)	0	100	100
19	SX	139/143 (97%)	129 (93%)	9 (6%)	1 (1%)	22	61
20	Sa	100/115 (87%)	92 (92%)	7 (7%)	1 (1%)	15	54
21	Sc	62/69 (90%)	48 (77%)	14 (23%)	0	100	100
22	Sd	53/56 (95%)	45 (85%)	7 (13%)	1 (2%)	8	39
23	Sg	311/317 (98%)	274 (88%)	37 (12%)	0	100	100
24	SC	220/293 (75%)	208 (94%)	11 (5%)	1 (0%)	29	67
25	SG	235/249 (94%)	221 (94%)	14 (6%)	0	100	100
26	SJ	183/194 (94%)	176 (96%)	7 (4%)	0	100	100
27	SM	120/132 (91%)	115 (96%)	5 (4%)	0	100	100
28	SN	148/151 (98%)	143 (97%)	5 (3%)	0	100	100
29	SO	138/151 (91%)	131 (95%)	7 (5%)	0	100	100
30	SW	127/130 (98%)	123 (97%)	4 (3%)	0	100	100
31	SY	129/133 (97%)	118 (92%)	11 (8%)	0	100	100
32	SZ	73/125 (58%)	56 (77%)	17 (23%)	0	100	100
33	Sb	81/84 (96%)	70 (86%)	11 (14%)	0	100	100
34	Se	56/59 (95%)	53 (95%)	3 (5%)	0	100	100
35	Sf	65/156 (42%)	59 (91%)	6 (9%)	0	100	100
36	CD	47/469 (10%)	41 (87%)	6 (13%)	0	100	100
37	3G	82/320 (26%)	81 (99%)	1 (1%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	4988/6290 (79%)	4618 (93%)	358 (7%)	12 (0%)	50 79

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
5	SD	178	ARG
9	SI	160	SER
19	SX	127	ASN
8	SH	15	LYS
13	SQ	43	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Ln	23/24 (96%)	23 (100%)	0	100 100
3	SA	183/243 (75%)	183 (100%)	0	100 100
4	SB	195/231 (84%)	195 (100%)	0	100 100
5	SD	190/202 (94%)	189 (100%)	1 (0%)	88 95
6	SE	224/225 (100%)	224 (100%)	0	100 100
7	SF	159/170 (94%)	159 (100%)	0	100 100
8	SH	166/174 (95%)	164 (99%)	2 (1%)	71 88
9	SI	178/180 (99%)	178 (100%)	0	100 100
10	SK	89/136 (65%)	89 (100%)	0	100 100
11	SL	137/142 (96%)	136 (99%)	1 (1%)	84 94
12	SP	107/130 (82%)	107 (100%)	0	100 100
13	SQ	119/121 (98%)	119 (100%)	0	100 100
14	SR	122/122 (100%)	120 (98%)	2 (2%)	62 84
15	SS	126/132 (96%)	126 (100%)	0	100 100
16	ST	113/115 (98%)	112 (99%)	1 (1%)	78 91
17	SU	94/107 (88%)	93 (99%)	1 (1%)	73 88

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
18	SV	67/67 (100%)	67 (100%)	0	100	100
19	SX	113/115 (98%)	113 (100%)	0	100	100
20	Sa	89/98 (91%)	89 (100%)	0	100	100
21	Sc	57/62 (92%)	57 (100%)	0	100	100
22	Sd	48/49 (98%)	48 (100%)	0	100	100
23	Sg	272/275 (99%)	272 (100%)	0	100	100
24	SC	188/225 (84%)	188 (100%)	0	100	100
25	SG	207/218 (95%)	206 (100%)	1 (0%)	88	95
26	SJ	161/168 (96%)	161 (100%)	0	100	100
27	SM	102/108 (94%)	99 (97%)	3 (3%)	42	74
28	SN	130/131 (99%)	130 (100%)	0	100	100
29	SO	110/119 (92%)	110 (100%)	0	100	100
30	SW	112/113 (99%)	112 (100%)	0	100	100
31	SY	113/115 (98%)	111 (98%)	2 (2%)	59	82
32	SZ	66/103 (64%)	66 (100%)	0	100	100
33	Sb	75/76 (99%)	75 (100%)	0	100	100
34	Se	47/48 (98%)	47 (100%)	0	100	100
35	Sf	60/140 (43%)	60 (100%)	0	100	100
36	CD	38/404 (9%)	38 (100%)	0	100	100
37	3G	70/277 (25%)	70 (100%)	0	100	100
All	All	4350/5365 (81%)	4336 (100%)	14 (0%)	92	96

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
17	SU	49	LYS
25	SG	119	LYS
31	SY	132	LYS
27	SM	121	LYS
31	SY	118	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
18	SV	35	ASN
24	SC	267	GLN
25	SG	13	GLN
11	SL	19	ASN
8	SH	165	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	S2	1701/1869 (91%)	413 (24%)	7 (0%)

5 of 413 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	S2	4	C
2	S2	14	C
2	S2	17	C
2	S2	25	A
2	S2	33	G

5 of 7 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
2	S2	543	C
2	S2	688	U
2	S2	1781	A
2	S2	1434	C
2	S2	541	U

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 26 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-38753. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 210




Y Index: 210




Z Index: 210


### 6.2.2 Raw map



X Index: 210



Y Index: 210




Z Index: 210


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 190




Y Index: 233




Z Index: 204


### 6.3.2 Raw map



X Index: 178



Y Index: 233




Z Index: 204

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1276  $\text{nm}^3$ ; this corresponds to an approximate mass of 1153 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of  $0.312 \text{ \AA}^{-1}$

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.17	3.79	3.22
Unmasked-calculated*	4.06	7.01	4.24

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.06 differs from the reported value 3.2 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-38753 and PDB model 8XXM. Per-residue inclusion information can be found in section [3](#) on page [11](#).


### 9.1 Map-model overlay [i](#)



X



Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 97% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ







The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9660	 0.4910
3G	 0.7940	 0.2310
CD	 0.9420	 0.4810
Ln	 0.9520	 0.4770
S2	 0.9910	 0.5100
SA	 0.9490	 0.5170
SB	 0.9530	 0.4950
SC	 0.9870	 0.5610
SD	 0.9650	 0.5160
SE	 0.9960	 0.5570
SF	 0.9610	 0.4660
SG	 0.9750	 0.4410
SH	 0.8950	 0.3940
SI	 0.9770	 0.4810
SJ	 0.9790	 0.5500
SK	 0.9710	 0.4630
SL	 0.9690	 0.5200
SM	 0.5070	 0.1740
SN	 0.9910	 0.5300
SO	 0.9390	 0.4900
SP	 0.9560	 0.4660
SQ	 0.9720	 0.5100
SR	 0.9430	 0.4710
SS	 0.9380	 0.4320
ST	 0.9860	 0.4930
SU	 0.9590	 0.4740
SV	 0.9840	 0.5250
SW	 0.9950	 0.5840
SX	 0.9960	 0.5600
SY	 0.9690	 0.5040
SZ	 0.9090	 0.3520
Sa	 0.9660	 0.5260
Sb	 0.9470	 0.4620
Sc	 0.9240	 0.4070
Sd	 0.9730	 0.5130



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
Se	 0.9620	 0.5090
Sf	 0.3840	 0.1460
Sg	 0.9410	 0.3830